

# AutoML: Evaluation

Background: Statistical Hypothesis Tests

Bernd Bischl   Frank Hutter   Lars Kotthoff  
Marius Lindauer   Joaquin Vanschoren

# Background: statistical hypothesis tests

- When we have a lot of data, we need to summarize it
  - ▶ But we already saw that summarization hides a lot of data
  - ▶ Ideally, we want to draw high-level conclusions (e.g., “A outperforms B on datasets of type X”)

# Background: statistical hypothesis tests

- When we have a lot of data, we need to summarize it
  - ▶ But we already saw that summarization hides a lot of data
  - ▶ Ideally, we want to draw high-level conclusions (e.g., “A outperforms B on datasets of type X”)
- Problem: we only have a finite number of observations
  - ▶ Can we attribute observed performance differences to chance?
  - ▶ Are we reasonably sure that a claim we make is reproducible?
  - ~> Statistical tests can help

# Statistical hypothesis testing

- 1 Define initial research hypothesis

# Statistical hypothesis testing

- 1 Define initial research hypothesis
- 2 Derive null  $H_0$  and alternative  $H_1$  hypothesis
  - ▶ Alternative hypothesis should be your research hypothesis

## First example: Courtroom Trial

- A prosecutor tries to prove the guilt of the defendant
- $H_0$ : The defendant is not guilty
  - ▶ Accepted for the moment  
(“not guilty as long as their guilt is not proven”)
- $H_1$ : The defendant is guilty
  - ▶ prosecutor hopes to support that

## First example: Courtroom Trial

- A prosecutor tries to prove the guilt of the defendant
- $H_0$ : The defendant is not guilty
  - ▶ Accepted for the moment  
(“not guilty as long as their guilt is not proven”)
- $H_1$ : The defendant is guilty
  - ▶ prosecutor hopes to support that

# First example: Courtroom Trial

- A prosecutor tries to prove the guilt of the defendant
- $H_0$ : The defendant is not guilty
  - ▶ Accepted for the moment  
(“not guilty as long as their guilt is not proven”)
- $H_1$ : The defendant is guilty
  - ▶ prosecutor hopes to support that

	Truly not guilty	Truly guilty
Found not guilty	Acquittal	Type II Error
Found guilty	Type I Error	Conviction

⇒ We want to minimize Type I error!



## Statistical hypothesis testing (cont'd)

- 1 Define initial research hypothesis
- 2 Derive null  $H_0$  and alternative  $H_1$  hypothesis
  - ▶ Alternative hypothesis should be your research hypothesis
- 3 Consider statistical assumptions
  - ▶ E.g., is your data Gaussian distributed?

# Statistical hypothesis testing (cont'd)

- ① Define initial research hypothesis
- ② Derive null  $H_0$  and alternative  $H_1$  hypothesis
  - ▶ Alternative hypothesis should be your research hypothesis
- ③ Consider statistical assumptions
  - ▶ E.g., is your data Gaussian distributed?
- ④ Decide test and test statistic  $T$ 
  - ▶ The correct test depends on your statistical assumptions.
  - ▶ Typically: if you use more assumptions, the test is more powerful (i.e., less Type-I error)

# Statistical hypothesis testing (cont'd)

- ① Define initial research hypothesis
- ② Derive null  $H_0$  and alternative  $H_1$  hypothesis
  - ▶ Alternative hypothesis should be your research hypothesis
- ③ Consider statistical assumptions
  - ▶ E.g., is your data Gaussian distributed?
- ④ Decide test and test statistic  $T$ 
  - ▶ The correct test depends on your statistical assumptions.
  - ▶ Typically: if you use more assumptions, the test is more powerful (i.e., less Type-I error)
- ⑤ Decide significance level  $\alpha$   
(i.e., acceptable Type-I error to reject null hypothesis)

# Statistical hypothesis testing (cont'd)

- ① Define initial research hypothesis
- ② Derive null  $H_0$  and alternative  $H_1$  hypothesis
  - ▶ Alternative hypothesis should be your research hypothesis
- ③ Consider statistical assumptions
  - ▶ E.g., is your data Gaussian distributed?
- ④ Decide test and test statistic  $T$ 
  - ▶ The correct test depends on your statistical assumptions.
  - ▶ Typically: if you use more assumptions, the test is more powerful (i.e., less Type-I error)
- ⑤ Decide significance level  $\alpha$   
(i.e., acceptable Type-I error to reject null hypothesis)
- ⑥ Compute observed  $t_{obs}$  of test statistic  $T$
- ⑦ Calculate  $p$ -value given  $t_{obs}$ 
  - ▶ i.e., probability under the null hypothesis of sampling a test statistic as extreme as observed (probability of Type-I error)

# Statistical hypothesis testing (cont'd)

- 1 Define initial research hypothesis
- 2 Derive null  $H_0$  and alternative  $H_1$  hypothesis
  - ▶ Alternative hypothesis should be your research hypothesis
- 3 Consider statistical assumptions
  - ▶ E.g., is your data Gaussian distributed?
- 4 Decide test and test statistic  $T$ 
  - ▶ The correct test depends on your statistical assumptions.
  - ▶ Typically: if you use more assumptions, the test is more powerful (i.e., less Type-I error)
- 5 Decide significance level  $\alpha$   
(i.e., acceptable Type-I error to reject null hypothesis)
- 6 Compute observed  $t_{obs}$  of test statistic  $T$
- 7 Calculate  $p$ -value given  $t_{obs}$ 
  - ▶ i.e., probability under the null hypothesis of sampling a test statistic as extreme as observed (probability of Type-I error)
- 8 If  $p < \alpha$ , reject null hypothesis in favor of alternative hypothesis
  - ▶ If  $p > \alpha$ , it doesn't tell you anything about the null hypothesis!

## Second example for a statistical test

- Claim: “the students in this course are more intelligent than average”

## Second example for a statistical test

- Claim: “the students in this course are more intelligent than average”
- Null Hypothesis  $H_0$ :  $\mu = 100$  ( $\mu$  is the population mean of this class)
- Alternative Hypothesis  $H_1$ :  $\mu > 100$  (one-sided hypothesis)

## Second example for a statistical test

- Claim: “the students in this course are more intelligent than average”
- Null Hypothesis  $H_0$ :  $\mu = 100$  ( $\mu$  is the population mean of this class)
- Alternative Hypothesis  $H_1$ :  $\mu > 100$  (one-sided hypothesis)
- IQ values are known to be normally distributed with  $X \sim \mathcal{N}(100, 15)$ 
  - ▶  $\rightarrow$  statistical assumption



## Second example for a statistical test

- Claim: “the students in this course are more intelligent than average”
- Null Hypothesis  $H_0$ :  $\mu = 100$  ( $\mu$  is the population mean of this class)
- Alternative Hypothesis  $H_1$ :  $\mu > 100$  (one-sided hypothesis)
- IQ values are known to be normally distributed with  $X \sim \mathcal{N}(100, 15)$ 
  - ▶  $\rightarrow$  statistical assumption
- Let's say we observed IQ values  $x_i$  of 9 students in the class:
  - ▶  $\{x_1, \dots, x_9\} = \{116, 128, 125, 119, 89, 99, 105, 116, 118\}$ .
  - ▶ The sample mean is  $\bar{x} = 112.8$
  - ▶ Does this data support the claim?

## Example continued


- Distribution of the test statistic

- ▶ Under  $H_0$ , we know that each  $x_i \sim \mathcal{N}(100, 15)$
- ▶ The test statistic that we measure is the sample mean  $\bar{x} = \frac{1}{9} \sum_{i=1}^9 x_i$


## Example continued

- Distribution of the test statistic

- ▶ Under  $H_0$ , we know that each  $x_i \sim \mathcal{N}(100, 15)$
- ▶ The test statistic that we measure is the sample mean  $\bar{x} = \frac{1}{9} \sum_{i=1}^9 x_i$
- ▶ Under  $H_0$ , the distribution of  $\bar{x}$  is  $\mathcal{N}(100, 15/\sqrt{9})$ 
  - Our observation  $\bar{x} = 112.8$  is quite extreme under that distribution




## General principle




- Compare the test statistic (here:  $\bar{x}$ ) to its sampling distribution under  $H_0$

# General principle




- Compare the test statistic (here:  $\bar{x}$ ) to its sampling distribution under  $H_0$
- **P-value:** probability  $p$  of observing values **at least as extreme as  $\bar{x}$**

## General principle



- Compare the test statistic (here:  $\bar{x}$ ) to its sampling distribution under  $H_0$
- **P-value:** probability  $p$  of observing values **at least as extreme as  $\bar{x}$**
- Compare  $p$  to pre-defined confidence level  $\alpha$  (usually  $\alpha = 0.05$ );  
if  $p < \alpha$ , **reject  $H_0$**

# General principle



- Compare the test statistic (here:  $\bar{x}$ ) to its sampling distribution under  $H_0$
- **P-value:** probability  $p$  of observing values **at least as extreme as  $\bar{x}$**
- Compare  $p$  to pre-defined confidence level  $\alpha$  (usually  $\alpha = 0.05$ );  
if  $p < \alpha$ , **reject  $H_0$**
- With  $\alpha = 0.01$ , would we reject  $H_0$  in this case?

## Summary of example

- We just used a so-called *Z-test*
- $H_0: \mu = \mu_0, H_1: \mu > \mu_0$
- Assumptions:  $X \sim \mathcal{N}(\mu, \sigma^2)$  , with known  $\mu$  and  $\sigma^2$



## Summary of example

- We just used a so-called *Z-test*
- $H_0: \mu = \mu_0, H_1: \mu > \mu_0$
- Assumptions:  $X \sim \mathcal{N}(\mu, \sigma^2)$  , with known  $\mu$  and  $\sigma^2$
- *Test statistic*: sample mean  $\bar{x}$ ; evaluate under  $\mathcal{N}(\mu = \mu_0, s = \sigma^2/\sqrt{n})$


## Summary of example

- We just used a so-called **Z-test**
- $H_0: \mu = \mu_0, H_1: \mu > \mu_0$
- Assumptions:  $X \sim \mathcal{N}(\mu, \sigma^2)$  , with known  $\mu$  and  $\sigma^2$
- **Test statistic**: sample mean  $\bar{x}$ ; evaluate under  $\mathcal{N}(\mu = \mu_0, s = \sigma^2/\sqrt{n})$
- Equivalent: compute the **Z-statistic**:  $Z = (\bar{x} - \mu_0)/s$  and evaluate cumulative distribution  $\Phi(Z)$  of  $Z$  under  $\mathcal{N}(0, 1)$

# Summary of example


- We just used a so-called *Z-test*
- $H_0: \mu = \mu_0, H_1: \mu > \mu_0$
- Assumptions:  $X \sim \mathcal{N}(\mu, \sigma^2)$  , with known  $\mu$  and  $\sigma^2$
- *Test statistic*: sample mean  $\bar{x}$ ; evaluate under  $\mathcal{N}(\mu = \mu_0, s = \sigma^2/\sqrt{n})$
- Equivalent: compute the *Z-statistic*:  $Z = (\bar{x} - \mu_0)/s$  and evaluate cumulative distribution  $\Phi(Z)$  of  $Z$  under  $\mathcal{N}(0, 1)$ 
  - ▶ There are standard tables to look up  $\Phi(Z)$  for different values of  $Z$
  - ▶ Nowadays, there are standard libraries to compute  $\Phi(Z)$

## Two-sided tests




- Similar to one-sided tests, but testing for extreme values in both tails
- Example Z-test: two-sided alternative hypothesis  $H_1: \mu \neq \mu_0$

## Two-sided tests



- Similar to one-sided tests, but testing for extreme values in both tails
- Example Z-test: two-sided alternative hypothesis  $H_1: \mu \neq \mu_0$
- Compute  $Z = (\bar{x} - \mu_0)/s$  as before
- Compute  $p$ -value as  $p = 2\Phi(Z)$ , to account for both tails

## Two-sided tests



- Similar to one-sided tests, but testing for extreme values in both tails
- Example Z-test: two-sided alternative hypothesis  $H_1: \mu \neq \mu_0$
- Compute  $Z = (\bar{x} - \mu_0)/s$  as before
- Compute  $p$ -value as  $p = 2\Phi(Z)$ , to account for both tails
- With  $\alpha = 0.01$ , would we reject  $H_0$  in this case?

# General points about statistical hypothesis tests

- What if  $p > \alpha$ ?
  - ▶ Failure to reject  $H_0$
  - ▶ We cannot conclude that we can accept  $H_0$ !

# General points about statistical hypothesis tests

- What if  $p > \alpha$ ?
  - ▶ Failure to reject  $H_0$
  - ▶ We cannot conclude that we can accept  $H_0$ !
- Beware (i): most tests make some assumptions
  - ▶ E.g.,  $Z$ -test and popular  $t$ -test assume **normality**
  - ▶ Our data is often far from normally-distributed
    - ↪ E.g., exponential runtime distributions of optimizers
    - ↪ E.g., distribution of fitting a neural network with different random seeds is not well studied



# General points about statistical hypothesis tests

- What if  $p > \alpha$ ?
  - ▶ Failure to reject  $H_0$
  - ▶ We cannot conclude that we can accept  $H_0$ !
- Beware (i): most tests make some assumptions
  - ▶ E.g.,  $Z$ -test and popular  $t$ -test assume **normality**
  - ▶ Our data is often far from normally-distributed
    - ↪ E.g., exponential runtime distributions of optimizers
    - ↪ E.g., distribution of fitting a neural network with different random seeds is not well studied
- Beware (ii): if you use cross-validation, observations are not independent (you cannot apply statistical tests that assume independence)