

Homework #6

Eric Tao
Math 123: Homework #6

March 5, 2023

Question 1. Let $L = D - W$ be the unnormalized graph Laplacian associated to a graph $\mathcal{G} = (V, W)$ on points $V = \{x_i\}_{i=1}^n$ with symmetric weight matrix W and diagonal degree matrix D . Let $\{C, \bar{C}\}$ be any partition of V , and let:

$$f_i^C = \begin{cases} -\sqrt{\text{Vol}(\bar{C})/\text{Vol}(C)} & \text{if } x_i \in C \\ -\sqrt{\text{Vol}(C)/\text{Vol}(\bar{C})} & \text{if } x_i \in \bar{C} \end{cases}$$

- (a) Prove that $\langle Df^C, \mathbb{1} \rangle = 0$.
- (b) Prove that $(f^C)^T Df^C = \text{Vol}(V)$.
- (c) Prove that $(f^C)^T Lf^C = \text{Vol}(V) \cdot \text{Ncut}(C, \bar{C})$.

Solution.

□

Question 2. Recall that one construction of the weight matrix for a graph on data points $\{x_i\}_{i=1}^n$ is to use the Gaussian kernel

$$W_{ij} = \begin{cases} \exp(-\|x_i - x_j\|_2^2 / \sigma^2) & i \neq j \\ 0 & i = j \end{cases}$$

for some choice of $\sigma > 0$.

- (a) What happens to the resulting Laplacian matrix as $\sigma \rightarrow 0$?
- (b) What happens to the resulting Laplacian matrix as $\sigma \rightarrow \infty$?

Solution.

□

Question 3.

Solution.

□