

Homework #3

Eric Tao
Math 237: Homework #3

March 30, 2024

Question 1. Show that the only function $f \in L^1(\mathbb{R})$ such that $f = f * f$ is $f = 0$ almost everywhere.

Solution. Let $f \in L^1(\mathbb{R})$. Suppose we have that $f = f * f$. We recall that the Fourier transform distributes pointwise over convolution, hence we have that:

$$\hat{f}(\zeta) = \widehat{f * f}(\zeta) = \hat{f}(\zeta)\hat{f}(\zeta)$$

Hence, we already have that at each ζ , $\hat{f} = 0, 1$, by solving the equation $\hat{f}(\zeta)^2 - \hat{f}(\zeta) = 0$.

From Lemma 9.2.3 in Heil, we have that \hat{f} is continuous. Moreover, from the Riemann-Lesbesgue lemma, since $f \in L^1$, we already know that $\hat{f} \in C_0(\mathbb{R})$. Hence, by continuity, since \hat{f} may only take on values 0, 1, and that it is continuous, $\hat{f} = 0$ everywhere.

Thus, by Corollary 9.2.12 in Heil, since $\hat{f} = 0$ everywhere, it is true almost everywhere of course, and hence $f = 0$ almost everywhere. □

Question 6. Suppose that $f \in AC(\mathbb{T})$, that is, 1 periodic and absolutely continuous on $[0, 1]$.

6.1)

Prove that $\hat{f}'(n) = 2\pi i n \hat{f}(n)$ for all $n \in \mathbb{Z}$. Conclude that $\lim_{|n| \rightarrow \infty} n \hat{f}(n) = 0$.

6.2)

Show that if $\int_0^1 f(x) dx = 0$, then:

$$\int_0^1 |f(x)|^2 dx \leq \frac{1}{4\pi^2} \int_0^1 |f'(x)|^2 dx$$

Solution. 6.1)

First, we recall that by Corollary 6.1.5 in Heil, that since f is absolutely continuous on $[0, 1]$, we have that f is differentiable almost everywhere, and $f' \in L^1[0, 1]$. Since f' is at least L^1 , we may look at its Fourier transform:

$$\hat{f}'(n) = \int_0^1 f'(x) \exp(-2\pi i n x) dx$$

We notice that for a fixed value of $n \in \mathbb{Z}$, that we may compute the derivative of $g = \exp(-2\pi i n x)$ as $g' = -2\pi i n \exp(-2\pi i n x)$. Hence, g is differentiable everywhere on $[0, 1]$, and of course, since $|-2\pi i n \exp(-2\pi i n x)| = 2\pi n$ always, it is bounded. Hence, by Lemma 5.2.5, g is Lipschitz on $[0, 1]$. Thus, since Lipschitz functions are absolutely continuous (6.1.3), we may apply integration by parts on f, g . We have then that:

$$\int_0^1 f'(x) \exp(-2\pi i n x) dx = f(1) \exp(-2\pi i n) - f(0) \exp(0) - \int_0^1 f(x) [-2\pi i n \exp(-2\pi i n x)] dx$$

Now, because f is 1-periodic, we have that $f(0) = f(1)$. Further, $\exp(-2\pi in) = \exp(0)$, of course, since $2\pi n$ is a multiple of 2π for any n . Hence, the first terms vanish.

We are left then with, after using the linearity of the integral:

$$2\pi in \int_0^1 f(x) \exp(-2\pi inx) dx$$

However, we recognize the integral as exactly the Fourier transform of f at n . Hence, we have our result, that:

$$\hat{f}'(n) = 2\pi in \hat{f}(n)$$

Again, by the Riemann-Lebesgue lemma, we have that since $f' \in L^1$, that $\lim_{|n| \rightarrow \infty} \hat{f}'(n) = 0$. But, from our result, we have that:

$$\lim_{|n| \rightarrow \infty} 2\pi in \hat{f}(n) = \lim_{|n| \rightarrow \infty} \hat{f}'(n) = 0 \implies \lim_{|n| \rightarrow \infty} n \hat{f}(n) = 0$$

6.2)

First, we use the result from Heil Problem 9.3.24 (b) that the Plancherel equality holds for $f \in L^1(\mathbb{T})$. and prove that later.

Then, since $f' \in L^1(\mathbb{T})$, we have that:

$$\sum_{n \in \mathbb{Z}} |\hat{f}'(n)|^2 = \|f'\|_2^2 = \int_0^1 |f'(x)|^2 dx$$

Now, from 6.1, we have that $\hat{f}'(n) = 2\pi in \hat{f}(n)$ for each n . Hence, we have that:

$$\sum_{n \in \mathbb{Z}} |\hat{f}'(n)|^2 = \sum_{n \in \mathbb{Z}} 4\pi^2 n^2 |\hat{f}(n)|^2 = 4\pi^2 \sum_{n \in \mathbb{Z}} n^2 |\hat{f}(n)|^2$$

Now, we notice that since $\int_0^1 f(x) dx = 0$, that evidently, $\hat{f}(0) = 0$, as $\hat{f}(0) = \int_0^1 f(x) \exp(-2\pi i 0 x) dx = \int_0^1 f(x) dx = 0$.

Because this is 0, and for all $n \in \mathbb{Z}, n \neq 0$, we have that $|\hat{f}(n)| \leq |n \hat{f}(n)|$, we can conclude that:

$$4\pi^2 \sum_{n \in \mathbb{Z}} n^2 |\hat{f}(n)|^2 \geq 4\pi^2 \sum_{n \in \mathbb{Z}} |\hat{f}(n)|^2$$

Finally, applying Plancherel's equality again for f , as $f \in L^1(\mathbb{T})$, we see that:

$$4\pi^2 \sum_{n \in \mathbb{Z}} |\hat{f}(n)|^2 = 4\pi^2 \|f\|_2^2 = 4\pi^2 \int_0^1 |f(x)|^2 dx$$

And rewriting all of these inequalities together, we have that:

$$4\pi^2 \int_0^1 |f(x)|^2 \leq \int_0^1 |f'(x)|^2 dx \implies \int_0^1 |f(x)|^2 \leq \frac{1}{4\pi^2} \int_0^1 |f'(x)|^2 dx$$

as desired.

Now, we return to proving 9.3.24.

First, we wish to show that if $f \in L^1(\mathbb{T})$, and $\hat{f} \in L^2(\mathbb{Z})$, then $f \in L^2(\mathbb{T})$.

Next, we want to show that the Plancherel equality holds, either both sides being finite or infinite. If $f \in L^2 \cap L^1$, then we're fine, by Corollary 9.3.14.

Then, we assume $f \notin L^2(\mathbb{T})$ but $f \in L^1(\mathbb{T})$. Then, by the previous result, we have that $\hat{f} \notin L^2(\mathbb{Z})$. Hence, we have that both sides of the Plancherel equality are infinite, and we are done. \square

Question 12. Fix a $g \in L^2(\mathbb{R})$. Let $a \in \mathbb{R}$, and define the operator T_a on g that sends $T_a(g(x)) \mapsto g(x-a)$. Prove that the family $\{T_a g\}_{a \in \mathbb{R}}$ is complete in L^2 if and only if $\hat{g}(\zeta) \neq 0$ almost everywhere.

Solution. \square

Question 14. Let $p(x) = \chi_{[0,1)}(x)$, $h(x) = \chi_{[0,1/2)}(x) - \chi_{[1/2,1)}(x)$. Let $j, k \in \mathbb{Z}$, and define $I_{jk} = [2^{-j}k, 2^{-j}(k+1))$. Further define the following functions:

$$\begin{cases} p_{jk} = 2^{j/2}p(2^j x - k) \\ h_{jk} = 2^{j/2}h(2^j x - k) \end{cases}$$

14.1)

Prove that $\{h_{jk}\}$ is an orthonormal sequence in L^2 .

14.2)

For each fixed $j \in \mathbb{Z}$, prove that $\{p_{jk}\}_{k \in \mathbb{Z}}$ is an orthonormal sequence in L^2 .

14.3)

Fix a $j \in \mathbb{Z}$. Let g_j be any step function, constant on each interval I_{jk} for $k \in \mathbb{Z}$. Show that we may express $g_j(x) = g_{j-1}(x) + r_{j-1}(x)$, where

$$r_{j-1}(x) = \sum_{k \in \mathbb{Z}} a_{j-1}(k) h_{j-1,k}(x)$$

for some coefficients $a_{j-1}(k)$ and some step function $g_{j-1}(x)$, constant on intervals $I_{j-1,k}$.

14.4)

Fix a $J \geq 0$. Consider the set:

$$\{p_{Jk} : 0 \leq k \leq 2^J - 1\} \cup \{h_{j,k} : j \geq J, 0 \leq k \leq 2^j - 1\}$$

Prove that this set is an orthonormal sequence in $L^2[0, 1]$.

14.5)

For $f \in L^2[0, 1]$, and a fixed $J \geq 0$, show that we may find g_j step functions for $j \geq J$, such that they are constant on each $I_{j,k}$, and that g_j approximates f in the L^2 norm.

Use this result and the result of 14.4 to show that the set in 14.4 is an orthonormal basis for $L^2[0, 1]$.

Solution. \square

Question 16. Let ϕ be a non-0 function in $L^2(\mathbb{R})$. For any $f \in L^2(\mathbb{R})$, define $V_\phi f$ via:

$$V_\phi(f)(x, \zeta) = \int_{\mathbb{R}} f(t) \overline{\phi(t-x)} \exp(-2\pi i t \zeta) dt$$

For $a, b \in \mathbb{R}$, let T_a be the translation operator that sends $T_a(f(x)) \mapsto f(x-a)$ and let M_b the modulation operator that sends $M_b(f(x)) \mapsto \exp(2\pi b x) f(x)$.

16.1)

Prove that for each $f \in L^2$, $V_\phi f$ is uniformly continuous on \mathbb{R}^2 , and that $\lim_{|(x, \zeta)| \rightarrow \infty} V_\phi f = 0$.

16.2)

Recall that the Schwarz space $\mathcal{S}(\mathbb{R})$ is defined as:

$$\mathcal{S}(\mathbb{R}) = \{f \in C^\infty(\mathbb{R}) : x^m f^{(n)}(x) \in L^\infty(\mathbb{R}) \text{ for all } m, n \geq 0\}$$

Prove that if $f \in \mathcal{S}(\mathbb{R})$, then $V_\phi \in \mathcal{S}(\mathbb{R}^2)$.

16.3)

Prove that V_ϕ acts as an isometry from $L^2(\mathbb{R})$ to $L^2(\mathbb{R}^2)$, and that $\|V_\phi f\|_{L^2(\mathbb{R}^2)} = \|\phi\|_{L^2(\mathbb{R})} \|f\|_{L^2(\mathbb{R})}$ for every $f \in L^2$.

16.4)

Show that the operator V_ϕ^* defined by:

$$V_\phi^* F(t) = \|\phi\|_2^{-2} \iint_{\mathbb{R}^2} F(x, \zeta) \exp(2\pi i \zeta t) \phi(t - x) dx d\zeta$$

takes $L^2(\mathbb{R}^2)$ to $L^2(\mathbb{R})$, and that for each $f \in L^2(\mathbb{R})$, we can make sense of the following inversion formula:

$$f(t) = \|\phi\|_2^{-2} \iint_{\mathbb{R}^2} V_\phi f(x, \zeta) \exp(2\pi i \zeta t) \phi(t - x) dx d\zeta$$

Solution.

□