The Sudbury Neutrino Observatory

Yutong Du

Physics H190 Research Presentation

October 2, 2024

What is SNO?

- From SNO themselves: "The goal of the observatory is to detect and study neutrinos emitted by the Sun and other celestial objects."
- A Heavy Water Cherenkov Detector
- Built in the Creighton Mine in Sudbury Ontario, 2km underground
- Largest underground cavity excavated at 2km depth

The Solar Neutrino Problem (SNP)

- Previous neutrino observation data was only $\frac{1}{3}$ of what was predicted, but the instrument was only sensitive to electron neutrinos ν_e .
- Hypothetical Solution: The neutrinos change flavor over the course of their journey from the Sun to the detector
- SNO's goal: verify this hypothesis

Rough Timeline

- 1985: Proposed solution to resolve the SNP via a Heavy-Water Cherenkov Detector
- 1989: US and Britain joins in on the project
- 1990: Construction begins
- November 1999: Construction completes, data collection (Phase I) begins.
 - May 2001: Phase II (NaCl → higher sensitivity)
 - February 2004: Phase III (Neutral Current Detector)
- November 2006: Data collection ends
- October 2015: Nobel Prize in Physics

Specs

- Spherical Chamber 12 m across
- 9456 Hamamatsu R4108 photomulplier tubes
- 1000 tons of Heavy Water (D₂O)
- 7000 tons of ultra-pure normal water





Detection Methods

Neutral Current (NC)

$$\nu_{\mathsf{x}} + \mathsf{d} \rightarrow \mathsf{p} + \mathsf{n} + \nu_{\mathsf{x}}$$

selects for all neutrinos equally

Charged Current (CC)

$$\nu_e + d \rightarrow p + p + e^-$$


Only selects for electron neutrinos

Elastic Scattering (ES)


$$\nu_{\rm X} + {\rm e}^- \rightarrow \nu_{\rm X} + {\rm e}^-$$

Selects for all neutrinos; ν_e selected 6x sensitivity

Feynman Diagrams



Charged current reaction, electron neutrinos only.



Neutral current reaction. all neutrinos.

Elastic scattering with any neutrino.

Results

- ullet Total number of neutrinos detected matches theory o solves SNP
- Proves that Neutrinos have mass due to oscillations.

$$P_{\alpha \to \beta} = \sin^2(2\theta) \sin^2\left(\frac{\Delta m^2 L}{4E}\right)$$

So $\Delta m > 0 \rightarrow$ neutrinos have mass!

• Data for solar mixing angle θ_{sol}

SNP Resolution




Figure: Events vs. effective kinetic energy

Evidence of Δm




Figure: Mixing angle $\tan^2 \theta_{12}$ vs. Δm_{21}^2

SNOLAB and Beyond

Since SNO, the mine was expanded to accommodate for more experiments, creating SNOLAB.

- SNO+: World's largest scintillator experiment
- SuperCDMS: Dark matter searches using Si and Ge crystals
- REPAIR: Effect of radiation on living organisms




Figure: SNO+ detector

References



Aharmim et al.

Combined analysis of all the phases of solar neutrino data from the sudbury neutrino observatory.

Physical Review C 50th Anniversay Milestones, 2013.



Linda Vu.

Historic sudbury neutrino observatory data, carried by esnet, lives on at nersc, February 2010.



The sudbury neutrino observatory, April 2023.



Boris Kayser.

Neutrino physics, August 2013.



Jelly et al.

The sudbury neutrino observatory.

Advance, November 2009.