

# Introduction to Mathematics of Quantum Computing

Erick Jesús Ríos González

March 13, 2024

Quantum mechanics describes states of systems at microscopic scales, which generally means that only speaking in terms of probabilities is allowed. The state in a quantum system can be described as an element of a vector space. It is possible that the state of the system can be written as a linear combination of a basis of the vector space: **superposition**. The observables of a quantum system are those physical quantities that in principle can be determined simultaneously and with a precision constraint: **Heisenberg's uncertainty principle**. In other words, they can be measurable.

Measurements on objects prepared "in the same way" are distributed with a relative frequency around a mean value.

$$\text{relative frequency} = \frac{\text{Number of measurements with result } a}{\text{Total number of measurements}}$$

$$\text{mean value} = \sum_{a \in \text{Measurements}} a \times (\text{relative frequency of measurements with result } a)$$

## 1 Hilbert Spaces

There are initial preparations whose state can be characterized with a vector in a Hilbert space (vector space).

1. States that can be characterized with a vector are called pure states.
2. Otherwise, we speak of "mixed states".

### Definition 1.1: Hilbert Space

A Hilbert Space  $\mathbb{H}$  is a complete complex vector space, that is,

$$\Psi, \Phi \in \mathbb{H} \text{ and } a, b \in \mathbb{C} \implies a\Psi + b\Phi \in \mathbb{H} \quad (1)$$

with a positive-definite scalar product:

$$\langle \cdot | \cdot \rangle : \mathbb{H} \times \mathbb{H} \rightarrow \mathbb{C} \quad (2)$$

such that  $\forall \Psi, \Phi, \Psi_1, \Psi_2 \in \mathbb{H}$  and  $a, b \in \mathbb{C}$

$$\langle \Psi | \Phi \rangle = \langle \Psi | \Phi \rangle$$