

KS141203 MATEMATIKA DISKRIT (*DISCRETE MATHEMATICS*)

Number Theory: Integers, Division, Prime Number

Ahmad Muklason, Ph.D.

Outline



1. Integer and Division
2. Primes
3. GCD (Great Common Divisor)
4. LCM (least Common Multiple)



Division

- **Definition:** if a and b are integers ($a \neq 0$), a divides b if $\exists c$ such that $b = ac$.
- When a divides b , we say that a is a factor of b and that b is a multiple of a .
- **Notation:**
 - $a|b$: a divides b (a habis membagi b ; b habis dibagi a)
 - $a \nmid b$: a does not divide b (a tidak habis membagi b ; b tidak habis dibagi a)
- **Example:**
 - $3|7$? $3|12$?
 - $3 \nmid 7$ since $7/3$ is not an integer
 - $3|12$ because $12/3 = 4$



Theorem 1

Let a , b , and c be integers, then:

- If a/b and a/c , then $a|(b+c)$
- If a/b , then a/bc for all integers c
- If a/b and b/c , then a/c

Proof:

If a/b and a/c , then $a|(b+c)$

- $b = ma$ and $c = na$
- $b + c = ma + na = (m+n)a$
- $b + c = (m+n)a$
- So, $a | (b+c)$

If a/b , then a/bc for all integers c

- $b = ma$, $bc = (ma)c = (mc)a$
- $bc = (mc)a$
- So, $a | bc$

If a/b and b/c , then a/c

- $b = ma$, $c = pb = p(ma) = (pm)a$
- $c = (pm)a$
- So, $a | c$



Corollary 1

$$a/b \text{ and } a/c \rightarrow a/mb + nc$$

Proof:

$$b = pa$$

$$c = qa$$

$$mb = (mp)a$$

$$nc = (nq)a$$

$$mb + nc = (mp + nq)a$$

$$\text{So, } a \mid mb + nc$$

***Corollary:** proposisi yang merupakan akibat langsung dari teorema yang dibuktikan

Division Algorithm

- **Theorem 2:** Let a be an integer and d a positive integer. Then there exist unique integers q and r , with $0 \leq r < d$, such that $a = dq + r$.
- **Definition**
 - $q = a \text{ div } d$; q = quotient (*hasil bagi*), d = divisor (*pembagi*), a = dividend (*yang dibagi*)
 - $r = a \text{ mod } d$; r = remainder (*sisanya*)



Division Algorithm Examples

- What are the quotient and remainder when 101 is divided by 11?
 - $101 = 11 \cdot 9 + 2$
 - The quotient is: $9 = 101 \text{ div } 11$
 - The remainder is: $2 = 101 \text{ mod } 11$
- What are the quotient and remainder when -11 is divided by 3?
 - $-11 = 3(-4) + 1$
 - The quotient is: $-4 = -11 \text{ div } 3$
 - The remainder is: $1 = -11 \text{ mod } 3$
 - Note: the remainder can't be negative
 - $-11 = 3(-3) - 2 \rightarrow r = -2$ doesn't satisfy $0 \leq r < 3$



Modular Arithmetic

- **Definition:** If a and b are integers and m is positive integer, then a is **congruent** to b modulo m if m divides $a - b$.
- **Notation:**
 - $a \equiv b \pmod{m}$; a is congruent to b modulo m
 - $a \not\equiv b \pmod{m}$; a and b are not congruent to modulo m
- **Theorem 3:**
 - $a \equiv b \pmod{m}$ iff $a \bmod m = b \bmod m$.
- **Example:**
 - $17 \equiv 12 \pmod{5}$, $17 \bmod 5 = 2$, $12 \bmod 5 = 2$



Modular Arithmetic

Theorem 4:

- Let m be a positive integer, $a \equiv b \pmod{m}$ iff $\exists k$ such that $a = b + km$.

Proof:

- If $a \equiv b \pmod{m}$, then $m \mid (a - b)$.
- This means that $\exists k$ such that $a - b = km$, so that $a = b + km$.
- Conversely, if $\exists k$ such that $a = b + km$, then $km = a - b$.
- Hence, m divides $a - b$, so that $a \equiv b \pmod{m}$



Modular Arithmetic

Theorem 5:

- Let m be a positive integer. If $a \equiv b \pmod{m}$ and $c \equiv d \pmod{m}$, then
 - $a + c \equiv b + d \pmod{m}$ and $ac \equiv bd \pmod{m}$

Proof:

- Because $a \equiv b \pmod{m}$ and $c \equiv d \pmod{m}$, there are integers s and t with $b = a + sm$ and $d = c + tm$. Hence:
 - $b + d = (a + sm) + (c + tm) = (a + c) + m(s + t)$
 - $bd = (a + sm)(c + tm) = ac + m(at + cs + stm)$
- Hence, $a + c \equiv b + d \pmod{m}$ and $ac \equiv bd \pmod{m}$



Corollary 2

Let m be a positive integer, a and b be integers. Then:

- $(a + b) \bmod m = ((a \bmod m) + (b \bmod m)) \bmod m$
- $ab \bmod m = ((a \bmod m)(b \bmod m)) \bmod m$



Exercise

What are the quotient and remainder when

- | | |
|------------------------------|--------------|
| 1. -111 is divided by 11 | 1. $-11, 10$ |
| 2. 789 is divided by 23 | 2. $34, 7$ |
| 3. 0 is divided by 19 | 3. $0, 0$ |
| 4. 3 is divided by 5 | 4. $0, 3$ |
| 5. -1 is divided by 3 | 5. $-1, 2$ |

What time does a 12-hour clock read

- | | |
|------------------------------------|----------|
| 1. 80 hours after it reads 11:00? | 1. 7:00 |
| 2. 40 hours before it reads 12:00? | 2. 8:00 |
| 3. 100 hours after it reads 6:00? | 3. 10:00 |

Caesar Cipher

Alphabet to number: $a \sim 0, b \sim 1, \dots, z \sim 25$.

Encryption: $f(p) = (p + k) \bmod 26$.

Decryption: $f^{-1}(p) = (p - k) \bmod 26$.

- Caesar used $k = 3$.

This is called a substitution cipher

- You are substituting one letter with another

Caesar Cipher Example

Encrypt “go cavaliers”

- Translate to numbers: g = 6, o = 14, etc.
 - Full sequence: 6, 14, 2, 0, 21, 0, 11, 8, 4, 17, 18
- Apply the cipher to each number: $f(6) = 9$, $f(14) = 17$, etc.
 - Full sequence: 9, 17, 5, 3, 24, 3, 14, 11, 7, 20, 21
- Convert the numbers back to letters 9 = j, 17 = r, etc.
 - Full sequence: jr wfdydolhuv

Decrypt “jr fdydolhuv”

- Translate to numbers: j = 9, r = 17, etc.
 - Full sequence: 9, 17, 5, 3, 24, 3, 14, 11, 7, 20, 21
- Apply the cipher to each number: $f^{-1}(9) = 6$, $f^{-1}(17) = 14$, etc.
 - Full sequence: 6, 14, 2, 0, 21, 0, 11, 8, 4, 17, 18
- Convert the numbers back to letters 6 = g, 14 = o, etc.
 - Full sequence: go cavaliers

Caesar Cipher Example

Encrypt “MEET YOU IN THE PARK”

- Translate to numbers:
 - Full sequence: 12, 4, 4, 19, 24, 14, 20, 8, 13, 19, 7, 4, 15, 0, 17, 10
- Apply the cipher to each number:
 - Full sequence: 15, 7, 7, 22, 1, 7, 23, 11, 16, 22, 10, 7, 18, 3, 20, 13
- Convert the numbers back to letters:
 - Full sequence: PHHW BRX LQ WKH SDUN



Caesar Cipher Example

What letter replaces the letter K when the function $f(p) = (7p + 3) \bmod 26$ is used for encryption?

Solution:

First, note that 10 represents K , then using the encryption function specified, it follows that $f(10) = (7 \cdot 10 + 3) \bmod 26 = 21$

Because 21 represents V , K is replaced by V in the encrypted message.

Prime Numbers

Definition: A positive integer p is **prime** if the only positive factors of p are 1 and p

- If there are other factors, it is **composite**
- Note that **1 is not prime!**
- It's not composite either – it's in its own class

Definition: An integer n is **composite** if and only if there exists an integer a such that a/n and $1 < a < n$

Fundamental theorem of arithmetic



Every positive integer greater than 1 can be **uniquely written as a prime** or **as the product of two or more primes** where the prime factors are written in order of non-decreasing size

Examples

- $100 = 2 * 2 * 5 * 5$
- $182 = 2 * 7 * 13$
- $29820 = 2 * 2 * 3 * 5 * 7 * 71$



Composite Factors

If n is a composite integer, then n has a prime divisor less than or equal to the square root of n



Showing a number is prime

Show that 113 is prime

Solution

- The only prime factors less than $\sqrt{113} = 10.63$ are 2, 3, 5, and 7
- Neither of these divide 113 evenly
- Thus, by the fundamental theorem of arithmetic, 113 must be prime

Showing a number is composite

Show that 899 is prime

Solution

- Divide 899 by successively larger primes (up to $\sqrt{899} = 29.98$), starting with 2
- We find that 29 and 31 divide 899

Primes are infinite

Theorem (by Euclid): There are infinitely many prime numbers

Proof by contradiction

Assume there are a finite number of primes

List them as follows: p_1, p_2, \dots, p_n .

Consider the number $q = p_1 p_2 \dots p_n + 1$

- This number is not divisible by any of the listed primes
 - If we divided p_i into q , there would result a remainder of 1
- We must conclude that q is a prime number, not among the primes listed above
 - This contradicts our assumption that all primes are in the list p_1, p_2, \dots, p_n .



The prime number theorem

The ratio of the number of primes not exceeding x and $x/\ln(x)$ approaches 1 as x grows without bound

- Rephrased: the number of prime numbers less than x is approximately $x/\ln(x)$
- Rephrased: the chance of an number x being a prime number is $1 / \ln(x)$

Consider 200 digit prime numbers

- $\ln(10^{200}) \approx 460$
- The chance of a 200 digit number being prime is $1/460$
- If we only choose odd numbers, the chance is $2/460 = 1/230$



Greatest common divisor

The greatest common divisor of two integers a and b is the largest integer d such that $d|a$ and $d|b$

- Denoted by $\gcd(a, b)$

Examples

- $\gcd(24, 36) = 12$
- $\gcd(17, 22) = 1$
- $\gcd(100, 17) = 1$

Relative primes

Two numbers are relatively prime if they don't have any common factors (other than 1)

- Rephrased: a and b are relatively prime if
$$\gcd(a, b) = 1$$

$\gcd(25, 39) = 1$, so 25 and 39 are relatively prime

Pairwise relative prime

A set of integers a_1, a_2, \dots, a_n are pairwise relatively prime if, for all pairs of numbers, they are relatively prime

- Formally: The integers a_1, a_2, \dots, a_n are pairwise relatively prime if $\gcd(a_i, a_j) = 1$ whenever $1 \leq i < j \leq n$.

Example: are 10, 17, and 21 pairwise relatively prime?

- $\gcd(10, 17) = 1$, $\gcd(17, 21) = 1$, and $\gcd(21, 10) = 1$
- Thus, they are pairwise relatively prime

Example: are 10, 19, and 24 pairwise relatively prime?

- Since $\gcd(10, 24) \neq 1$, they are not



More on gcd's

Given two numbers a and b , rewrite them as:

$$a = p_1^{a_1} p_2^{a_2} \dots p_n^{a_n}, b = p_1^{b_1} p_2^{b_2} \dots p_n^{b_n}$$

Example: gcd (120, 500)

- $120 = 2^3 * 3 * 5 = 2^3 * 3^1 * 5^1$
- $500 = 2^2 * 5^3 = 2^2 * 3^0 * 5^3$

Then compute the gcd by the following formula:

$$\text{gcd}(a, b) = p_1^{\min(a_1, b_1)} p_2^{\min(a_2, b_2)} \dots p_n^{\min(a_n, b_n)}$$

- Example: $\text{gcd}(120, 500) = 2^{\min(3, 2)} 3^{\min(1, 0)} 5^{\min(1, 3)} = 2^2 3^0 5^1 = 20$

Least common multiple

The least common multiple of the positive integers a and b is the smallest positive integer that is divisible by both a and b .

- Denoted by $\text{lcm}(a, b)$

$$\text{lcm}(a, b) = p_1^{\max(a_1, b_1)} p_2^{\max(a_2, b_2)} \dots p_n^{\max(a_n, b_n)}$$

Example: $\text{lcm}(10, 25) = 50$

What is $\text{lcm}(95256, 432)$?

- $95256 = 2^3 3^5 7^2$; $432 = 2^4 3^3$
- $\text{lcm}(2^3 3^5 7^2, 2^4 3^3) = 2^{\max(3, 4)} 3^{\max(5, 3)} 7^{\max(2, 0)} = 2^4 3^5 7^2 = 190512$

lcm and gcd theorem

Let a and b be positive integers. Then

$$a * b = \gcd(a, b) * \text{lcm}(a, b)$$

Example: $\gcd(10, 25) = 5$, $\text{lcm}(10, 25) = 50$

- $10 * 25 = 5 * 50$

Example: $\gcd(95256, 432) = 216$, $\text{lcm}(95256, 432) = 190512$

- $95256 * 432 = 216 * 190512$