

Fibrin glue system for adjuvant brachytherapy of brain tumors with ¹⁸⁸Re and ¹⁸⁶Re-labeled microspheres.

Authors: Hafeli U.O., Pauer G.J., Unnithan J., Prayson R.A.

Publication Date: 2007

Abstract:

Brain tumors such as glioblastoma reappear in their original location in almost 50% of cases. To prevent this recurrence, we developed a radiopharmaceutical system that consists of a gel applied immediately after surgical resection of a brain tumor to deliver local radiation booster doses. The gel, which strongly adheres to tissue in the treatment area, consists of fibrin glue containing the beta-emitters rhenium-188 and rhenium-186 in microsphere-bound form. Such microspheres can be prepared by short (2 h or less) neutron activation even in low neutron flux reactors, yielding a mixture of the two beta-emitters rhenium-188 ($E_{\text{max}} = 2.1$ MeV, half life = 17 h) and rhenium-186 ($E_{\text{max}} = 1.1$ MeV, half life = 90.6 h). The dosimetry of this rhenium-188/rhenium-186 fibrin glue system was determined using gafchromic film measurements. The treatment efficacy of the radioactive fibrin glue was measured in a 9L-glioblastoma rat model. All animals receiving the non-radioactive fibrin glue died within 17 +/- 3 days, whereas 60% of the treated animals survived 36 days, the final length of the experiment. Control animals that were treated with the same amount of radioactive fibrin glue, but had not received a previous tumor cell injection, showed no toxic effects over one year. The beta-radiation emitting rhenium-188/rhenium-186-based gel thus provides an effective method of delivering high doses of local radiation to tumor tissue, particularly to wet areas where high adhesive strength and long-term radiation (with or without drug) delivery are needed. © 2006 Elsevier B.V. All rights reserved.