Antitumor effect of fibrin glue containing temozolomide against malignant glioma.

Authors: Anai S., Hide T., Takezaki T., Kuroda J.-I., Shinojima N., Makino K., Nakamura H., Yano

S., Kuratsu J.-I.

Publication Date: 2014

Abstract:

Temozolomide (TMZ), used to treat glioblastoma and malignant glioma, induces autophagy,

apoptosis and senescence in cancer cells. We investigated fibrin glue (FG) as a drug delivery

system for the local administration of high-concentration TMZ aimed at preventing glioma

recurrence. Our high-power liquid chromatography studies indicated that FG containing TMZ

(TMZ-FG) manifested a sustained drug release potential. We prepared a subcutaneous tumor model

by injecting groups of mice with three malignant glioma cell lines and examined the antitumor effect

of TMZ-FG. We estimated the tumor volume and performed immunostaining and immunoblotting

using antibodies to Ki-67, cleaved caspase 3, LC3 and p16. When FG sheets containing TMZ

(TMZ-FGS) were inserted beneath the tumors, their growth was significantly suppressed. In mice

treated with peroral TMZ plus TMZ-FGS the tumors tended to be smaller than in mice whose tumors

were treated with TMZ-FGS or peroral TMZ alone. The TMZ-FGS induced autophagy, apoptosis

and senescence in subcutaneous glioma tumor cells. To assess the safety of TMZ-FG for normal

brain, we placed it directly on the brain of living mice and stained tissue sections obtained in the

acute and chronic phase immunohistochemically. In both phases, TMZ-FG failed to severely

damage normal brain tissue. TMZ-FG may represent a safe new drug delivery system with

sustained drug release potential to treat malignant glioma. © 2014 The Authors.