

L-PRP/L-PRF in esthetic plastic surgery, regenerative medicine of the skin and chronic wounds.

Authors: Cieslik-Bieleck A., Choukroun J., Odin G., Dohan Ehrenfest D.M.

Publication Date: 2012

Abstract:

The use of platelet concentrates for topical use is of particular interest for the promotion of skin wound healing. Fibrin-based surgical adjuvants are indeed widely used in plastic surgery since many years in order to improve scar healing and wound closure. However, the addition of platelets and their associated growth factors opened a new range of possibilities, particularly for the treatment of chronic skin ulcers and other applications of regenerative medicine on the covering tissues. In the 4 families of platelet concentrates available, 2 families were particularly used and tested in this clinical field: L-PRP (Leukocyte- and Platelet-rich Plasma) and L-PRF (Leukocyte- and Platelet-Rich Fibrin). These 2 families have in common the presence of significant concentrations of leukocytes, and these cells are important in the local cleaning and immune regulation of the wound healing process. The main difference between them is the fibrin architecture, and this parameter considerably influences the healing potential and the therapeutical protocol associated to each platelet concentrate technology. In this article, we describe the historical evolutions of these techniques from the fibrin glues to the current L-PRP and L-PRF, and discuss the important functions of the platelet growth factors, the leukocyte content and the fibrin architecture in order to optimize the numerous potential applications of these products in regenerative medicine of the skin. Many outstanding perspectives are appearing in this field and require further research. © 2012 Bentham Science Publishers.