

Treatment of experimental peri-implantitis using autogenous bone grafts and platelet-enriched fibrin glue in dogs.

Authors: You T.-M., Choi B.-H., Zhu S.-J., Jung J.-H., Lee S.-H., Huh J.-Y., Lee H.-J., Li J.

Publication Date: 2007

Abstract:

Objective: The purpose of this study was to evaluate the effects of autogenous bone grafts and platelet-enriched fibrin glue in the treatment of peri-implantitis. **Study design:** Thirty-six screw-type commercially pure titanium implants with rough acid-etched surfaces were inserted into 6 mongrel dogs 3 months after extraction of mandibular premolars. After 3 months of healing, peri-implantitis was induced by placing gauze and wire around the implants. Once peri-implantitis was created, surgical treatments involving a combination of autogenous bone grafts and platelet-enriched fibrin glue, autogenous bone grafts alone, or a conventional flap procedure only (control) were carried out. Six months later, biopsies of the implant sites were taken and prepared for ground sectioning and analysis. **Results:** The amount of reosseointegration was significantly higher in peri-implantitis defects treated with combined autogenous bone grafts and platelet-enriched fibrin glue as compared with the other 2 treatment procedures. A mean bone-to-implant contact of 50.1% was obtained in the peri-implantitis lesions treated with combined autogenous bone grafts and platelet-enriched fibrin glue. The corresponding values for the autogenous bone grafts and control groups were 19.3% and 6.5%, respectively. **Conclusion:** The present study demonstrates that surgical treatment involving the combined use of autogenous bone grafts and platelet-enriched fibrin glue might effectively promote reosseointegration in lesions resulting from peri-implantitis. © 2007 Mosby, Inc. All rights reserved.