

HWs

Eric Liu

# CONTENTS

## CHAPTER 1

1.1 HW1

GENERAL ANALYSIS HW \_\_\_\_\_ PAGE 2 \_\_\_\_\_

2

## CHAPTER 2

2.1 HW1

COMPLEX ANALYSIS HW \_\_\_\_\_ PAGE 5 \_\_\_\_\_

5

## CHAPTER 3

3.1 HW1

PDE INTRO HW \_\_\_\_\_ PAGE 8 \_\_\_\_\_

8

# Chapter 1

## General Analysis HW

### 1.1 HW1

**Theorem 1.1.1.**  $\mathbb{R}^n$  is complete.

*Proof.* Let  $\mathbf{x}_k$  be an arbitrary Cauchy sequence in  $\mathbb{R}^n$ . We are required to show  $\mathbf{x}_k$  converge in  $\mathbb{R}^n$ . For each  $k$ , denote  $\mathbf{x}_k$  by  $(x_{(1,k)}, \dots, x_{(n,k)})$ . We claim that for each  $i \in \{1, \dots, n\}$

$x_{(i,k)}$  is a Cauchy sequence

Fix  $i$  and  $\epsilon > 0$ . To show  $x_{(i,k)}$  is a Cauchy sequence, we are required to find  $N \in \mathbb{N}$  such that for all  $r, m \geq N$  we have

$$|x_{(i,r)} - x_{(i,m)}| \leq \epsilon$$

Because  $\mathbf{x}_k$  is a Cauchy sequence in  $\mathbb{R}^n$ , we know there exists  $N \in \mathbb{N}$  such that for all  $r, m \geq N$ , we have

$$|\mathbf{x}_r - \mathbf{x}_m| < \epsilon$$

Fix such  $N$  and arbitrary  $r, m \geq M$ . Observe

$$|x_{(i,r)} - x_{(i,m)}| \leq \sqrt{\sum_{j=1}^n |x_{(j,r)} - x_{(j,m)}|^2} = |\mathbf{x}_r - \mathbf{x}_m| < \epsilon$$

We have proved that for each  $i \in \{1, \dots, n\}$ , the real sequence  $x_{(i,k)}$  is Cauchy. We now claim that for each  $i \in \{1, \dots, n\}$ , we have

$$\limsup_{r \rightarrow \infty} x_{(i,r)} \in \mathbb{R} \text{ and } \lim_{k \rightarrow \infty} x_{(i,k)} = \limsup_{r \rightarrow \infty} x_{(i,r)}$$

Again fix  $i$ . Because  $x_{(i,k)}$  is a Cauchy sequence, we know there exists some  $N$  such that for all  $r, m \geq N$ , we have

$$|x_{(i,r)} - x_{(i,m)}| < 1$$

This implies that for all  $r \geq N$ , we have

$$x_{(i,r)} < x_{(i,N)} + 1 \quad (1.1)$$

Equation 1.1 then tell us

$$x_{(i,N)} + 1 \text{ is an upper bound of } \{x_{(i,r)} : r \geq N\}$$

Then by definition of sup, we have

$$\sup\{x_{(i,r)} : r \geq N\} \leq x_{(i,N)} + 1 \in \mathbb{R}$$

This then implies  $\limsup_{r \rightarrow \infty} x_{(i,r)} \in \mathbb{R}$ . We now prove

$$\lim_{k \rightarrow \infty} x_{(i,k)} = \limsup_{r \rightarrow \infty} x_{(i,r)} \quad (1.2)$$

Fix  $\epsilon > 0$ . We are required to find  $N$  such that

$$\forall k \geq N, \left| x_{(i,k)} - \limsup_{r \rightarrow \infty} x_{(i,r)} \right| \leq \epsilon$$

Because  $\{x_{(i,k)}\}_{k \in \mathbb{N}}$  is a Cauchy sequence, we can let  $N_0$  satisfy

$$\forall k, m \geq N_0, |x_{(i,k)} - x_{(i,m)}| < \frac{\epsilon}{2}$$

Because  $\sup\{x_{(i,k)} : k \geq N'\} \searrow \limsup_{r \rightarrow \infty} x_{(i,r)}$  as  $N' \rightarrow \infty$ , we know there exists  $N_1 > N_0$  such that

$$\limsup_{r \rightarrow \infty} x_{(i,r)} - \frac{\epsilon}{2} < \sup\{x_{(i,k)} : k \geq N_0\} \leq \limsup_{r \rightarrow \infty} x_{(i,r)} + \frac{\epsilon}{2}$$

Then because  $\limsup_{r \rightarrow \infty} x_{(i,r)} - \frac{\epsilon}{2}$  is strictly smaller than the smallest upper bound of  $\{x_{(i,k)} : k \geq N_1\}$ , we see  $\limsup_{n \rightarrow \infty} x_{(i,r)} - \frac{\epsilon}{2}$  is not an upper bound of  $\{x_{(i,k)} : k \geq N_1\}$ . This implies the existence of some  $N$  such that  $N \geq N_1$  and

$$\limsup_{r \rightarrow \infty} x_{(i,r)} - \frac{\epsilon}{2} < x_{(i,N)} \leq \limsup_{r \rightarrow \infty} x_{(i,r)} + \frac{\epsilon}{2}$$

Now observe that for all  $k \geq N$ , because  $N \geq N_1 \geq N_0$

$$\limsup_{r \rightarrow \infty} x_{(i,r)} - \epsilon < x_{(i,N)} - \frac{\epsilon}{2} < x_{(i,k)} < x_{(i,N)} + \frac{\epsilon}{2} \leq \limsup_{r \rightarrow \infty} x_{(i,r)} + \epsilon$$

This implies for all  $k \geq N$ , we have

$$\left| x_{(i,k)} - \limsup_{r \rightarrow \infty} x_{(i,r)} \right| \leq \epsilon$$

We have just proved [Equation 1.2](#). Lastly, to close out the proof, we show

$$\lim_{k \rightarrow \infty} \mathbf{x}_k = \left( \lim_{k \rightarrow \infty} x_{(1,k)}, \dots, \lim_{k \rightarrow \infty} x_{(n,k)} \right) \quad (1.3)$$

Fix  $\epsilon > 0$ . For each  $i \in \{1, \dots, n\}$ , let  $N_i$  satisfy

$$\forall r \geq N_i, \left| x_{(i,r)} - \lim_{k \rightarrow \infty} x_{(i,k)} \right| \leq \frac{\epsilon}{\sqrt{n}}$$

Observe that for all  $r \geq \max_{i \in \{1, \dots, n\}} N_i$ , we have

$$\begin{aligned} \left| \mathbf{x}_r - \left( \lim_{k \rightarrow \infty} x_{(1,k)}, \dots, \lim_{k \rightarrow \infty} x_{(n,k)} \right) \right| &= \sqrt{\sum_{i=1}^n \left| x_{(i,r)} - \lim_{k \rightarrow \infty} x_{(i,k)} \right|^2} \\ &\leq \sqrt{\sum_{i=1}^n \frac{\epsilon^2}{n}} = \epsilon \end{aligned}$$

We have proved [Equation 1.3](#). ■

### **Theorem 1.1.2.** ( $\mathbb{Q}$ is dense in $\mathbb{R}$ )

*Proof.* Fix  $x \in \mathbb{R}$  and  $\epsilon > 0$ . To show  $\mathbb{Q}$  is dense in  $\mathbb{R}$ , we have to find  $q \in \mathbb{Q}$  such that  $|x - q| < \epsilon$ .

Let  $m \in \mathbb{N}$  satisfy  $\frac{1}{m} < \epsilon$ . Let  $n$  be the largest integer such that  $n \leq mx$ . Because  $n$  is the largest integer such that  $n \leq mx$ , we know  $mx - n < 1$ , otherwise we can deduce  $n + 1 \leq mx$ , which is impossible, since  $n + 1$  is an integer and  $n$  is the largest integer such that  $n \leq mx$ . We now see that

$$\frac{n}{m} \in \mathbb{Q} \text{ and } \left| x - \frac{n}{m} \right| = \frac{mx - n}{m} < \frac{1}{m} < \epsilon$$
■

# Chapter 2

## Complex Analysis HW

### 2.1 HW1

**Theorem 2.1.1.**

$$(1+i)^n, \frac{(1+i)^n}{n}, \frac{n!}{(1+i)^n} \text{ all diverge as } n \rightarrow \infty$$

*Proof.* Note that

$$|(1+i)^n| = 2^{\frac{n}{2}} \rightarrow \infty \text{ as } n \rightarrow \infty$$

This implies  $(1+i)$  is unbounded, thus diverge.

Note that

$$\left| \frac{(1+i)^n}{n} \right| = \frac{(\sqrt{2})^n}{n}$$

Observe

$$\begin{aligned} \frac{(\sqrt{2})^n}{n} &= \frac{[(\sqrt{2}-1)+1]^n}{n} = \frac{\sum_{k=0}^n \binom{n}{k} (\sqrt{2}-1)^k}{n} \\ &\geq \frac{\binom{n}{2} (\sqrt{2}-1)^2}{n} = (n-1) \left[ \frac{(\sqrt{2}-1)^2}{2} \right] \rightarrow \infty \text{ as } n \rightarrow \infty \end{aligned}$$

This implies  $\frac{(1+i)^n}{n}$  is unbounded, thus diverge.

Note that

$$\left| \frac{n!}{(1+i)^n} \right| = \frac{n!}{(\sqrt{2})^n}$$

Note that for all  $k \geq 8$ , we have

$$\frac{k}{\sqrt{2}} \geq \frac{\sqrt{8}}{\sqrt{2}} = 2$$

This implies

$$\frac{n!}{(\sqrt{2})^n} = \prod_{k=1}^n \frac{k}{\sqrt{2}} = \frac{7!}{(\sqrt{2})^7} \prod_{k=8}^n \frac{k}{\sqrt{2}} \geq \frac{7!}{(\sqrt{2})^7} \prod_{k=8}^n 2 \geq \frac{7!2^{n-8+1}}{(\sqrt{2})^7} \rightarrow \infty$$

which implies  $\frac{n!}{(1+i)^n}$  is unbounded, thus diverge. ■

**Theorem 2.1.2.**

$$n!z^n \text{ converge} \iff z = 0$$

*Proof.* If  $z = 0$ , then  $n!z^n = 0$  for all  $n$ , which implies  $n!z^n \rightarrow 0$ . Now, suppose  $z \neq 0$ . Let  $M \in \mathbb{N}$  satisfy  $|z| > \frac{1}{M}$ . Observe

$$|n!z^n| = \left| \prod_{k=1}^n kz \right| = \left| \prod_{k=1}^{2M-1} kz \right| \left| \prod_{k=2M}^n kz \right| \geq \left| \prod_{k=1}^{2M-1} kz \right| \left| \prod_{k=2M}^n 2Mz \right| \geq \left| \prod_{k=1}^{2M-1} kz \right| 2^{n-2M+1} \rightarrow \infty$$

This implies  $n!z^n$  is unbounded, thus diverge. ■

**Theorem 2.1.3.**

$$u_n \rightarrow u \implies v_n \triangleq \sum_{k=1}^n \frac{u_k}{n} \rightarrow u$$

*Proof.* Because

$$\sum_{k=1}^n \frac{u_k}{n} = \sum_{k \leq \sqrt{n}} \frac{u_k}{n} + \sum_{\sqrt{n} < k \leq n} \frac{u_k}{n}$$

It suffices to prove

$$\sum_{k \leq \sqrt{n}} \frac{u_k}{n} \rightarrow 0 \text{ and } \sum_{\sqrt{n} < k \leq n} \frac{u_k}{n} \rightarrow u \text{ as } n \rightarrow \infty$$

Because  $u_n$  converge, we can let  $M$  bound  $|u_n|$ . Observe

$$\left| \sum_{k \leq \sqrt{n}} \frac{u_k}{n} \right| \leq \sum_{k \leq \sqrt{n}} \left| \frac{u_k}{n} \right| \leq \sum_{k \leq \sqrt{n}} \frac{M}{n} \leq \frac{M\sqrt{n}}{n} = \frac{M}{\sqrt{n}} \rightarrow 0 \text{ as } n \rightarrow \infty \text{ (done)}$$

Because

$$\sum_{\sqrt{n} < k \leq n} \frac{u_k}{n} = \frac{n - \lceil \sqrt{n} \rceil + 1}{n} \sum_{\sqrt{n} < k \leq n} \frac{u_k}{n - \lceil \sqrt{n} \rceil + 1}$$

and

$$\lim_{n \rightarrow \infty} \frac{n - \lceil \sqrt{n} \rceil + 1}{n} = 1$$

We can reduce the problem into proving

$$\lim_{n \rightarrow \infty} \sum_{\sqrt{n} < k \leq n} \frac{u_k}{n - \lceil \sqrt{n} \rceil + 1} = u$$

Fix  $\epsilon$ . Let  $N$  satisfy that for all  $n \geq N$ , we have  $|u_n - u| < \epsilon$ . Then for all  $n \geq N^2$ , we have

$$\begin{aligned} \left| \left( \sum_{\sqrt{n} < k \leq n} \frac{u_k}{n - \lceil \sqrt{n} \rceil + 1} \right) - u \right| &= \left| \sum_{\sqrt{n} < k \leq n} \frac{u_k - u}{n - \lceil \sqrt{n} \rceil + 1} \right| \\ &\leq \sum_{\sqrt{n} < k \leq n} \frac{|u_k - u|}{n - \lceil \sqrt{n} \rceil + 1} \\ &\leq \sum_{\sqrt{n} < k \leq n} \frac{\epsilon}{n - \lceil \sqrt{n} \rceil + 1} = \epsilon \text{ (done)} \end{aligned}$$

■



# Chapter 3

## PDE intro HW

### 3.1 HW1

**Theorem 3.1.1.**

Show  $u \mapsto u_x + uu_y$  is non-linear

*Proof.* See that

$$2u \mapsto 2u_x + 4uu_y \neq 2(u_x + uu_y) \quad (3.1)$$

■

**Theorem 3.1.2.**

Solve  $(1 + x^2)u_x + u_y = 0$

*Proof.* The characteristic curve has the derivative

$$\frac{dy}{dx} = \frac{1}{1 + x^2}$$


The solution to this ODE is

$$y = \arctan x + C$$

We now see that the solution to the PDE in [Equation 3.1](#) is

$u = f((\arctan x) - y)$  where  $f : \mathbb{R} \rightarrow \mathbb{R}$  is an arbitrary smooth function

A characteristic curve is as followed.



■

### Theorem 3.1.3.

$$\text{Solve } au_x + bu_y + cu = 0 \quad (3.2)$$

*Proof.* Fix

$$\begin{cases} x' \triangleq ax + by \\ y' \triangleq bx - ay \end{cases}$$

This map is clearly a diffeomorphism. Compute

$$\begin{cases} u_x = \frac{\partial u}{\partial x'} \frac{\partial x'}{\partial x} + \frac{\partial u}{\partial y'} \frac{\partial y'}{\partial x} = au_{x'} + bu_{y'} \\ u_y = \frac{\partial u}{\partial x'} \frac{\partial x'}{\partial y} + \frac{\partial u}{\partial y'} \frac{\partial y'}{\partial y} = bu_{x'} - au_{y'} \end{cases}$$

Plugging it back into the PDE in [Equation 3.2](#), we have

$$cu + (a^2 + b^2)u_{x'} = 0 \quad (3.3)$$

If  $c = a^2 + b^2 = 0$ , then all smooth functions are solution. If  $a^2 + b^2 = 0$  but  $c \neq 0$ , then clearly the only solution is  $u = \tilde{0}$ . If  $a^2 + b^2 \neq 0$  but  $c = 0$ , then  $u_{x'} = \tilde{0}$ , which implies  $u = f(y')$  where  $y' = bx - ay$  and  $f$  can be arbitrary smooth function.

Now, suppose  $a^2 + b^2 \neq 0 \neq c$ , note that the PDE in [Equation 3.3](#) is just an ODE of the form

$$y + \frac{a^2 + b^2}{c}y' = 0$$

9

The general solution to this ODE is

$$y = Ce^{\frac{-ct}{a^2+b^2}}$$

In other words, the general solution of the PDE in **Equation 3.3** is

$$u = Ce^{\frac{-cx'}{a^2+b^2}} = Ce^{\frac{-c(ax+by)}{a^2+b^2}}$$

