

Name: _____

University of Illinois

Spring 2020

CS 446/ECE 449 Machine Learning
Homework 6: Structured Prediction

Due on Thursday March 12 2020, noon Central Time

1. [28 points] Structured Prediction

We are interested in jointly predicting/modeling two discrete random variables $y = (y_1, y_2) \in \mathcal{Y}$ with $y_i \in \mathcal{Y}_i = \{0, 1\}$ for $i \in \{1, 2\}$ and $\mathcal{Y} = \prod_{i \in \{1, 2\}} \mathcal{Y}_i$. We define the joint probability distribution to be $p(y) = p(y_1, y_2) = \frac{1}{Z} \exp F(y)$.

- (a) (3 points) What is the value of Z (in terms of $F(y)$) and what is Z called? How many configurations do we need to sum over? Provide the expression using \mathcal{Y}_i .

Your answer:

- (b) (6 points) Next we want to solve (for any hyperparameter ϵ)

$$\max_{\hat{p} \in \Delta_{\mathcal{Y}}} \sum_{y \in \mathcal{Y}} \hat{p}(y) F(y) - \sum_{y \in \mathcal{Y}} \epsilon \hat{p}(y) \log \hat{p}(y), \quad (1)$$

where $\Delta_{\mathcal{Y}}$ denotes the probability simplex, *i.e.*, \hat{p} is a valid probability distribution over its domain \mathcal{Y} . Using general notation, write down the Lagrangian and compute its derivative w.r.t. $\hat{p}(y) \forall y \in \mathcal{Y}$. Subsequently, find the optimal \hat{p}^* . What is the resulting optimal cost function value for the program given in Eq. (1)? How does this result relate to part (a)?

Your answer:

Name: _____

- (c) (3 points) For the program in Eq. (1) assume now $\epsilon = 0$, *i.e.*, we are searching for that configuration $y^* = \arg \max_{\hat{y} \in \mathcal{Y}} F(\hat{y})$ which maximizes $F(y)$. Assume $F(y) = f_1(y_1) + f_2(y_2) + f_{1,2}(y_1, y_2)$. How many different values can the functions f_1 , f_2 and $f_{1,2}$ result in?

Your answer:

- (d) (9 points) As discussed in class, finding the global maximizer can be equivalently written as the following integer linear program:

$$\max_b \sum_{r, y_r} b_r(y_r) f_r(y_r) \quad \text{s.t.} \quad \begin{cases} b_r(y_r) \in \{0, 1\} & \forall r, y_r \\ \sum_{y_r} b_r(y_r) = 1 & \forall r \\ \sum_{y_p \setminus y_r} b_p(y_p) = b_r(y_r) & \forall r, p \in P(r), y_r \end{cases} \quad (2)$$

Using the decomposition $F(y) = f_1(y_1) + f_2(y_2) + f_{1,2}(y_1, y_2)$, *i.e.*, for $r \in \{\{1\}, \{2\}, \{1, 2\}\}$, explicitly state the integer linear program and all its constraints for the special case that $\mathcal{Y}_i = \{0, 1\}$ for $i \in \{1, 2\}$. (**Hint:** The parent sets are as follows: $P(\{1\}) = \{1, 2\}$ and $P(\{2\}) = \{1, 2\}$. Use notation such as $f_1(y_1 = 0)$ and $b_1(y_1 = 0)$.)

Your answer:

Name: _____

- (e) (3 points) Let b be the vector

$$b = [\quad b_1(y_1 = 0), b_1(y_1 = 1), b_2(y_2 = 0), b_2(y_2 = 1), \\ b_{1,2}(y_1 = 0, y_2 = 0), b_{1,2}(y_1 = 1, y_2 = 0), b_{1,2}(y_1 = 0, y_2 = 1), b_{1,2}(y_1 = 1, y_2 = 1)]^\top.$$

Specify all but the integrality constraints of part (d) using matrix vector notation, *i.e.*, provide A and c for $Ab = c$.

Your answer:

- (f) (4 points) Complete `A6_Structure.py` where we approximately solve the integer linear program using the linear programming relaxation. Implement the constraints. Why do we provide $-f$ as input to the solver? What is the obtained result b for the relaxation of the program given in Eq. (2) and its cost function value? Is this the configuration y^* which has the largest score?

Your answer: