

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

Diffusion processes with applications in diffusion networks

W. Genuist^{1,2}, É. Savin^{1,3}
{wilfried.genuist,eric.savin}@centralesupelec.fr

¹Computer Science Dept.
ONERA, France

²Laboratoire de Mécanique Paris-Saclay
CentraleSupélec, France

³Mechanical and Environmental Engineering Dept.
CentraleSupélec, France

May 5, 2024

Outline

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- 1 Stochastic differential equations (SDE)
 - First-order stochastic systems driven by noise
 - Stochastic integrals
 - Diffusion processes
- 2 Numerical simulations of SDE
 - Stochastic modeling with SDE
 - Numerical schemes
- 3 Generative Adversarial Networks (GAN)
- 4 Variational Auto-Encoders (VAE)
- 5 Diffusion models in ML
 - Continuous setting
 - Denoising Diffusion Probabilistic Modeling (DDPM)
 - Score Matching with Langevin Dynamics (SMLD)

Outline

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- 1 Stochastic differential equations (SDE)
 - First-order stochastic systems driven by noise
 - Stochastic integrals
 - Diffusion processes
- 2 Numerical simulations of SDE
 - Stochastic modeling with SDE
 - Numerical schemes
- 3 Generative Adversarial Networks (GAN)
- 4 Variational Auto-Encoders (VAE)
- 5 Diffusion models in ML
 - Continuous setting
 - Denoising Diffusion Probabilistic Modeling (DDPM)
 - Score Matching with Langevin Dynamics (SMLD)

First-order stochastic differential equation

SDE

W. Genuist,
É. Savin

SDE

First-order
systems

Stochastic
integrals

Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

A general first-order stochastic differential equation for the process U indexed on \mathbb{R}_+ with values in \mathbb{R}^q :

$$\dot{U}(t) = \underline{b}(U, t) + \sigma(U, t)F(t), \quad U(0) = U_0,$$

with the data:

- $u, t \mapsto \underline{b}(u, t) : \mathbb{R}^q \times \mathbb{R}_+ \rightarrow \mathbb{R}^q$ the *drift* function;
- $u, t \mapsto \sigma(u, t) : \mathbb{R}^q \times \mathbb{R}_+ \rightarrow \mathbb{M}_{q,p}(\mathbb{R})$ the *scattering* operator;
- U_0 is an r.v. in \mathbb{R}^q with known marginal PDF $\pi_0(u_0)$;
- $F(t) = (F_1(t), \dots, F_p(t))$ is a second-order Gaussian random process indexed on \mathbb{R} with values in \mathbb{R}^p , also centered, stationary, such that $F_1(t), \dots, F_p(t)$ are mutually independent and mean-square continuous, with:

$$S_F(\omega) = S_0 \mathbf{1}_{[-B, B]}(\omega) [I_p], \quad S_0 > 0, \quad B > 0.$$

First-order systems driven by noise

SDE

W. Genuist,
É. Savin

SDE

First-order
systems

Stochastic
integrals

Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- $B < +\infty$: colored noise, hot topic!

$$\mathbf{U}(t) = \mathbf{U}_0 + \int_0^t \underline{\mathbf{b}}(\mathbf{U}(s), s) ds + \int_0^t \boldsymbol{\sigma}(\mathbf{U}(s), s) \mathbf{F}(s) ds .$$

- $B \rightarrow +\infty$: $\mathbf{F} \rightarrow \dot{\mathbf{W}}$ the *normalized Gaussian white noise*, and the solution of the first-order SDE holds as a “stochastic integral”:

$$\mathbf{U}(t) = \mathbf{U}_0 + \int_0^t \underline{\mathbf{b}}(\mathbf{U}(s), s) ds + \int_0^t \boldsymbol{\sigma}(\mathbf{U}(s), s) \circ d\mathbf{W}(s) .$$

- *Causality*: the family of r.v. $\{\mathbf{U}(s), 0 \leq s \leq t\}$ is independent of the family of r.v. $\{\mathbf{F}(\tau), \tau > t\}$ or $\{d\mathbf{W}(\tau), \tau > t\}$.

White noise

Definition

SDE

W. Genuist,
É. Savin

SDE

First-order
systems

Stochastic
integrals

Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case

DDPM
SMLD

Definition

- *The normalized Gaussian white noise $\mathbf{B}(t) \equiv \dot{\mathbf{W}}(t)$ with values in \mathbb{R}^p is the Gaussian stochastic process indexed on \mathbb{R} , centered, stationary, with the spectral density matrix:*

$$\mathbf{S}_{\mathbf{B}}(\omega) = \frac{1}{2\pi} \mathbf{I}_p.$$

- *Since $B_1(t), \dots, B_p(t)$ are uncorrelated and jointly Gaussian, they are mutually independent.*
- *\mathbf{B} is not second order $\|\mathbf{B}(t)\|^2 = \int \text{Tr } \mathbf{S}_{\mathbf{B}}(\omega) d\omega = +\infty$.*

This definition holds in the sense of generalized stochastic processes $\varphi \mapsto \mathbf{B}(\varphi) : \mathcal{D}(T) \rightarrow L^2(\Omega, \mathbb{R}^p)$ where $\mathcal{D}(T)$ is the set of \mathcal{C}^∞ functions having a compact support within $T \subseteq \mathbb{R}$.

White noise

Definition

SDE

W. Genuist,
É. Savin

SDE

First-order
systems

Stochastic
integrals

Diffusion
processes

Numerical
solutions

Stochastic
modeling

Numerical
schemes

GAN


VAE

Diffusion &
ML

Continuous
case

DDPM

SMLD



White noise.

Wiener process

Definition

SDE

W. Genuist,
É. Savin

SDE

First-order
systems

Stochastic
integrals

Diffusion
processes

Numerical
solutions

Stochastic
modeling

Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case

DDPM

SMLD

The white noise is the (generalized) derivative of the Wiener process, or *Brownian motion*.

Definition

The (normalized) Wiener process $\mathbf{W}(t)$ with values in \mathbb{R}^p is the stochastic process indexed on \mathbb{R}_+ , such that:

- $W_1(t), \dots, W_p(t)$ are mutually independent;
- $\mathbf{W}(0) = \mathbf{0}$ almost surely (a.s.);
- If $0 \leq s < t < +\infty$ let $\Delta \mathbf{W}(s, t) = \mathbf{W}(t) - \mathbf{W}(s)$, then:
 - $\forall m$ and $0 < t_1 < t_2 < \dots < t_m < +\infty$, $\mathbf{W}(0)$, $\Delta \mathbf{W}(0, t_1)$, $\Delta \mathbf{W}(t_1, t_2)$, ... $\Delta \mathbf{W}(t_{m-1}, t_m)$ are mutually independent r.v. (independent increments);
 - $\Delta \mathbf{W}(s, t)$ is a Gaussian, centered, second-order r.v. with $C_{\Delta \mathbf{W}}(s, t) = (t - s)\mathbf{I}_p$.

Wiener process

Characterization

SDE

W. Genuist,
É. Savin

SDE

First-order
systems

Stochastic
integrals

Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

Consequently it can be shown that:

- $\mathbf{W}(t)$ is a second-order Gaussian, centered, mean-square continuous, non stationary stochastic process;
- the covariance and conditional PDF for $0 \leq t, s < +\infty$:

$$\mathbf{C}_{\mathbf{W}}(t, s) = \text{Min}(t, s) \mathbf{I}_p,$$

$$\pi_{\tau}(\mathbf{v}'; t + s | \mathbf{v}; t) = (2\pi s)^{-\frac{p}{2}} e^{-\frac{\|\mathbf{v}' - \mathbf{v}\|^2}{2s}};$$

- $\mathbf{W}(t)$ has a.s. continuous sample paths;
- sample paths $t \mapsto \mathbf{W}(t, \theta)$, $\theta \in \Omega_{\theta}$, are non differentiable a.s.

As a generalized derivative with $d\mathbf{W} = (dW_1, \dots, dW_p)$:

$$d\mathbf{W}(\varphi) = \mathbf{B}(-\dot{\varphi}), \quad \forall \varphi \in \mathcal{D}(\mathbb{R}).$$

Wiener process

Characterization

SDE

W. Genuist,
É. Savin

SDE

First-order
systems

Stochastic
integrals
Diffusion
processes

Numerical
solutions


Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD



Wiener process in \mathbb{R}^3 .

Stochastic integrals

Definition

SDE

W. Genuist,
É. Savin

SDE

First-order
systems

Stochastic
integrals

Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- Let $\mathbf{X}(t)$ be a stochastic process indexed by \mathbb{R}_+ with a.s. continuous sample paths.
- Assume the r.v. $\{\mathbf{X}(s), 0 \leq s \leq t\}$ are independent of the r.v. $\{\Delta \mathbf{W}(t, \tau), \tau > t\}$: a *non anticipative* process, then

$$\begin{aligned} & \int_0^t \mathbf{X}(s) d_\lambda \mathbf{W}(s) \\ &= \text{l. i. p.} \sum_{K \rightarrow +\infty}^K [(1 - \lambda) \mathbf{X}(t_k) + \lambda \mathbf{X}(t_{k+1})] \Delta \mathbf{W}(t_k, t_{k+1}), \end{aligned}$$

for any partition $0 = t_1 < t_2 < \dots < t_{K+1} = t$ of $[0, t]$
with $\max_{1 \leq k \leq K} (t_{k+1} - t_k) \xrightarrow{K \rightarrow +\infty} 0$.

Stochastic integrals

Application to stochastic differential calculus

SDE

W. Genuist,
É. Savin

SDE

First-order
systems

Stochastic
integrals

Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- A simple example—remind $\Delta W \propto \Delta t^{\frac{1}{2}}$ for the real-valued Wiener process W :

$$\int_0^t W(s) d_\lambda W(s) = \frac{1}{2} W(t)^2 + \left(\lambda - \frac{1}{2} \right) t,$$

from which one deduces the stochastic differential:

$$d_\lambda(W(t)^2) = 2W(t)d_\lambda W(t) + (1 - 2\lambda)dt.$$

- More generally ($\lambda = 0$ is called the *Itô formula*):

$$d_\lambda(f(W(t))) = f'(W(t))d_\lambda W(t) + \left(\frac{1}{2} - \lambda \right) f''(W(t))dt.$$

Stochastic integrals

Application to stochastic differential calculus

SDE

Setting $W(t_k) = W_k$:

$$\begin{aligned}\int_0^t W(s) d_\lambda W(s) &= \text{l.i.p.} \sum_{K \rightarrow +\infty}^K [(1-\lambda)W_k + \lambda W_{k+1}] (W_{k+1} - W_k) \\&= \text{l.i.p.} \sum_{K \rightarrow +\infty}^K [(1-2\lambda)W_k W_{k+1} - \lambda W_{k+1}^2 - (1-\lambda)W_k^2] \\&= \text{l.i.p.} \sum_{K \rightarrow +\infty}^K \frac{1}{2} [(2\lambda-1)(W_{k+1} - W_k)^2 + W_{k+1}^2 - W_k^2] \\&= \frac{1}{2}(2\lambda-1) \text{l.i.p.} \sum_{K \rightarrow +\infty}^K (W_{k+1} - W_k)^2 + \frac{1}{2}(W_{K+1}^2 - W_1^2) \\&= \frac{1}{2}(2\lambda-1)t + \frac{1}{2}W(t)^2,\end{aligned}$$

since $W_1 = W(0) = 0$ and $W_{K+1} = W(t)$. Besides:

$$\mathbb{E} \left\{ \int_0^t W(s) d_\lambda W(s) \right\} = \lambda t.$$

Stochastic integrals

Stratonovich-Itô

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- If $\lambda = \frac{1}{2}$ the usual differential calculus applies, and the solution of SDE holds as a *Stratonovich integral* (1966):

$$U(t) = U_0 + \int_0^t \underline{b}(U(s), s) ds + \int_0^t \sigma(U(s), s) \circ dW(s).$$

- If $\lambda = 0$, its solution holds as an *Itô integral* (1944):

$$U(t) = U_0 + \int_0^t \mathbf{b}(U(s), s) ds + \int_0^t \sigma(U(s), s) dW(s),$$

where:

$$\mathbf{b} = \underline{b} + \frac{1}{2}(D_u \sigma) \sigma^\top.$$

- $U(t)$ is a *Markov process*.

Stochastic integrals

Itô's formula

SDE

W. Genuist,
É. Savin

SDE

First-order
systems

Stochastic
integrals

Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- Let $\mathbf{U}(t) \in \mathbb{R}^q$ be the solution of the ISDE:

$$\mathbf{U}(t) = \mathbf{U}_0 + \int_0^t \mathbf{b}(\mathbf{U}(s), s) ds + \int_0^t \boldsymbol{\sigma}(\mathbf{U}(s), s) d\mathbf{W}(s).$$

- Let $\phi : \mathbb{R}^q \times \mathbb{R} \rightarrow \mathbb{R}$ be a smooth function. Then *Itô's formula* states that:

$$\begin{aligned} \phi(\mathbf{U}(t), t) &= \phi(\mathbf{U}_0, 0) + \int_0^t \frac{\partial \phi}{\partial t}(\mathbf{U}(s), s) ds \\ &\quad + \int_0^t \nabla_{\mathbf{u}} \phi(\mathbf{U}(s), s) \cdot d\mathbf{U}(s) \\ &\quad + \frac{1}{2} \int_0^t \boldsymbol{\sigma}(\mathbf{U}(s), s) \boldsymbol{\sigma}(\mathbf{U}(s), s)^\top : \nabla_{\mathbf{u}} \otimes \nabla_{\mathbf{u}} \phi(\mathbf{U}(s), s) ds, \end{aligned}$$

where $d\mathbf{U}(s) = \mathbf{b}(\mathbf{U}(s), s) ds + \boldsymbol{\sigma}(\mathbf{U}(s), s) d\mathbf{W}(s)$.

Markov processes

Definition

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

Definition

The conditional probability given $t_0 < \dots < t_m < t$:

$$\pi_{\tau}(\mathbf{u}; t | \mathbf{u}_0, \dots, \mathbf{u}_m; t_0, \dots, t_m) = \frac{\pi(\mathbf{u}_0, \dots, \mathbf{u}_m, \mathbf{u}; t_0, \dots, t_m, t)}{\pi(\mathbf{u}_0, \dots, \mathbf{u}_m; t_0, \dots, t_m)}.$$

Definition

Let $\mathbf{U}(t)$ be a stochastic process defined on (Ω, \mathcal{E}, P) and indexed on \mathbb{R}_+ with values in \mathbb{R}^q . It is a Markov process if:

- *for all $0 \leq t_1 < \dots < t_m < t$ and $\mathbf{u}_1, \dots, \mathbf{u}_m, \mathbf{u}$ in \mathbb{R}^q*

$$\pi_{\tau}(\mathbf{u}; t | \mathbf{u}_0, \dots, \mathbf{u}_m; t_0, \dots, t_m) = \pi_{\tau}(\mathbf{u}; t | \mathbf{u}_m; t_m);$$

- *the marginal PDF $\pi_0(\mathbf{u}_0)$ of $\mathbf{U}(0)$ can be any PDF.*

Markov processes

Chapman-Kolmogorov equation

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- A Markov process is fully characterized by:
 - its marginal PDF $\pi(\mathbf{u}; t)$,
 - and its *transition PDF* $\pi_{\tau}(\mathbf{u}; t | \mathbf{v}; s)$, $0 \leq s < t < +\infty$,
with

$$\pi(\mathbf{u}; t) = \int_{\mathbb{R}^q} \pi_{\tau}(\mathbf{u}; t | \mathbf{v}; s) \pi(\mathbf{v}; s) d\mathbf{v}.$$

- π_{τ} satisfies the *Chapman-Kolmogorov equation*:

$$\pi_{\tau}(\mathbf{u}; t | \mathbf{u}'; t') = \int_{\mathbb{R}^q} \pi_{\tau}(\mathbf{u}; t | \mathbf{v}; s) \pi_{\tau}(\mathbf{v}; s | \mathbf{u}'; t') d\mathbf{v}, \quad t' < s < t.$$

- Homogeneous Markov process:

$$\pi_{\tau}(\mathbf{u}; t | \mathbf{v}; s) = \pi_{\tau}(\mathbf{u}; t - s | \mathbf{v}; 0), \quad 0 \leq s < t < +\infty.$$

- The Brownian motion is a Markov process.

Diffusion processes

Definition

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

Definition

The \mathbb{R}^q -valued continuous-time Markov process $\mathbf{U}(t)$ with a.s. continuous sample paths and transition PDF $\pi_\tau(\mathbf{v}; s|\mathbf{u}; t)$ is a diffusion process if $\forall \epsilon > 0$ (but not necessarily small), $\forall \mathbf{u} \in \mathbb{R}^q$ the first moments of its increments are such that for $h > 0$:

$$\int_{\|\mathbf{v}-\mathbf{u}\| \geq \epsilon} \pi_\tau(d\mathbf{v}; t+h|\mathbf{u}; t) = o(h),$$

$$\int_{\|\mathbf{v}-\mathbf{u}\| < \epsilon} (\mathbf{v} - \mathbf{u}) \pi_\tau(d\mathbf{v}; t+h|\mathbf{u}; t) = h\mathbf{b}(\mathbf{u}, t) + o(h),$$

$$\int_{\|\mathbf{v}-\mathbf{u}\| < \epsilon} (\mathbf{v} - \mathbf{u}) \otimes (\mathbf{v} - \mathbf{u}) \pi_\tau(d\mathbf{v}; t+h|\mathbf{u}; t) = h\mathbf{a}(\mathbf{u}, t) + o(h),$$

where $\mathbf{b} \in \mathbb{R}^q$ and $\mathbf{a} \in \mathbb{R}^{q \times q}$ symmetric, positive.

Diffusion processes

Interpretation

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- *Continuity*: particles moving on sample paths of a diffusion process only make small jumps, or the probability of moving a distance ϵ goes to zero as h goes to zero no matter how small ϵ is.
- *Drift*: those particles can have a net mean velocity \mathbf{b} .
- *Diffusion*: particles spread as time increases with the rate $\text{Tr } \mathbf{a}$. Entropy increases while the phase space contracts, thus some information (energy) gets lost.

$$\mathbf{U}(t+h) - \mathbf{U}(t) \approx h\mathbf{b}(\mathbf{U}(t), t) + \mathbf{a}^{\frac{1}{2}}(\mathbf{U}(t), t)\Delta\mathbf{W}(t, t+h).$$

Diffusion processes

Infinitesimal generator

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- *Infinitesimal generator*: for a time-continuous process $U(t)$ and a suitably regular function ϕ , the infinitesimal generator \mathcal{A}_t is

$$\mathcal{A}_t \phi(\mathbf{u}, t) = \lim_{h \downarrow 0} \frac{\mathbb{E}\{\phi(\mathbf{U}(t+h), t+h) | \mathbf{U}(t) = \mathbf{u}\} - \phi(\mathbf{u}, t)}{h}.$$

- For a diffusion process:

$$\mathcal{A}_t \phi(\mathbf{u}, t) = \partial_t \phi + \mathbf{b}(\mathbf{u}, t) \cdot \nabla_{\mathbf{u}} \phi + \frac{1}{2} \mathbf{a}(\mathbf{u}, t) : \nabla_{\mathbf{u}} \otimes \nabla_{\mathbf{u}} \phi,$$

with (formal) adjoint operator:

$$\mathcal{A}_t^* \phi(\mathbf{u}, t) = -\partial_t \phi - \nabla_{\mathbf{u}} \cdot (\mathbf{b}(\mathbf{u}, t) \phi) + \frac{1}{2} \nabla_{\mathbf{u}} \otimes \nabla_{\mathbf{u}} : (\mathbf{a}(\mathbf{u}, t) \phi).$$

Fokker-Planck equation

SDE

W. Genuist,
É. Savin

The marginal PDF π and transition PDF π_τ of a diffusion process satisfy the *Fokker-Planck equation* $\mathcal{A}_t^* \pi = 0$, or:

$$\partial_t \pi + \nabla_{\mathbf{u}} \cdot \left(\pi \mathbf{b} - \frac{1}{2} \nabla_{\mathbf{u}} \cdot (\pi \mathbf{a}) \right) = 0,$$

with $\pi(\mathbf{u}_0; 0) = \pi_0(\mathbf{u}_0)$ and $\lim_{h \downarrow 0} \pi_\tau(\mathbf{u}; t+h|\mathbf{v}; t) = \delta(\mathbf{u} - \mathbf{v})$.

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

$$\begin{aligned} \int_{\mathbb{R}^q} f(\mathbf{u}) \partial_t \pi_\tau(\mathbf{u}; t|\mathbf{v}; s) d\mathbf{u} &= \lim_{h \downarrow 0} \frac{1}{h} \int_{\mathbb{R}^q} f(\mathbf{u}) (\pi_\tau(\mathbf{u}; t+h|\mathbf{v}; s) - \pi_\tau(\mathbf{u}; t|\mathbf{v}; s)) d\mathbf{u} \\ &= \lim_{h \downarrow 0} \frac{1}{h} \int_{\mathbb{R}^q} \pi_\tau(\mathbf{u}; t|\mathbf{v}; s) \left[\int_{\mathbb{R}^q} f(\mathbf{u}') \pi_\tau(\mathbf{u}'; t+h|\mathbf{u}; t) d\mathbf{u}' - f(\mathbf{u}) \right] d\mathbf{u} \quad (\text{C-K}) \\ &= \lim_{h \downarrow 0} \frac{1}{h} \int_{\mathbb{R}^q} \pi_\tau(\mathbf{u}; t|\mathbf{v}; s) \int_{\mathbb{R}^q} (f(\mathbf{u}') - f(\mathbf{u})) \pi_\tau(\mathbf{u}'; t+h|\mathbf{u}; t) d\mathbf{u}' d\mathbf{u} \quad (\text{norm.}) \\ &= \lim_{h \downarrow 0} \frac{1}{h} \int_{\mathbb{R}^q} \pi_\tau(\mathbf{u}; t|\mathbf{v}; s) \int_{\|\mathbf{u}' - \mathbf{u}\| < \epsilon} (f(\mathbf{u}') - f(\mathbf{u})) \pi_\tau(\mathbf{u}'; t+h|\mathbf{u}; t) d\mathbf{u}' d\mathbf{u}, \quad \forall f \in \mathcal{C}_0^2. \end{aligned}$$

Then use a Taylor expansion for f , definitions of drift and diffusion, and integrate by parts.

Backward Kolmogorov equation

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

The transition PDF $\pi_\tau(\cdot|\mathbf{v}; s)$ of a diffusion process also satisfies the *Backward Kolmogorov equation* $\mathcal{A}_s\pi_\tau = 0$, or:

$$\begin{aligned} \partial_s \pi_\tau(\cdot|\mathbf{v}; s) + \mathbf{b}(\mathbf{v}, s) \cdot \nabla_{\mathbf{v}} \pi_\tau(\cdot|\mathbf{v}; s) \\ + \frac{1}{2} \mathbf{a}(\mathbf{v}, s) : \nabla_{\mathbf{v}} \otimes \nabla_{\mathbf{v}} \pi_\tau(\cdot|\mathbf{v}; s) = 0, \end{aligned}$$

with $\lim_{h \downarrow 0} \pi_\tau(\mathbf{u}; s|\mathbf{v}; s-h) = \delta(\mathbf{v} - \mathbf{u})$.

Itô's stochastic differential equations (ISDE)

Solutions

SDE

W. Genuist,
É. Savin

$$dU = b(U, t)dt + \sigma(U, t)dW, \quad U(0) = U_0,$$

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

with the regularity assumptions:

$$\begin{aligned} \|b(u, t)\| + \|\sigma(u, t)\| &\leq K(1 + \|u\|), \\ \|b(u', t) - b(u, t)\| + \|\sigma(u', t) - \sigma(u, t)\| &\leq K \|u' - u\|. \end{aligned}$$

- 1 Then the SDE has a unique solution, with a.s. continuous sample paths. If in addition b and σ are independent of t , $U(t)$ is homogeneous.
- 2 If $t \mapsto b(u, t) \in \mathbb{R}^q$ and $t \mapsto \sigma(u, t) \in \mathbb{R}^{q \times p}$ are continuous, $U(t)$ is also a diffusion process with $a = \sigma \sigma^\top$.

Itô's stochastic differential equations (ISDE)

Example: Black-Scholes¹ model

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- The relative variation of a stock $U(t)$ with constant (annualized) drift rate μ and volatility σ :

$$\frac{dU}{U} = \mu dt + \sigma dW, \quad U(0) = U_0.$$

- Transformation to a Stratonovich SDE:

$$\frac{dU}{U} = \left(\mu - \frac{\sigma^2}{2} \right) dt + \sigma \circ dW, \quad U(0) = U_0,$$

for which “normal rules of integration” apply:

$$U(t) = U_0 e^{(\mu - \frac{\sigma^2}{2})t + \sigma W(t)}.$$

- The Fokker-Planck equation:

$$\partial_t \pi + \mu \partial_u (\pi u) - \frac{\sigma^2}{2} \partial_u^2 (\pi u^2) = 0, \quad \pi(u; 0) = \pi_0(u).$$

¹Fischer Black (1938–1995), Myron Scholes (1941–): American financial economists. M. Scholes received the Sveriges Riksbank Prize in Economic Sciences in Memory of A. Nobel in 1997 for this model for valuing options, together with Robert Merton (1944–).

Time reversal of diffusions²

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- Let $T > 0$:

$$\bar{b}(u, t) = -b(u, T - t) + \frac{\nabla u \cdot (\pi(u, T - t)a(u, T - t))}{\pi(u, T - t)},$$

$$\bar{\sigma}(u, t) = \sigma(u, T - t),$$

and:

$$\bar{\mathcal{A}}_t \phi(u) = \bar{b}(u, t) \cdot \nabla u \phi + \frac{1}{2} \bar{a}(u, t) : \nabla u \otimes \nabla u \phi,$$

with $\bar{a} = \bar{\sigma} \bar{\sigma}^\top$.

- Then $\bar{U}(t) = U(T - t)$ is a Markov diffusion process with infinitesimal generator $\bar{\mathcal{A}}_t$, such that:

$$d\bar{U} = \bar{b}(\bar{U}, t)dt + \bar{\sigma}(\bar{U}, t)dW, \quad 0 \leq t < T.$$

²U. G. Haussmann, É. Pardoux. Time reversal of diffusions. *Ann. Probab.* **14**(4), 1188-1205 (1986).

Time reversal of diffusions

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

Discretized forward diffusion $\mathbf{U}_{i+1} = \mathbf{U}_i + \mathbf{b}_i \Delta t + \sigma_i \sqrt{\Delta t} \mathbf{Z}$, $\mathbf{Z} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$,
from which one deduces:

$$\begin{aligned}\mathbb{P}(\mathbf{u}_i | \mathbf{u}_{i+1}) &= \frac{\mathbb{P}(\mathbf{u}_{i+1} | \mathbf{u}_i) \mathbb{P}(\mathbf{u}_i)}{\mathbb{P}(\mathbf{u}_{i+1})} \\ &= \frac{\mathbb{P}(\mathbf{u}_i)}{\mathbb{P}(\mathbf{u}_{i+1})} \mathcal{N}(\mathbf{u}_{i+1}; \mathbf{u}_i + \mathbf{b}_i \Delta t, \sigma_i^2 \Delta t); \end{aligned}$$

But:

$$\begin{aligned}\mathbb{P}(\mathbf{u}_i) &= \mathbb{P}(\mathbf{u}_{i+1}) + (\mathbf{u}_i - \mathbf{u}_{i+1}) \cdot \nabla_{\mathbf{u}} \mathbb{P}(\mathbf{u}_{i+1}) + \dots \\ \frac{\mathbb{P}(\mathbf{u}_i)}{\mathbb{P}(\mathbf{u}_{i+1})} &= 1 + (\mathbf{u}_i - \mathbf{u}_{i+1}) \cdot \nabla_{\mathbf{u}} \log \mathbb{P}(\mathbf{u}_{i+1}) + \dots \\ &\simeq \exp [(\mathbf{u}_i - \mathbf{u}_{i+1}) \cdot \nabla_{\mathbf{u}} \log \mathbb{P}(\mathbf{u}_{i+1})], \end{aligned}$$

from which one deduces:

$$\begin{aligned}\mathbb{P}(\mathbf{u}_i | \mathbf{u}_{i+1}) &\propto \exp \left[(\mathbf{u}_i - \mathbf{u}_{i+1}) \cdot \nabla_{\mathbf{u}} \log \mathbb{P}(\mathbf{u}_{i+1}) - \frac{\|\mathbf{u}_{i+1} - \mathbf{u}_i - \mathbf{b}_i \Delta t\|^2}{2\sigma_i^2 \Delta t} \right] \\ &= \exp \left[- \frac{\|\mathbf{u}_i - (\mathbf{u}_{i+1} - \mathbf{b}_i \Delta t + \sigma_i^2 \nabla_{\mathbf{u}} \log \mathbb{P}(\mathbf{u}_{i+1}) \Delta t)\|^2}{2\sigma_i^2 \Delta t} \right]. \end{aligned}$$

Probability flow ODE

SDE

W. Genuist,
É. Savin

SDE

First-order
systems

Stochastic
integrals

Diffusion
processes

Numerical
solutions

Stochastic
modeling

Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case

DDPM

SMLD

- Since $\nabla_{\mathbf{u}} \cdot (\pi \mathbf{a}) = \pi \nabla_{\mathbf{u}} \cdot \mathbf{a} + \mathbf{a} \nabla_{\mathbf{u}} \pi$, the Fokker-Planck equation also reads:

$$\partial_t \pi + \nabla_{\mathbf{u}} \cdot (\pi \mathbf{b}^\dagger) = 0,$$

with:

$$\mathbf{b}^\dagger(\mathbf{u}, t) = \mathbf{b}(\mathbf{u}, t) - \frac{1}{2} \frac{\nabla_{\mathbf{u}} \cdot (\pi(\mathbf{u}, t) \mathbf{a}(\mathbf{u}, t))}{\pi(\mathbf{u}, t)}.$$

- It is to the Fokker-Planck equation associated to the SDE:

$$d\mathbf{U} = \mathbf{b}^\dagger(\mathbf{U}, t) dt, \quad \mathbf{U}(0) = \mathbf{U}_0,$$

which is coined "probability flow ODE" (Ordinary Differential Equation).

Feynman-Kac formula

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- Consider the *backward* PDE with final condition at $T > 0$ and (scalar) solution $\mathbf{x}, t \mapsto u(\mathbf{x}, t)$:

$$\partial_t u + \mathbf{b}(\mathbf{x}) \cdot \nabla_{\mathbf{x}} u + \frac{1}{2} \boldsymbol{\sigma}(\mathbf{x}) \boldsymbol{\sigma}(\mathbf{x})^\top : \nabla_{\mathbf{x}} \otimes \nabla_{\mathbf{x}} u = 0, \\ u(\mathbf{x}, T) = \phi(\mathbf{x});$$

- Consider the process \mathbf{X}_t on $[\tau, T]$ solving:

$$d\mathbf{X}_t = \mathbf{b}(\mathbf{X}_t)dt + \boldsymbol{\sigma}(\mathbf{X}_t)d\mathbf{W}_t;$$

- Then:

$$u(\boldsymbol{\xi}, \tau) = \mathbb{E}\{\phi(\mathbf{X}_T) | \mathbf{X}_\tau = \boldsymbol{\xi}\}.$$

Feynman-Kac formula

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- Consider the *forward* PDE with initial condition at $t = 0$ and (scalar) solution $\mathbf{x}, t \mapsto u(\mathbf{x}, t)$:

$$\begin{aligned}\partial_t u &= \mathbf{b}(\mathbf{x}) \cdot \nabla_{\mathbf{x}} u + \frac{1}{2} \boldsymbol{\sigma}(\mathbf{x}) \boldsymbol{\sigma}(\mathbf{x})^\top : \nabla_{\mathbf{x}} \otimes \nabla_{\mathbf{x}} u, \\ u(\mathbf{x}, 0) &= \phi(\mathbf{x});\end{aligned}$$

- Consider the process \mathbf{X}_t on $[0, \tau]$ solving:

$$d\mathbf{X}_t = \mathbf{b}(\mathbf{X}_t)dt + \boldsymbol{\sigma}(\mathbf{X}_t)d\mathbf{W}_t;$$

- Then:

$$u(\boldsymbol{\xi}, \tau) = \mathbb{E}\{\phi(\mathbf{X}_\tau) | \mathbf{X}_0 = \boldsymbol{\xi}\}.$$

Feynman-Kac formula

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion & ML

Continuous
case
DDPM
SMLD

Backward case. Using Itô's formula for $u(\mathbf{X}_t, t)$:

$$\begin{aligned} du &= (\partial_t u)dt + \nabla_{\mathbf{x}} u \cdot d\mathbf{X}_t + \frac{1}{2} \nabla_{\mathbf{x}} u \otimes \nabla_{\mathbf{x}} u : d\mathbf{X}_t \otimes d\mathbf{X}_t \\ &= \left(\partial_t u + \nabla_{\mathbf{x}} u \cdot \mathbf{b}(\mathbf{X}_t) + \frac{1}{2} \nabla_{\mathbf{x}} u \otimes \nabla_{\mathbf{x}} u : \boldsymbol{\sigma}(\mathbf{X}_t) \boldsymbol{\sigma}(\mathbf{X}_t)^\top \right) dt + \nabla_{\mathbf{x}} u \cdot \boldsymbol{\sigma}(\mathbf{X}_t) dW_t \\ &= \nabla_{\mathbf{x}} u \cdot \boldsymbol{\sigma}(\mathbf{X}_t) dW_t ; \end{aligned}$$

Integrating between τ and T :

$$u(\mathbf{X}_T, T) - u(\mathbf{X}_\tau, \tau) = \phi(\mathbf{X}_T) - u(\mathbf{X}_\tau, \tau) = \int_\tau^T \nabla_{\mathbf{x}} u \cdot \boldsymbol{\sigma}(\mathbf{X}_s) dW_s ;$$

Taking the expectation the right-hand side vanishes and then:

$$u(\boldsymbol{\xi}, \tau) = \mathbb{E}\{\phi(\mathbf{X}_T) | \mathbf{X}_\tau = \boldsymbol{\xi}\} .$$

Forward case. Change of variable $t \rightarrow T - t$.

Outline

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion & ML

Continuous
case
DDPM
SMLD

- 1 Stochastic differential equations (SDE)
 - First-order stochastic systems driven by noise
 - Stochastic integrals
 - Diffusion processes
- 2 Numerical simulations of SDE
 - Stochastic modeling with SDE
 - Numerical schemes
- 3 Generative Adversarial Networks (GAN)
- 4 Variational Auto-Encoders (VAE)
- 5 Diffusion models in ML
 - Continuous setting
 - Denoising Diffusion Probabilistic Modeling (DDPM)
 - Score Matching with Langevin Dynamics (SMLD)

First-order stochastic differential equation

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

A general first-order stochastic differential equation for the process U indexed on \mathbb{R}_+ with values in \mathbb{R}^q :

$$\begin{cases} \dot{U}(t) = \mathbf{b}(U, t) + \sigma(U, t)\mathbf{F}(t), & t > 0, \\ U(0) = U_0, \end{cases}$$

with the data:

- $\mathbf{u}, t \mapsto \mathbf{b}(\mathbf{u}, t) : \mathbb{R}^q \times \mathbb{R}_+ \rightarrow \mathbb{R}^q$ the *drift* function;
- $\mathbf{u}, t \mapsto \sigma(\mathbf{u}, t) : \mathbb{R}^q \times \mathbb{R}_+ \rightarrow \mathbb{M}_{q,p}(\mathbb{R})$ the *scattering* operator;
- U_0 is an r.v. in \mathbb{R}^q with known marginal PDF $\pi_0(\mathbf{u}_0)$;
- $\mathbf{F}(t) = (F_1(t), \dots, F_p(t))$ is a second-order Gaussian random process indexed on \mathbb{R}^+ with values in \mathbb{R}^p , centered, mean-square continuous.

Markovian realization

Definition

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

Definition

$\mathbf{F}(t)$ indexed on \mathbb{R}^+ with values in \mathbb{R}^p , second-order, Gaussian, centered and mean-square continuous admits a Markovian realization if:

$$\left\{ \begin{array}{ll} \mathbf{F}(t) = \mathbf{H}\mathbf{V}(t), & t \geq 0, \\ \dot{\mathbf{V}}(t) = \mathbf{P}\mathbf{V}(t) + \mathbf{Q}\mathbf{B}(t), & t > 0, \\ \mathbf{V}(0) = \mathbf{V}_0 & a.s. \end{array} \right.$$

where \mathbf{V}_0 is a Gaussian r.v. in \mathbb{R}^n , $\mathbf{V}(t)$ is a diffusion process indexed on \mathbb{R}_+ with values in \mathbb{R}^n , $\mathbf{P}, \mathbf{Q} \in \mathbb{M}_n(\mathbb{R})$, $\mathbf{H} \in \mathbb{M}_{p,n}(\mathbb{R})$, $\Re\{\lambda_j(\mathbf{P})\} < 0$.

- This is equivalent to a linear Itô stochastic differential equation.
- $\mathbf{V}_0 \sim \mathcal{N}(\mathbf{0}, \mathbf{\Sigma}_0)$ where $\mathbf{\Sigma}_0 = \int_0^{+\infty} e^{\tau\mathbf{P}} \mathbf{Q}\mathbf{Q}^\top e^{\tau\mathbf{P}^\top} d\tau$.

Physically realizable process

Definition

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

Definition

$\mathbf{F}(t)$ indexed on \mathbb{R} with values in \mathbb{R}^p , second-order, mean-square stationary and continuous, centered, is physically realizable if $\exists \mathbb{H} \in L^2(\mathbb{R})$, $\text{supp } \mathbb{H} \subseteq \mathbb{R}_+$, such that:

$$\mathbf{F}(t) = \int_{-\infty}^t \mathbb{H}(t - \tau) \mathbf{B}(\tau) d\tau,$$

or equivalently $\mathbf{S}_{\mathbf{F}}(\omega) = \frac{1}{2\pi} \hat{\mathbb{H}}(\omega) \hat{\mathbb{H}}(\omega)^, \forall \omega \in \mathbb{R}$.*

A necessary and sufficient condition (Rozanov 1967):

$$\int_{\mathbb{R}} \frac{\ln(\det \mathbf{S}_{\mathbf{F}}(\omega))}{1 + \omega^2} d\omega > -\infty.$$

Markovian realization

Existence for a physically realizable process

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

Theorem

A necessary and sufficient condition:

$$\mathbf{S}_{\mathbf{F}}(\omega) = \frac{\mathbf{R}(\mathrm{i}\omega)\mathbf{R}(\mathrm{i}\omega)^*}{2\pi|P(\mathrm{i}\omega)|^2}, \quad \text{or} \quad \mathbb{H}(\omega) = \frac{\mathbf{R}(\mathrm{i}\omega)}{P(\mathrm{i}\omega)},$$

where:

- $P(z)$ is a polynomial of degree d on \mathbb{C} with real coefficients and roots in the half-plane $\Re(z) < 0$,
- $\mathbf{R}(z)$ is a polynomial on \mathbb{C} with coefficients in $\mathbb{M}_{p,n}(\mathbb{R})$ and degree $r < n$.

The Markovian realization always exists in infinite dimension $n = +\infty$.

First-order SDE (cont'd)

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

A non linear first-order stochastic differential equation for the process $\mathbf{Z}(t) = (\mathbf{U}(t), \mathbf{V}(t))$ indexed on \mathbb{R}_+ with values in \mathbb{R}^ν , $\nu = q + n$:

$$\begin{cases} d\mathbf{Z}(t) = \mathbf{b}_z(\mathbf{Z}, t)dt + \boldsymbol{\sigma}_z d\mathbf{W}, & t > 0, \\ \mathbf{Z}(0) = \mathbf{Z}_0, \end{cases}$$

where $\mathbf{Z}_0 = (\mathbf{U}_0, \mathbf{V}_0)$,

$$\mathbf{b}_z(\mathbf{u}, \mathbf{v}, t) = \begin{bmatrix} \mathbf{b}(\mathbf{u}, t) + \boldsymbol{\sigma}(\mathbf{u}, t)\mathbf{H}\mathbf{v} \\ \mathbf{P}\mathbf{v} \end{bmatrix}, \quad \boldsymbol{\sigma}_z = \begin{bmatrix} \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{Q} \end{bmatrix},$$

and $\mathbf{W}(t)$ is the Wiener process in \mathbb{R}^ν .

Numerical integration of SDE

Strong convergence

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

$$\begin{cases} dU(t) = b(U, t)dt + \sigma(U, t)dW(t), & t > 0, \\ U(0) = U_0 & \text{a.s.} \end{cases}$$

Definition

An approximation $(\tilde{U}_j)_j$ converges with strong order $k > 0$ if $\exists K_j > 0$:

$$\mathbb{E} \left\{ \left| U(j\Delta t) - \tilde{U}_j \right| \right\} \leq K_j (\Delta t)^k.$$

The sample paths of the approximation \tilde{U} should be close to those of the Itô process.

Numerical integration of SDE

Weak convergence

SDE

W. Genuist,
É. Savin

$$\begin{cases} dU(t) = b(U, t)dt + \sigma(U, t)dW(t), & t > 0, \\ U(0) = U_0 & \text{a.s.} \end{cases}$$

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

Definition

An approximation $(\tilde{U}_j)_j$ converges with weak order $k > 0$ if for any polynomial $g \exists K_{g,j} > 0$:

$$\left| \mathbb{E} \{g(U(j\Delta t))\} - \mathbb{E} \{g(\tilde{U}_j)\} \right| \leq K_{g,j}(\Delta t)^k.$$

The probability distribution of the approximation should be close to that of the Itô process in order to get a good estimate of the expectation ($g(u) = u$) or the variance ($g(u) = u^2$), for example.

Time discrete approximations

Explicit 0.5-order methods

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion & ML

Continuous
case
DDPM
SMLD

- Assume that v and a are independent of time t (thus $U(t)$ is a diffusion process), and let $t_j = j\Delta t$, $b_j = b(\tilde{U}_j)$, $\sigma_j = \sigma(\tilde{U}_j)$, $U_0 \sim \pi_0(du_0)$, $G \sim \mathcal{N}(0, 1)$.

- Itô SDE: the *Euler-Maruyama scheme* (1955),

$$\begin{aligned}\tilde{U}_{j+1} &= \tilde{U}_j + b_j \Delta t + \sigma_j \sqrt{\Delta t} G, \\ \tilde{U}_0 &= U_0.\end{aligned}$$

- Stratonovich SDE: the *Euler-Heun scheme* (1982),

$$\begin{aligned}\tilde{U}_{j+1} &= \tilde{U}_j + b_j \Delta t + \tilde{\sigma}_j \sqrt{\Delta t} G, \\ \tilde{\sigma}_j &= \frac{1}{2} \left[\sigma_j + \sigma \left(\tilde{U}_j + \sigma_j \sqrt{\Delta t} G \right) \right], \\ \tilde{U}_0 &= U_0.\end{aligned}$$

- Both have a strong order $k = \frac{1}{2}$ (vs. $k = 1$ for ordinary differential equations) and a weak order $k = 1$.

Time discrete approximations

Explicit 1-order methods

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- The *Milstein scheme* (1974):

$$\tilde{U}_{j+1} = \tilde{U}_j + b_{\lambda,j}\Delta t + \sigma_j\sqrt{\Delta t}G + \frac{1}{2}\sigma_j\sigma'_j\Delta t(G^2 + 2\lambda - 1),$$

$$\tilde{U}_0 = U_0,$$

where $\lambda = 0$ (Itô SDE) or $\lambda = \frac{1}{2}$ (Stratonovich SDE).

- The *Runge-Kutta Milstein scheme* (1984):

$$\tilde{U}_{j+1} = \tilde{U}_j + b_{\lambda,j}\Delta t + \sigma_j\sqrt{\Delta t}G + \frac{1}{2}\sigma_j\tilde{\sigma}'_j\Delta t(G^2 + 2\lambda - 1),$$

$$\sigma_j\tilde{\sigma}'_j = (\Delta t)^{-\frac{1}{2}} \left[\sigma \left(\tilde{U}_j + \sigma_j\sqrt{\Delta t} \right) - \sigma_j \right],$$

$$\tilde{U}_0 = U_0.$$

- Both have strong and weak orders $k = 1$ (under mild conditions on b and σ).

Time discrete approximations

Stochastic Taylor approximations

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- Higher-order schemes may be derived using stochastic Taylor expansions:

$$\begin{aligned} U_{j+1} - U_j &= \int_{t_j}^{t_{j+1}} b(U) dt + \int_{t_j}^{t_{j+1}} \sigma(U) dW \\ &\simeq \int_{t_j}^{t_{j+1}} (b(U_j) + b'(U_j) \Delta U_j) dt + \int_{t_j}^{t_{j+1}} (\sigma(U_j) + \sigma'(U_j) \Delta U_j) dW, \end{aligned}$$

where $\Delta U_j = \int_{t_j}^t b(U) d\tau + \int_{t_j}^t \sigma(U) dW$.

- Then $\int_{t_j}^{t_{j+1}} \int_{t_j}^t d_\lambda W(s) d_\lambda W(t) = \frac{1}{2}(\Delta W)^2 + (\lambda - \frac{1}{2})\Delta t$.
- Higher-order expansions involve additional r.v.
 $\Delta Z_j = \int_{t_j}^{t_{j+1}} \int_{t_j}^t dW dt$ with $\mathbb{E}\{(\Delta Z_j)^2\} \propto \Delta t^3$ etc.
- Weak Taylor approximations $U_0 \sim \hat{U}_0$, $\Delta W \sim \Delta \hat{W}$,
 $\Delta Z_j \sim \Delta \hat{Z}_j$ with approximately the same moment properties.

Outline

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion & ML

Continuous
case
DDPM
SMLD

- 1 Stochastic differential equations (SDE)
 - First-order stochastic systems driven by noise
 - Stochastic integrals
 - Diffusion processes
- 2 Numerical simulations of SDE
 - Stochastic modeling with SDE
 - Numerical schemes
- 3 Generative Adversarial Networks (GAN)
- 4 Variational Auto-Encoders (VAE)
- 5 Diffusion models in ML
 - Continuous setting
 - Denoising Diffusion Probabilistic Modeling (DDPM)
 - Score Matching with Langevin Dynamics (SMLD)

Setting

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical solutions

Stochastic
modeling
Numerical
schemes


GAN

VAE

Diffusion & ML

Continuous
case
DDPM
SMLD

- *Unsupervised learning*: infer $\mathbf{X}^* \sim \mathbb{P}(\mathbf{x})$ where $\mathbb{P}(\mathbf{x})$ is only partially known through the dataset $\{\mathbf{x}_n\}_{n=1}^N$;
- *Discriminator*: $\mathbb{R}^q \rightarrow \mathbb{R} : \mathbf{x} \mapsto d_\phi(\mathbf{x})$;
- *Generator*: $\mathbb{R}^p \rightarrow \mathbb{R}^q : \mathbf{z} \mapsto \mathbf{g}_\theta(\mathbf{z})$ with $p \ll q$.



Training and inference

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE


Diffusion &
ML

Continuous
case
DDPM
SMLD

- **Loss function:** $\mathbf{Z}_i \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$, say, and

$$\theta^*, \phi^* = \arg \min_{\theta} \max_{\phi} \frac{1}{N} \sum_{n=1}^N [d_{\phi}(\mathbf{x}_i) - d_{\phi}(\mathbf{g}_{\theta}(\mathbf{Z}_i))] \\ (+ \text{gradient penalty});$$

- **Inference:** $\mathbf{X}^* = \mathbf{g}_{\theta^*}(\mathbf{Z})$ where $\mathbf{Z} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$, say.



Outline

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- 1 Stochastic differential equations (SDE)
 - First-order stochastic systems driven by noise
 - Stochastic integrals
 - Diffusion processes
- 2 Numerical simulations of SDE
 - Stochastic modeling with SDE
 - Numerical schemes
- 3 Generative Adversarial Networks (GAN)
- 4 Variational Auto-Encoders (VAE)
- 5 Diffusion models in ML
 - Continuous setting
 - Denoising Diffusion Probabilistic Modeling (DDPM)
 - Score Matching with Langevin Dynamics (SMLD)

Setting

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion & ML

Continuous
case
DDPM
SMLD

- *Unsupervised learning*: infer $\mathbf{X}^* \sim \mathbb{P}(\mathbf{x})$ where $\mathbb{P}(\mathbf{x})$ is only partially known through the dataset $\{\mathbf{x}_n\}_{n=1}^N$;
- *Encoder* (forward relationship $\mathbb{R}^q \rightarrow \mathbb{R}^p$ with $p \ll q$):

$$\mathbf{Z} = \text{encoder}_{\phi}(\mathbf{X}) \sim \mathbb{Q}_{\phi}(\mathbf{z}|\mathbf{x}) \approx \mathbb{P}(\mathbf{z}|\mathbf{x});$$

- *Decoder* (backward relationship $\mathbb{R}^p \rightarrow \mathbb{R}^q$):

$$\mathbf{X} = \text{decoder}_{\theta}(\mathbf{Z}) \sim \mathbb{P}_{\theta}(\mathbf{x}|\mathbf{z}) \approx \mathbb{P}(\mathbf{x}|\mathbf{z});$$

- The *latent variable* is e.g. $\mathbf{Z} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$.
- *Example*: in **jpeg** the encoder is the discrete cosine transform, the decoder is its inverse, and \mathbf{z} are the projection coefficients.

Evidence

SDE

- Straightforward inference is $\mathbf{X}^* = \arg \max \mathbb{P}(\mathbf{x})$ but:

$$\begin{aligned}\log \mathbb{P}(\mathbf{x}) &= \log \frac{\mathbb{P}(\mathbf{x}, \mathbf{z})}{\mathbb{P}(\mathbf{z}|\mathbf{x})} \\ &= \log \frac{\mathbb{P}(\mathbf{x}|\mathbf{z})\mathbb{P}(\mathbf{z})\mathbb{Q}_{\phi}(\mathbf{z}|\mathbf{x})}{\mathbb{P}(\mathbf{z}|\mathbf{x})\mathbb{Q}_{\phi}(\mathbf{z}|\mathbf{x})} \\ &= \log \mathbb{P}(\mathbf{x}|\mathbf{z}) - \log \frac{\mathbb{Q}_{\phi}(\mathbf{z}|\mathbf{x})}{\mathbb{P}(\mathbf{z})} + \log \frac{\mathbb{Q}_{\phi}(\mathbf{z}|\mathbf{x})}{\mathbb{P}(\mathbf{z}|\mathbf{x})}\end{aligned}$$

where $\mathbb{Q}_{\phi}(\mathbf{z}|\mathbf{x})$ could actually be anything else.

- Then (remind $\mathbb{D}_{\text{KL}}(\mathbb{P}||\mathbb{Q}) \stackrel{\text{def}}{=} \mathbb{E}_{\mathbb{P}}\{\log \frac{\mathbb{P}}{\mathbb{Q}}\} \geq 0$):

$$\begin{aligned}\log \mathbb{P}(\mathbf{x}) &= \mathbb{E}_{\mathbb{Q}_{\phi}(\mathbf{z}|\mathbf{x})}\{\log \mathbb{P}(\mathbf{x})\} \\ &= \mathbb{E}_{\mathbb{Q}_{\phi}(\mathbf{z}|\mathbf{x})}\{\log \mathbb{P}(\mathbf{x}|\mathbf{Z})\} - \mathbb{D}_{\text{KL}}(\mathbb{Q}_{\phi}(\mathbf{z}|\mathbf{x})||\mathbb{P}(\mathbf{z})) + \mathbb{D}_{\text{KL}}(\mathbb{Q}_{\phi}(\mathbf{z}|\mathbf{x})||\mathbb{P}(\mathbf{z}|\mathbf{x})) \\ &\geq \mathbb{E}_{\mathbb{Q}_{\phi}(\mathbf{z}|\mathbf{x})}\{\log \mathbb{P}(\mathbf{x}|\mathbf{Z})\} - \mathbb{D}_{\text{KL}}(\mathbb{Q}_{\phi}(\mathbf{z}|\mathbf{x})||\mathbb{P}(\mathbf{z})) \\ &\simeq \mathbb{E}_{\mathbb{Q}_{\phi}(\mathbf{z}|\mathbf{x})}\{\log \mathbb{P}_{\theta}(\mathbf{x}|\mathbf{Z})\} - \mathbb{D}_{\text{KL}}(\mathbb{Q}_{\phi}(\mathbf{z}|\mathbf{x})||\mathbb{P}(\mathbf{z})).\end{aligned}$$

Evidence Lower BOund (ELBO)

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- The *Evidence Lower BOund* (ELBO), or *variational bound*, is:

$$\begin{aligned}\text{ELBO}(\mathbf{x}; \boldsymbol{\theta}, \phi) &\stackrel{\text{def}}{=} \mathbb{E}_{\mathbb{Q}_{\phi}(\mathbf{z}|\mathbf{x})} \left\{ \log \frac{\mathbb{P}_{\boldsymbol{\theta}}(\mathbf{x}, \mathbf{Z})}{\mathbb{Q}_{\phi}(\mathbf{Z}|\mathbf{x})} \right\} \\ &= \mathbb{E}_{\mathbb{Q}_{\phi}(\mathbf{z}|\mathbf{x})} \{ \log \mathbb{P}_{\boldsymbol{\theta}}(\mathbf{x}|\mathbf{Z}) \} - \mathbb{D}_{\text{KL}}(\mathbb{Q}_{\phi}(\mathbf{z}|\mathbf{x}) || \mathbb{P}(\mathbf{z}));\end{aligned}$$

- Sampling $\mathbf{Z} \sim \mathbb{Q}_{\phi}(\mathbf{z}|\mathbf{x})$ and maximizing $\text{ELBO}(\mathbf{x}; \boldsymbol{\theta}, \phi)$, we almost do maximize $\log \mathbb{P}(\mathbf{x})$ on average;
- *Prior matching*: $\mathbb{D}_{\text{KL}}(\mathbb{Q}_{\phi}(\mathbf{z}|\mathbf{x}) || \mathbb{P}(\mathbf{z}))$ tells how good the encoder is *vs.* a prior belief held over latent variables, and has to be minimized to maximize the ELBO;
- *Reconstruction*: $\mathbb{E}_{\mathbb{Q}_{\phi}(\mathbf{z}|\mathbf{x})} \{ \log \mathbb{P}_{\boldsymbol{\theta}}(\mathbf{x}|\mathbf{Z}) \}$ tells how good the decoder is, and has to be maximized to maximize the ELBO.

Training and inference

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- **Encoder:** we choose $\mathbb{Q}_\phi(z|\mathbf{x}) = \mathcal{N}(z; \boldsymbol{\mu}_\phi(\mathbf{x}), \sigma_\phi^2(\mathbf{x})\mathbf{I})$, where $\boldsymbol{\mu}_\phi(\mathbf{x})$ and $\sigma_\phi^2(\mathbf{x})$ are Neural Networks;
- **Decoder:** we choose $\mathbb{P}_\theta(\mathbf{x}|z) = \mathcal{N}(\mathbf{x}; \boldsymbol{\mu}_\theta(z), \sigma^2\mathbf{I})$, where $\boldsymbol{\mu}_\theta(z)$ is a Neural Network and σ is a parameter;
- **Loss function:**

$$\boldsymbol{\theta}^*, \phi^* = \arg \max_{\boldsymbol{\theta}, \phi} \frac{1}{N} \sum_{n=1}^N (\log \mathbb{P}_\theta(\mathbf{x}_n | z_n) - \mathbb{D}_{\text{KL}}(\mathbb{Q}_\phi(z | \mathbf{x}_n) || \mathbb{P}(z))) ,$$

where $z_n \sim \mathbb{Q}_\phi(z | \mathbf{x}_n)$, and for *e.g.* $\mathbb{P}(z) = \mathcal{N}(\mathbf{0}, \mathbf{I})$ in a latent space of dimension p :

$$\mathbb{D}_{\text{KL}}(\mathbb{Q}_\phi(z | \mathbf{x}) || \mathbb{P}(z)) = \frac{1}{2} \left(\|\boldsymbol{\mu}_\phi(\mathbf{x})\|^2 + \sigma_\phi^{2p}(\mathbf{x}) - p \log \sigma_\phi^2(\mathbf{x}) \right) ;$$

- **Inference:** $\mathbf{X}^* = \boldsymbol{\mu}_\theta^*(\mathbf{Z})$ where $\mathbf{Z} \sim \mathbb{P}(z)$.

Outline

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion & ML

Continuous
case
DDPM
SMLD

- 1 Stochastic differential equations (SDE)
 - First-order stochastic systems driven by noise
 - Stochastic integrals
 - Diffusion processes
- 2 Numerical simulations of SDE
 - Stochastic modeling with SDE
 - Numerical schemes
- 3 Generative Adversarial Networks (GAN)
- 4 Variational Auto-Encoders (VAE)
- 5 Diffusion models in ML
 - Continuous setting
 - Denoising Diffusion Probabilistic Modeling (DDPM)
 - Score Matching with Langevin Dynamics (SMLD)

Unsupervised learning by noising/denoising

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions



Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD



© J. Ho, A. Jain, P. Abbeel

Langevin diffusion

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- Langevin diffusion:

$$d\mathbf{X}_t = -\nabla_{\mathbf{x}}\mathcal{U}(\mathbf{X}_t)dt + \sqrt{2}d\mathbf{W}_t, \quad \mathbb{R}^p \ni \mathbf{X}_0 \sim \pi_0.$$

- Regardless of π_0 , \mathbf{X}_t converges in law towards a density $\propto e^{-\mathcal{U}(\mathbf{x})}$ as $t \rightarrow \infty$. The Fokker-Planck equation reads:

$$\partial_t \pi = \nabla_{\mathbf{x}} \cdot (\pi \nabla_{\mathbf{x}} \mathcal{U}) + \Delta \pi.$$

- **Example:** Ornstein-Uhlenbeck process, $\mathcal{U}(\mathbf{x}) = \frac{1}{2} \|\mathbf{x}\|^2$, thus

$$d\mathbf{X}_t = -\mathbf{X}_t dt + \sqrt{2}d\mathbf{W}_t, \quad \mathbf{X}_0 \sim \pi_0,$$

and

$$\partial_t \pi = \nabla_{\mathbf{x}} \cdot (\pi \mathbf{x}) + \Delta \pi.$$

Ornstein-Uhlenbeck process

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- In integral form:

$$\mathbf{X}_t = e^{-t} \left(\mathbf{X}_0 + \sqrt{2} \int_0^t e^s d\mathbf{W}_s \right),$$

with ground-truth density $\pi = \pi_0(\cdot/e^t) \star \mathcal{N}(0, 1 - e^{-2t})$.

Let $\mathbf{Y}_t = e^t \mathbf{X}_t$. Then by Itô's formula with $\phi(x, t) = e^t x$:

$$\begin{aligned} d\mathbf{Y}_t &= (\partial_t \phi) dt + (D_x \phi) d\mathbf{X}_t + \frac{1}{2} (D_x^2 \phi) d\mathbf{X}_t \otimes d\mathbf{X}_t \\ &= e^t \mathbf{X}_t dt + e^t d\mathbf{X}_t \quad (\partial_t \phi = \phi, D_x \phi = e^t I, D_x^2 \phi = 0) \\ &= e^t \mathbf{X}_t dt + e^t (-\mathbf{X}_t dt + \sqrt{2} d\mathbf{W}_t) \\ &= \sqrt{2} e^t d\mathbf{W}_t. \end{aligned}$$

Thus:

$$\mathbf{Y}_t = \mathbf{X}_0 + \sqrt{2} \int_0^t e^s d\mathbf{W}_s.$$

Ornstein-Uhlenbeck process

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- It is a Gaussian process with conditional moments:

$$\mathbb{E}\{\mathbf{X}_t|\mathbf{X}_0\} = e^{-t}\mathbf{X}_0,$$

$$\mathbb{V}\{\mathbf{X}_t|\mathbf{X}_0\} = (1 - e^{-2t})\mathbf{I}.$$

$$\begin{aligned}\mathbb{E}\{(\mathbf{X}_t - \mathbb{E}\{\mathbf{X}_t\}) \otimes (\mathbf{X}_{t'} - \mathbb{E}\{\mathbf{X}_{t'}\})|\mathbf{X}_0\} \\ = 2e^{-t-t'}\mathbb{E}\left\{\left(\int_0^t e^s d\mathbf{W}_s\right) \otimes \left(\int_0^{t'} e^s d\mathbf{W}_s\right)\right\}.\end{aligned}$$

Then apply Itô's isometry formula:

$$\mathbb{E}\left\{\left(\int_0^t Y_s d\mathbf{W}_s\right) \otimes \left(\int_0^{t'} Z_s d\mathbf{W}_s\right)\right\} = \mathbb{E}\left\{\int_0^{\min(t,t')} Y_s Z_s ds\right\} \mathbf{I}$$

to obtain:

$$\mathbb{E}\left\{\left(\int_0^t e^s d\mathbf{W}_s\right) \otimes \left(\int_0^{t'} e^s d\mathbf{W}_s\right)\right\} = \mathbb{E}\left\{\int_0^{\min(t,t')} e^{2s} ds\right\} = \frac{1}{2} \left(e^{2\min(t,t')} - 1\right).$$

Backward Ornstein-Uhlenbeck process

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- Let $\rho(\mathbf{x}, t) = \pi(\mathbf{x}, T - t)$ starting from some large enough $T \gg 0$, it satisfies the Fokker-Planck equation:

$$\partial_t \rho = -\partial_t \pi = -\nabla_{\mathbf{x}} \cdot (\rho \mathbf{x}) - \Delta \rho.$$

- But:

$$-\Delta \rho = \Delta \rho - 2\nabla \cdot (\rho \nabla \log \rho),$$

such that the Fokker-Planck equation also reads:

$$\partial_t \rho = -\nabla_{\mathbf{x}} \cdot (\rho \mathbf{x} + 2\rho \nabla_{\mathbf{x}} \log \rho) + \Delta \rho.$$

- This is the law density of the process \mathbf{Y}_t that follows backward Langevin diffusion:

$$d\mathbf{Y}_t = (\mathbf{Y}_t + 2\nabla_{\mathbf{x}} \log \rho(\mathbf{Y}_t, t))dt + \sqrt{2}d\mathbf{W}_t,$$

starting from $\mathbf{Y}_0 = \mathbf{X}_T \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$.

Score function

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- The *score function*:

$$\mathbf{s}(\mathbf{x}, t) = \nabla_{\mathbf{x}} \log \pi(\mathbf{x}, t) = \nabla_{\mathbf{x}} \log \rho(\mathbf{x}, T - t)$$

is approximated from samples of \mathbf{X}_t from the forward flow, $\mathbf{s}(\mathbf{x}, t) \approx \mathbf{s}_{\theta}(\mathbf{x}, t)$.

- For Ornstein-Uhlenbeck $\mathbf{X}_t \sim e^{-t} \mathbf{X}_0 + \sqrt{1 - e^{-2t}} \mathbf{Z}$, $\mathbf{Z} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$, then setting $\mathbb{P}_{\mathbf{X}_t | \mathbf{X}_0}(\mathbf{x} | \mathbf{x}_0) \stackrel{\text{def}}{=} \mathbb{P}(\mathbf{x} | \mathbf{x}_0; t)$:

$$\begin{aligned} \nabla_{\mathbf{x}} \log \mathbb{P}(\mathbf{x} | \mathbf{x}_0; t) &= -\frac{\mathbf{x} - e^{-t} \mathbf{x}_0}{1 - e^{-2t}} \\ &= \nabla_{\mathbf{x}} \log \frac{\mathbb{P}(\mathbf{x}, \mathbf{x}_0; t)}{\mathbb{P}(\mathbf{x}_0)} \\ &= \frac{\nabla_{\mathbf{x}} \mathbb{P}(\mathbf{x}, \mathbf{x}_0; t)}{\mathbb{P}(\mathbf{x}, \mathbf{x}_0; t)}. \end{aligned}$$

Denoising score matching

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- Since $\mathbb{P}(\mathbf{x}; t) = \int \mathbb{P}(\mathbf{x}, \mathbf{x}_0; t) d\mathbf{x}_0$ one has:

$$\begin{aligned}\nabla_{\mathbf{x}} \log \mathbb{P}(\mathbf{x}; t) &= \frac{\nabla_{\mathbf{x}} \mathbb{P}(\mathbf{x}; t)}{\mathbb{P}(\mathbf{x}; t)} \\&= \frac{1}{\mathbb{P}(\mathbf{x}; t)} \int \nabla_{\mathbf{x}} \mathbb{P}(\mathbf{x}, \mathbf{x}_0; t) d\mathbf{x}_0 \\&= \frac{1}{\mathbb{P}(\mathbf{x}; t)} \int \mathbb{P}(\mathbf{x}, \mathbf{x}_0; t) \nabla_{\mathbf{x}} \log \mathbb{P}(\mathbf{x} | \mathbf{x}_0; t) d\mathbf{x}_0 \quad (\text{previous slide}) \\&= \int \nabla_{\mathbf{x}} \log \mathbb{P}(\mathbf{x} | \mathbf{x}_0; t) d\mathbb{P}(\mathbf{x}_0 | \mathbf{x}; t) \quad \left(\frac{\mathbb{P}(\mathbf{x}, \mathbf{x}_0; t)}{\mathbb{P}(\mathbf{x}; t)} \stackrel{\text{def}}{=} \mathbb{P}(\mathbf{x}_0 | \mathbf{x}; t) \right) \\&= \mathbb{E}_{\mathbf{X}_0 | \mathbf{X}_t} \{ \nabla_{\mathbf{x}} \log \mathbb{P}(\mathbf{x} | \mathbf{x}_0; t) \} \cdot \left(\mathbb{E}_{\mathbf{X}_0 | \mathbf{X}_t} \{ \mathbf{z} \} \stackrel{\text{def}}{=} \int \mathbf{z} d\mathbb{P}(\mathbf{x}_0 | \mathbf{x}; t) \right)\end{aligned}$$

- The score function is $\mathbf{s}(\mathbf{x}, t) \approx \mathbf{s}_{\boldsymbol{\theta}^*}(\mathbf{x}, t)$ where:

$$\boldsymbol{\theta}^* = \arg \min_{\boldsymbol{\theta}} \mathbb{E}_{\mathbf{X}_0} \{ \mathbb{E}_{\mathbf{X}_t | \mathbf{X}_0} \{ \| \mathbf{s}_{\boldsymbol{\theta}}(\mathbf{x}, t) - \nabla_{\mathbf{x}} \log \mathbb{P}(\mathbf{x} | \mathbf{x}_0; t) \|^2 \} \},$$

averaging vs. \mathbf{X}_t ($\mathbb{E}_{\mathbf{X}_t} \mathbb{E}_{\mathbf{X}_0 | \mathbf{X}_t} = \mathbb{E}_{\mathbf{X}_0} \mathbb{E}_{\mathbf{X}_t | \mathbf{X}_0} = \mathbb{E}_{\mathbf{X}_0, \mathbf{X}_t}$).

Denoising score matching with diffusion

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- The score function is $\mathbf{s}(\mathbf{x}, t) \approx \mathbf{s}_{\theta^*}(\mathbf{x}, t)$ where:

$$\theta^* = \arg \min_{\theta} \int_0^T \mathbb{E}_{\mathbf{X}_0, \mathbf{X}_t} \left\{ \left\| \mathbf{s}_{\theta}(\mathbf{x}, t) - \frac{e^{-t} \mathbf{x}_0 - \mathbf{x}}{1 - e^{-2t}} \right\|^2 \right\} \lambda(dt),$$

with some weighting scheme λ w.r.t. time.

- The function $\mathbf{s}_{\theta}(\cdot, t) : \mathbb{R}^p \rightarrow \mathbb{R}^p$ is typically a neural network from \mathbb{R}^p into himself, *e.g.* a U-net for images.

Remark: implicit score matching³

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- Score matching reads:

$$\theta^* = \arg \min_{\theta} J(\theta),$$

where:

$$\begin{aligned} J(\theta) &= \mathbb{E}_{\mathbf{X} \sim \mathbb{P}} \left\{ \frac{1}{2} \|\mathbf{s}_{\theta}(\mathbf{X}) - \nabla_{\mathbf{x}} \log \mathbb{P}(\mathbf{X})\|^2 \right\} \\ &= \int \left(\frac{1}{2} \|\mathbf{s}_{\theta}(\mathbf{x})\|^2 + \frac{1}{2} \|\nabla_{\mathbf{x}} \log \mathbb{P}(\mathbf{x})\|^2 - \mathbf{s}_{\theta}(\mathbf{x}) \cdot \nabla_{\mathbf{x}} \log \mathbb{P}(\mathbf{x}) \right) \mathbb{P}(\mathbf{x}) d\mathbf{x} \\ &= \int \left(\frac{1}{2} \|\mathbf{s}_{\theta}(\mathbf{x})\|^2 - \frac{\mathbf{s}_{\theta}(\mathbf{x}) \cdot \nabla_{\mathbf{x}} \mathbb{P}(\mathbf{x})}{\mathbb{P}(\mathbf{x})} \right) \mathbb{P}(\mathbf{x}) d\mathbf{x} + C \\ &= \int \left(\frac{1}{2} \|\mathbf{s}_{\theta}(\mathbf{x})\|^2 + \nabla_{\mathbf{x}} \cdot \mathbf{s}_{\theta}(\mathbf{x}) \right) \mathbb{P}(\mathbf{x}) d\mathbf{x} + C \\ &= \mathbb{E}_{\mathbf{X} \sim \mathbb{P}} \left\{ \frac{1}{2} \|\mathbf{s}_{\theta}(\mathbf{X})\|^2 + \nabla_{\mathbf{x}} \cdot \mathbf{s}_{\theta}(\mathbf{X}) \right\} + C; \end{aligned}$$

- The blue term is called *implicit score matching*.

³A. Hyvärinen. Estimation of non-normalized statistical models by score matching. *J. Mach. Learn. Res.* **6**(4), 695-709 (2005).

DDPM in continuous setting⁴

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- Forward Ornstein-Uhlenbeck process with time-varying noise $\beta(t) > 0$:

$$d\mathbf{X}_t = -\frac{1}{2}\beta(t)\mathbf{X}_t dt + \sqrt{\beta(t)}d\mathbf{W}_t, \quad \mathbf{X}_0 \sim \pi_0.$$

- The solution reads:

$$\mathbf{X}_t = e^{-\frac{1}{2}\int_0^t \beta(s)ds} \left(\mathbf{X}_0 + \int_0^t e^{\frac{1}{2}\int_0^s \beta(s')ds'} \sqrt{\beta(s)} d\mathbf{W}_s \right),$$

with moments:

$$\mathbb{E}\{\mathbf{X}_t|\mathbf{X}_0\} = e^{-\frac{1}{2}\int_0^t \beta(s)ds} \mathbf{X}_0,$$

$$\mathbb{V}\{\mathbf{X}_t|\mathbf{X}_0\} = \left(1 - e^{-\int_0^t \beta(s)ds}\right) \mathbf{I}.$$

Apply again Itô's isometry formula with $Y_s = Z_s = e^{\frac{1}{2}\int_0^s \beta(s')ds'} \sqrt{\beta(s)}$.

⁴Called Variance Preserving (VP) SDE in: Y. Song, J. Sohl-Dickstein, D. P. Kingma, A. Kumar, S. Ermon, B. Poole. Score-based generative modeling through stochastic differential equations. [arXiv:2011.13456](https://arxiv.org/abs/2011.13456) (2020).

DDPM in continuous setting

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- Backward Ornstein-Uhlenbeck process:

$$d\mathbf{Y}_t = -\frac{1}{2}\beta(t) (\mathbf{Y}_t + 2\mathbf{s}(\mathbf{Y}_t, t)) dt + \sqrt{\beta(t)}d\mathbf{W}_t,$$

starting from $\mathbf{Y}_0 = \mathbf{X}_T \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$.

DDPM in discrete setting

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case

DDPM
SMLD

- Forward process: apply *e.g.* Euler-Maruyama scheme,

$$\begin{aligned}\mathbf{X}_{i+1} &= \left(1 - \frac{1}{2}\beta(t_i)\Delta t\right) \mathbf{X}_i + \sqrt{\beta(t_i)\Delta t} \mathbf{Z}, \quad \mathbf{Z} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}), \\ &\simeq \sqrt{1 - \beta_i} \mathbf{X}_i + \sqrt{\beta_i} \mathbf{Z} \\ &= \sqrt{\alpha_i} \mathbf{X}_i + \sqrt{1 - \alpha_i} \mathbf{Z}\end{aligned}$$

setting $\beta(t_i)\Delta t = \beta_i = 1 - \alpha_i$ such that $\beta_i \ll 1$.

- Backward process:

$$\begin{aligned}\mathbf{X}_{i-1} &= \left(1 + \frac{1}{2}\beta_i\right) \mathbf{X}_i + \beta_i \mathbf{s}(\mathbf{X}_i, t_i) + \sqrt{\beta_i} \mathbf{Z} \\ &\simeq \left(1 + \frac{1}{2}\beta_i\right) (\mathbf{X}_i + \beta_i \mathbf{s}(\mathbf{X}_i, t_i)) + \sqrt{\beta_i} \mathbf{Z} \\ &\simeq \frac{1}{\sqrt{1 - \beta_i}} (\mathbf{X}_i + \beta_i \mathbf{s}(\mathbf{X}_i, t_i)) + \sqrt{\beta_i} \mathbf{Z} \\ &= \frac{1}{\sqrt{\alpha_i}} (\mathbf{X}_i + (1 - \alpha_i) \mathbf{s}(\mathbf{X}_i, t_i)) + \sqrt{1 - \alpha_i} \mathbf{Z}.\end{aligned}$$

DDPM encoder

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- Given a *noise schedule* $\phi = (\alpha_1, \alpha_2, \dots, \alpha_T)$, the *encoder* is the Markov chain $\mathbf{X}_{1:T}$:

$$\mathbf{X}_i = \sqrt{\alpha_i} \mathbf{X}_{i-1} + \sqrt{1 - \alpha_i} \mathbf{Z}, \quad \mathbf{Z} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}),$$

starting from $\mathbf{X}_0 \sim \pi_0$, with transition probabilities:

$$\mathbb{Q}_\phi(\mathbf{x}_i | \mathbf{x}_{i-1}) = \mathcal{N}(\mathbf{x}_i; \sqrt{\alpha_i} \mathbf{x}_{i-1}, (1 - \alpha_i) \mathbf{I}),$$

$$\mathbb{Q}_\phi(\mathbf{x}_i | \mathbf{x}_0) = \mathcal{N}(\mathbf{x}_i; \sqrt{A_i} \mathbf{x}_0, (1 - A_i) \mathbf{I}),$$

where $A_i = \prod_{j=1}^i \alpha_j$.

- Choose $\alpha_1 > \dots > \alpha_T$ so that $\mathbb{Q}_\phi(\mathbf{x}_T | \mathbf{x}_0) \approx \mathcal{N}(\mathbf{x}_T; \mathbf{0}, \mathbf{I})$ independently of \mathbf{x}_0 .

DDPM encoder

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- $\mathbf{X}_{1:T}$ also has reverse transition probability:

$$\mathbb{Q}_{\phi}(\mathbf{x}_{i-1}|\mathbf{x}_i, \mathbf{x}_0) = \mathcal{N}(\mathbf{x}_{i-1}; \boldsymbol{\mu}_i(\mathbf{x}_i, \mathbf{x}_0), \tau_i^2 \mathbf{I}) ,$$

where:

$$\boldsymbol{\mu}_i(\mathbf{x}_i, \mathbf{x}_0) = \tau_i^2 \left(\frac{\sqrt{\alpha_i}}{1 - \alpha_i} \mathbf{x}_i + \frac{\sqrt{A_{i-1}}}{1 - A_{i-1}} \mathbf{x}_0 \right) , \quad \tau_i^2 = \frac{(1 - \alpha_i)(1 - A_{i-1})}{1 - A_i} .$$

Apply Bayes rule:

$$\begin{aligned} \mathbb{Q}_{\phi}(\mathbf{x}_i|\mathbf{x}_{i-1}) &= \mathbb{Q}_{\phi}(\mathbf{x}_i|\mathbf{x}_{i-1}, \mathbf{x}_0) = \frac{\mathbb{Q}_{\phi}(\mathbf{x}_{i-1}|\mathbf{x}_i, \mathbf{x}_0)\mathbb{Q}_{\phi}(\mathbf{x}_i|\mathbf{x}_0)}{\mathbb{Q}_{\phi}(\mathbf{x}_{i-1}|\mathbf{x}_0)} \\ \Rightarrow \mathbb{Q}_{\phi}(\mathbf{x}_{i-1}|\mathbf{x}_i, \mathbf{x}_0) &= \frac{\mathbb{Q}_{\phi}(\mathbf{x}_i|\mathbf{x}_{i-1})\mathbb{Q}_{\phi}(\mathbf{x}_{i-1}|\mathbf{x}_0)}{\mathbb{Q}_{\phi}(\mathbf{x}_i|\mathbf{x}_0)} \\ &= \frac{\mathcal{N}(\mathbf{x}_i; \sqrt{\alpha_i}\mathbf{x}_{i-1}, (1 - \alpha_i)\mathbf{I})\mathcal{N}(\mathbf{x}_{i-1}; \sqrt{A_{i-1}}\mathbf{x}_0, (1 - A_{i-1})\mathbf{I})}{\mathcal{N}(\mathbf{x}_i; \sqrt{A_i}\mathbf{x}_0, (1 - A_i)\mathbf{I})} . \end{aligned}$$

ELBO

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case

DDPM
SMLD

- *Unsupervised learning*: infer $\mathbf{X}_0^* \sim \pi_0 = d\mathbb{P}(\mathbf{x}_0)$ using $\mathbf{X}_{1:T}$ as latent variables,

$$\text{ELBO}(\mathbf{x}_0; \boldsymbol{\theta}, \phi) \stackrel{\text{def}}{=} \mathbb{E}_{\mathbb{Q}_{\phi}(\mathbf{x}_{1:T}|\mathbf{x}_0)} \left\{ \log \frac{\mathbb{P}_{\boldsymbol{\theta}}(\mathbf{x}_0, \mathbf{X}_{1:T})}{\mathbb{Q}_{\phi}(\mathbf{X}_{1:T}|\mathbf{x}_0)} \right\}.$$

- Sampling $\mathbf{X}_{1:T} \sim \mathbb{Q}_{\phi}(\mathbf{x}_{1:T}|\mathbf{x}_0)$ and maximizing $\text{ELBO}(\mathbf{x}_0; \boldsymbol{\theta}, \phi)$, we almost do maximize $\log \mathbb{P}(\mathbf{x}_0)$ on average.

$$\begin{aligned} \log \mathbb{P}(\mathbf{x}_0) &= \log \int \mathbb{P}(\mathbf{x}_0, d\mathbf{x}_{1:T}) \\ &= \log \mathbb{E}_{\mathbb{Q}_{\phi}(\mathbf{x}_{1:T}|\mathbf{x}_0)} \left\{ \frac{\mathbb{P}(\mathbf{x}_0, \mathbf{X}_{1:T})}{\mathbb{Q}_{\phi}(\mathbf{X}_{1:T}|\mathbf{x}_0)} \right\} \\ &\geq \mathbb{E}_{\mathbb{Q}_{\phi}(\mathbf{x}_{1:T}|\mathbf{x}_0)} \left\{ \log \frac{\mathbb{P}(\mathbf{x}_0, \mathbf{X}_{1:T})}{\mathbb{Q}_{\phi}(\mathbf{X}_{1:T}|\mathbf{x}_0)} \right\}, \end{aligned}$$

using Jensen's inequality $f(\mathbb{E}\{X\}) \geq \mathbb{E}\{f(X)\}$ with $f(x) = \log x$.

ELBO

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- The ELBO also reads:

$$\text{ELBO}(\mathbf{x}_0; \boldsymbol{\theta}, \phi) = \mathbb{E}_{\mathbb{Q}_{\phi}(\mathbf{x}_1|\mathbf{x}_0)} \{ \log \mathbb{P}_{\boldsymbol{\theta}}(\mathbf{x}_0|\mathbf{X}_1) \} - \mathbb{D}_{\text{KL}}(\mathbb{Q}_{\phi}(\mathbf{x}_T|\mathbf{x}_0) || \mathbb{P}(\mathbf{x}_T)) \\ - \sum_{i=2}^T \mathbb{E}_{\mathbb{Q}_{\phi}(\mathbf{x}_i|\mathbf{x}_0)} \{ \mathbb{D}_{\text{KL}}(\mathbb{Q}_{\phi}(\mathbf{x}_{i-1}|\mathbf{X}_i, \mathbf{x}_0) || \mathbb{P}_{\boldsymbol{\theta}}(\mathbf{x}_{i-1}|\mathbf{X}_i)) \}.$$

- *Prior matching*: $\mathbb{D}_{\text{KL}}(\mathbb{Q}_{\phi}(\mathbf{x}_T|\mathbf{x}_0) || \mathbb{P}(\mathbf{x}_T))$ tells how good the encoder is *vs.* a target distribution of the latent \mathbf{X}_T , and has to be minimized to maximize the ELBO;
- *Reconstruction*: $\mathbb{E}_{\mathbb{Q}_{\phi}(\mathbf{x}_1|\mathbf{x}_0)} \{ \log \mathbb{P}_{\boldsymbol{\theta}}(\mathbf{x}_0|\mathbf{X}_1) \}$ tells how good the *decoder* (denoiser) is, and has to be maximized to maximize the ELBO;
- *Diffusion loss*: $\mathbb{D}_{\text{KL}}(\mathbb{Q}_{\phi}(\mathbf{x}_{i-1}|\mathbf{X}_i, \mathbf{x}_0) || \mathbb{P}_{\boldsymbol{\theta}}(\mathbf{x}_{i-1}|\mathbf{X}_i))$, $2 \leq i \leq T$, tell how much forward/backward transitions to latent variables are consistent, and have to be minimized to maximize the ELBO.

Training

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- **Encoder:** choose the noise schedule ϕ , in which case there is nothing left to learn in $\mathbb{D}_{\text{KL}}(\mathbb{Q}_{\phi}(\mathbf{x}_T|\mathbf{x}_0)||\mathbb{P}(\mathbf{x}_T))$;
- **Decoder:** choose $\mathbb{P}_{\theta}(\mathbf{x}_{i-1}|\mathbf{x}_i) = \mathcal{N}(\mathbf{x}_{i-1}; \boldsymbol{\mu}_{\theta}(\mathbf{x}_i), \tau_i^2 \mathbf{I})$, where $\boldsymbol{\mu}_{\theta}(\mathbf{x}_i)$ is a Neural Network;
- **Loss function:** then the consistency \mathbb{D}_{KL} 's and reconstruction term read

$$\mathbb{D}_{\text{KL}}(\mathbb{Q}_{\phi}(\mathbf{x}_{i-1}|\mathbf{X}_i, \mathbf{x}_0)||\mathbb{P}_{\theta}(\mathbf{x}_{i-1}|\mathbf{X}_i)) = \frac{1}{2\tau_i^2} \|\boldsymbol{\mu}_{\theta}(\mathbf{X}_i) - \boldsymbol{\mu}_i(\mathbf{X}_i, \mathbf{x}_0)\|^2,$$

$$\log \mathbb{P}_{\theta}(\mathbf{x}_0|\mathbf{X}_1) = -\frac{1}{2\tau_1^2} \|\boldsymbol{\mu}_{\theta}(\mathbf{X}_1) - \mathbf{x}_0\|^2 + C,$$

such that (since $\boldsymbol{\mu}_1(\mathbf{X}_1, \mathbf{x}_0) = \mathbf{x}_0$ with $\alpha_0 = 1$)

$$\begin{aligned} \text{ELBO}(\mathbf{x}_0; \theta, \phi) &= -\frac{1}{2} \sum_{i=1}^T \mathbb{E}_{\mathbb{Q}_{\phi}(\mathbf{x}_i|\mathbf{x}_0)} \left\{ \frac{1}{\tau_i^2} \|\boldsymbol{\mu}_{\theta}(\mathbf{X}_i) - \boldsymbol{\mu}_i(\mathbf{X}_i, \mathbf{x}_0)\|^2 \right\} \\ &\simeq -\frac{T}{2} \mathbb{E}_{\mathbb{Q}_{\phi}(\mathbf{x}_i|\mathbf{x}_0), i \sim \mathcal{U}(1, T)} \left\{ \frac{1}{\tau_i^2} \|\boldsymbol{\mu}_{\theta}(\mathbf{X}_i) - \boldsymbol{\mu}_i(\mathbf{X}_i, \mathbf{x}_0)\|^2 \right\}. \end{aligned}$$

Inference

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- **Inference:** with optimal parameters θ^* of the Neural Network μ_θ ,

$$\theta^* = \arg \max_{\theta} \mathbb{E}_{\mathbf{X}_0} \{ \text{ELBO}(\mathbf{X}_0; \theta, \phi) \}$$

$$= \arg \min_{\theta} \mathbb{E}_{\mathbb{Q}_{\phi}(\mathbf{x}_0, \mathbf{x}_i), i \sim \mathcal{U}(1, T)} \left\{ \frac{1}{2\tau_i^2} \|\mu_{\theta}(\mathbf{X}_i) - \mu_i(\mathbf{X}_i, \mathbf{X}_0)\|^2 \right\},$$

for $i = T, T-1, \dots, 1$ do

$$\mathbf{X}_{i-1} = \mu_{\theta^*}(\mathbf{X}_i) + \tau_i \mathbf{Z}, \quad \mathbf{Z} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}),$$

starting with $\mathbf{X}_T \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$.

Alternative training #1

"Predict image"

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- Since:

$$\mu_i(\mathbf{x}_i, \mathbf{x}_0) = \tau_i^2 \left(\frac{\sqrt{\alpha_i}}{1 - \alpha_i} \mathbf{x}_i + \frac{\sqrt{A_{i-1}}}{1 - A_{i-1}} \mathbf{x}_0 \right),$$

choose the reparameterization:

$$\mu_{\theta}(\mathbf{x}_i) = \tau_i^2 \left(\frac{\sqrt{\alpha_i}}{1 - \alpha_i} \mathbf{x}_i + \frac{\sqrt{A_{i-1}}}{1 - A_{i-1}} \hat{\mathbf{x}}_{\theta}(\mathbf{x}_i) \right);$$

- Then:

$$\|\mu_{\theta}(\mathbf{X}_i) - \mu_i(\mathbf{X}_i, \mathbf{x}_0)\|^2 = \frac{(1 - \alpha_i)^2 A_{i-1}}{(1 - A_i)^2} \|\hat{\mathbf{x}}_{\theta}(\mathbf{X}_i) - \mathbf{x}_0\|^2,$$

and with $\Delta_i = \text{SNR}_{i-1} - \text{SNR}_i > 0$, $\text{SNR}_i = \frac{A_i}{1 - A_i}$:

$$\text{ELBO}(\mathbf{x}_0; \theta, \phi) = -\frac{1}{2} \sum_{i=1}^T \mathbb{E}_{\mathbb{Q}_{\phi}(\mathbf{x}_i | \mathbf{x}_0)} \{ \Delta_i \|\hat{\mathbf{x}}_{\theta}(\mathbf{X}_i) - \mathbf{x}_0\|^2 \}.$$

Alternative training⁵ #1

Learning the noise schedule

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- The *signal-to-noise ratio* SNR_i of $\mathbf{X}_i \sim \mathbb{Q}_\phi(\mathbf{x}_i|\mathbf{x}_0)$ is represented by a monotonically increasing Neural Network $\omega_\phi(t)$:

$$\text{SNR}_i \stackrel{\text{def}}{=} \frac{A_i}{1 - A_i} = e^{-\omega_\phi(t_i)},$$

such that $A_i = \sigma(-\omega_\phi(t_i))$ and $1 - A_i = \sigma(\omega_\phi(t_i))$,
where:

$$\sigma(x) = \frac{1}{1 + e^{-x}}$$

is the usual sigmoid, or logistic activation function;

- Then $\mathbf{x} \rightarrow \hat{\mathbf{x}}_\theta(\mathbf{x})$ and $t \rightarrow \omega_\phi(t)$ are learnt altogether.

⁵D. P. Kingma, T. Salimans, B. Poole, J. Ho. Variational diffusion models.
arXiv:2107.00630 (2021).

Alternative training⁶ #2

"Predict noise"

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- Since $\mathbf{X}_i = \sqrt{A_i}\mathbf{X}_0 + \sqrt{1 - A_i}\mathbf{Z}$:

$$\boldsymbol{\mu}_i(\mathbf{x}_i, \mathbf{x}_0) \rightarrow \boldsymbol{\mu}_i(\mathbf{x}_i, \mathbf{z}) = \frac{1}{\sqrt{\alpha_i}} \left(\mathbf{x}_i - \frac{1 - \alpha_i}{\sqrt{1 - A_i}} \mathbf{z} \right),$$

choose the reparameterization:

$$\boldsymbol{\mu}_{\boldsymbol{\theta}}(\mathbf{x}_i) = \frac{1}{\sqrt{\alpha_i}} \left(\mathbf{x}_i - \frac{1 - \alpha_i}{\sqrt{1 - A_i}} \hat{\mathbf{z}}_{\boldsymbol{\theta}}(\mathbf{x}_i) \right);$$

- Then:

$$\|\boldsymbol{\mu}_{\boldsymbol{\theta}}(\mathbf{X}_i) - \boldsymbol{\mu}_i(\mathbf{X}_i, \mathbf{z})\|^2 = \frac{(1 - \alpha_i)^2}{\alpha_i(1 - A_i)} \|\hat{\mathbf{z}}_{\boldsymbol{\theta}}(\mathbf{X}_i) - \mathbf{z}\|^2,$$

and with $\lambda_i = \frac{\Delta_i}{\text{SNR}_i} = e^{\omega_{\phi}(t_i) - \omega_{\phi}(t_{i-1})} - 1$:

$$\text{ELBO}(\mathbf{x}_0; \boldsymbol{\theta}, \phi) = -\frac{1}{2} \sum_{i=1}^T \mathbb{E}_{\mathbf{Z} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})} \left\{ \lambda_i \left\| \hat{\mathbf{z}}_{\boldsymbol{\theta}} \left(\sqrt{A_i} \mathbf{x}_0 + \sqrt{1 - A_i} \mathbf{Z} \right) - \mathbf{Z} \right\|^2 \right\}.$$

⁶J. Ho, A. Jain, P. Abbeel. Denoising diffusion probabilistic models. [arXiv:2006.11239](https://arxiv.org/abs/2006.11239) (2020).

Alternative training #3

"Predict score"

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- Since $\mathbf{X}_i | \mathbf{X}_0 \sim \mathcal{N}(\sqrt{A_i} \mathbf{X}_0, (1 - A_i) \mathbf{I})$, *Tweedie's formula* yields $\mathbb{E}\{\sqrt{A_i} \mathbf{X}_0 | \mathbf{X}_i\} = \mathbf{X}_i + (1 - A_i) \mathbf{s}(\mathbf{X}_i, t_i)$ such that the reverse transition mean can be written:

$$\boldsymbol{\mu}_i(\mathbf{x}_i, \mathbf{x}_0) \rightarrow \boldsymbol{\mu}_i(\mathbf{x}_i, \mathbf{s}) = \frac{1}{\sqrt{\alpha_i}} (\mathbf{x}_i + (1 - \alpha_i) \mathbf{s}(\mathbf{x}_i, t_i)) ,$$

and one can choose the reparameterization:

$$\boldsymbol{\mu}_{\boldsymbol{\theta}}(\mathbf{x}_i) = \frac{1}{\sqrt{\alpha_i}} (\mathbf{x}_i + (1 - \alpha_i) \hat{\mathbf{s}}_{\boldsymbol{\theta}}(\mathbf{x}_i)) ;$$

- Then:

$$\|\boldsymbol{\mu}_{\boldsymbol{\theta}}(\mathbf{X}_i) - \boldsymbol{\mu}_i(\mathbf{X}_i, \mathbf{s})\|^2 = \frac{(1 - \alpha_i)^2}{\alpha_i} \|\hat{\mathbf{s}}_{\boldsymbol{\theta}}(\mathbf{X}_i) - \mathbf{s}(\mathbf{X}_i, t_i)\|^2 ,$$

and with $\lambda'_i = (1 - \alpha_i) \frac{\text{SNR}_{i-1}}{\text{SNR}_i}$:

$$\text{ELBO}(\mathbf{x}_0; \boldsymbol{\theta}, \phi) = -\frac{1}{2} \sum_{i=1}^T \mathbb{E}_{\mathbb{Q}_{\phi}(\mathbf{x}_i | \mathbf{x}_0)} \left\{ \lambda'_i \|\hat{\mathbf{s}}_{\boldsymbol{\theta}}(\mathbf{X}_i) - \mathbf{s}(\mathbf{X}_i, t_i)\|^2 \right\} .$$

Tweedie's formula

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- Let $\mathbf{Y} = \mathbf{X} + \mathcal{N}(\mathbf{0}, \sigma^2 \mathbf{I})$ where $\mathbf{X} \sim \pi_{\mathbf{X}}$;
- Then $\pi_{\mathbf{Y}}(\mathbf{y}) \propto \int \pi_{\mathbf{X}}(\mathbf{x}) \exp(-\frac{\|\mathbf{y}-\mathbf{x}\|^2}{2\sigma^2}) d\mathbf{x}$ and:

$$\begin{aligned} \frac{\mathbb{E}\{\mathbf{X}|\mathbf{Y}=\mathbf{y}\}-\mathbf{y}}{\sigma^2} &= \frac{\int \mathbf{x} \frac{\pi_{\mathbf{X},\mathbf{Y}}(\mathbf{x},\mathbf{y}) d\mathbf{x}}{\pi_{\mathbf{Y}}(\mathbf{y})} - \mathbf{y}}{\sigma^2} \\ &= \frac{\int (\frac{\mathbf{x}-\mathbf{y}}{\sigma^2}) \pi_{\mathbf{X}}(\mathbf{x}) \exp(-\frac{\|\mathbf{y}-\mathbf{x}\|^2}{2\sigma^2}) d\mathbf{x}}{\int \pi_{\mathbf{X}}(\mathbf{x}) \exp(-\frac{\|\mathbf{y}-\mathbf{x}\|^2}{2\sigma^2}) d\mathbf{x}} \\ &= \nabla_{\mathbf{y}} \log \int \pi_{\mathbf{X}}(\mathbf{x}) \exp\left(-\frac{\|\mathbf{y}-\mathbf{x}\|^2}{2\sigma^2}\right) d\mathbf{x} \\ &= \nabla_{\mathbf{y}} \log \pi_{\mathbf{Y}}(\mathbf{y}) \\ &= \mathbf{s}_{\mathbf{Y}}(\mathbf{y}); \end{aligned}$$

- Thus $\mathbb{E}\{\mathbf{X}|\mathbf{Y}\} = \mathbf{Y} + \sigma^2 \mathbf{s}_{\mathbf{Y}}(\mathbf{Y})$, which allows to estimate the score $\mathbf{s}_{\mathbf{Y}}$ given the data $\{\mathbf{x}_i, \mathbf{y}_i\}_{i=1}^N$.

SMLD in continuous setting⁷

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- Forward process with time-varying noise $\beta(t) > 0$:

$$d\mathbf{X}_t = \sqrt{\beta(t)}d\mathbf{W}_t, \quad \mathbf{X}_0 \sim \pi_0.$$

- The solution reads:

$$\mathbf{X}_t = \mathbf{X}_0 + \int_0^t \sqrt{\beta(s)}d\mathbf{W}_s,$$

with moments:

$$\mathbb{E}\{\mathbf{X}_t|\mathbf{X}_0\} = \mathbf{X}_0,$$

$$\mathbb{V}\{\mathbf{X}_t|\mathbf{X}_0\} = \left(\int_0^t \beta(s)ds\right) \mathbf{I}.$$

⁷Called Variance Exploding (VE) SDE in: Y. Song, J. Sohl-Dickstein, D. P. Kingma, A. Kumar, S. Ermon, B. Poole. Score-based generative modeling through stochastic differential equations. [arXiv:2011.13456](https://arxiv.org/abs/2011.13456) (2020).

SMLD in continuous setting

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- Backward process:

$$d\mathbf{Y}_t = -\beta(t)\mathbf{s}(\mathbf{Y}_t, t)dt + \sqrt{\beta(t)}d\mathbf{W}_t, \quad \mathbf{Y}_0 = \mathbf{X}_T,$$

which looks like (stochastic) gradient descent on $\log \pi$ with learning rate $\beta(t)$ since Stein's score function reads $\mathbf{s}(\mathbf{x}, t) = \nabla_{\mathbf{x}} \log \pi(\mathbf{x}, t)$.

SMLD encoder

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- Given a *variance schedule* $\phi = (\sigma_1^2, \sigma_2^2, \dots, \sigma_T^2)$, the *encoder* is the Markov chain $\mathbf{X}_{1:T}$:

$$\mathbf{X}_i = \mathbf{X}_{i-1} + \sqrt{\sigma_i^2 - \sigma_{i-1}^2} \mathbf{Z}, \quad \mathbf{Z} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}),$$

starting from $\mathbf{X}_0 \sim \pi_0$, with transition probabilities:

$$\mathbb{Q}_\phi(\mathbf{x}_i | \mathbf{x}_{i-1}) = \mathcal{N}(\mathbf{x}_i; \mathbf{x}_{i-1}, (\sigma_i^2 - \sigma_{i-1}^2) \mathbf{I}),$$

$$\mathbb{Q}_\phi(\mathbf{x}_i | \mathbf{x}_0) = \mathcal{N}(\mathbf{x}_i; \mathbf{x}_0, \sigma_i^2 \mathbf{I});$$

- $\mathbf{X}_{1:T}$ also has reverse transition probability:

$$\mathbb{Q}_\phi(\mathbf{x}_{i-1} | \mathbf{x}_i, \mathbf{x}_0) = \mathcal{N}(\mathbf{x}_{i-1}; \boldsymbol{\mu}_i(\mathbf{x}_i, \mathbf{x}_0), \tau_i^2 \mathbf{I}),$$

where:

$$\boldsymbol{\mu}_i(\mathbf{x}_i, \mathbf{x}_0) = \frac{1}{\sigma_i^2} (\sigma_{i-1}^2 \mathbf{x}_i + (\sigma_i^2 - \sigma_{i-1}^2) \mathbf{x}_0), \quad \tau_i^2 = \frac{\sigma_{i-1}^2}{\sigma_i^2} (\sigma_i^2 - \sigma_{i-1}^2).$$

Training

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- **Encoder:** choose the variance schedule ϕ , so there is nothing left to learn in $\mathbb{D}_{\text{KL}}(\mathbb{Q}_{\phi}(\mathbf{x}_T|\mathbf{x}_0)||\mathbb{P}(\mathbf{x}_T))$;
- **Decoder:** choose $\mathbb{P}_{\theta}(\mathbf{x}_{i-1}|\mathbf{x}_i) = \mathcal{N}(\mathbf{x}_{i-1}; \boldsymbol{\mu}_{\theta}(\mathbf{x}_i), \tau_i^2 \mathbf{I})$, where $\boldsymbol{\mu}_{\theta}(\mathbf{x}_i)$ is a Neural Network;
- **Loss function:** then the consistency \mathbb{D}_{KL} 's and reconstruction term read

$$\mathbb{D}_{\text{KL}}(\mathbb{Q}_{\phi}(\mathbf{x}_{i-1}|\mathbf{X}_i, \mathbf{x}_0)||\mathbb{P}_{\theta}(\mathbf{x}_{i-1}|\mathbf{X}_i)) = \frac{1}{2\tau_i^2} \|\boldsymbol{\mu}_{\theta}(\mathbf{X}_i) - \boldsymbol{\mu}_i(\mathbf{X}_i, \mathbf{x}_0)\|^2 ,$$

$$\log \mathbb{P}_{\theta}(\mathbf{x}_0|\mathbf{X}_1) = -\frac{1}{2\tau_1^2} \|\boldsymbol{\mu}_{\theta}(\mathbf{X}_1) - \mathbf{x}_0\|^2 + C ,$$

such that (since $\boldsymbol{\mu}_1(\mathbf{X}_1, \mathbf{x}_0) = \mathbf{x}_0$ with $\sigma_0 = 0$)

$$\begin{aligned} \text{ELBO}(\mathbf{x}_0; \theta, \phi) &= -\frac{1}{2} \sum_{i=1}^T \mathbb{E}_{\mathbb{Q}_{\phi}(\mathbf{x}_i|\mathbf{x}_0)} \left\{ \frac{1}{\tau_i^2} \|\boldsymbol{\mu}_{\theta}(\mathbf{X}_i) - \boldsymbol{\mu}_i(\mathbf{X}_i, \mathbf{x}_0)\|^2 \right\} \\ &\simeq -\frac{T}{2} \mathbb{E}_{\mathbb{Q}_{\phi}(\mathbf{x}_i|\mathbf{x}_0), i \sim \mathcal{U}(1, T)} \left\{ \frac{1}{\tau_i^2} \|\boldsymbol{\mu}_{\theta}(\mathbf{X}_i) - \boldsymbol{\mu}_i(\mathbf{X}_i, \mathbf{x}_0)\|^2 \right\} . \end{aligned}$$

Alternative training #1

”Predict image”

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- Since:

$$\mu_i(\mathbf{x}_i, \mathbf{x}_0) = \frac{1}{\sigma_i^2} \left(\sigma_{i-1}^2 \mathbf{x}_i + (\sigma_i^2 - \sigma_{i-1}^2) \mathbf{x}_0 \right) ,$$

choose the reparameterization:

$$\mu_{\theta}(\mathbf{x}_i) = \frac{1}{\sigma_i^2} \left(\sigma_{i-1}^2 \mathbf{x}_i + (\sigma_i^2 - \sigma_{i-1}^2) \hat{\mathbf{x}}_{\theta}(\mathbf{x}_i) \right) ;$$

- Then:

$$\|\mu_{\theta}(\mathbf{X}_i) - \mu_i(\mathbf{X}_i, \mathbf{x}_0)\| = \frac{\sigma_i^2 - \sigma_{i-1}^2}{\sigma_i^2} \|\hat{\mathbf{x}}_{\theta}(\mathbf{X}_i) - \mathbf{x}_0\| ,$$

and with $\Delta_i = 1/\sigma_{i-1}^2 - 1/\sigma_i^2$:

$$\text{ELBO}(\mathbf{x}_0; \theta, \phi) = -\frac{1}{2} \sum_{i=1}^T \mathbb{E}_{\mathbb{Q}_{\phi}(\mathbf{x}_i|\mathbf{x}_0)} \{ \Delta_i \|\hat{\mathbf{x}}_{\theta}(\mathbf{X}_i) - \mathbf{x}_0\|^2 \} .$$

Alternative training⁸ #2

"Predict noise"

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- Since $\mathbf{X}_i = \mathbf{X}_0 + \sigma_i \mathbf{Z}$:

$$\mu_i(\mathbf{x}_i, \mathbf{x}_0) \rightarrow \mu_i(\mathbf{x}_i, \mathbf{z}) = \mathbf{x}_i - \frac{\sigma_i^2 - \sigma_{i-1}^2}{\sigma_i} \mathbf{z},$$

choose the reparameterization:

$$\mu_{\theta}(\mathbf{x}_i) = \mathbf{x}_i - \frac{\sigma_i^2 - \sigma_{i-1}^2}{\sigma_i} \hat{\mathbf{z}}_{\theta}(\mathbf{x}_i);$$

- Then:

$$\|\mu_{\theta}(\mathbf{X}_i) - \mu_i(\mathbf{X}_i, \mathbf{z})\| = \frac{\sigma_i^2 - \sigma_{i-1}^2}{\sigma_i} \|\hat{\mathbf{z}}_{\theta}(\mathbf{X}_i) - \mathbf{z}\|,$$

and with $\lambda_i = \sigma_i^2 / \sigma_{i-1}^2 - 1$:

$$\text{ELBO}(\mathbf{x}_0; \theta, \phi) = -\frac{1}{2} \sum_{i=1}^T \mathbb{E}_{\mathbf{Z} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})} \left\{ \lambda_i \|\hat{\mathbf{z}}_{\theta}(\mathbf{x}_0 + \sigma_i \mathbf{Z}) - \mathbf{Z}\|^2 \right\}.$$

⁸Y. Song, S. Ermon. Generative modeling by estimating gradients of the data distribution. [arXiv:1907.05600](https://arxiv.org/abs/1907.05600) (2019).

Inference

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- **Inference:** with optimal parameters θ^* of the Neural Network \hat{z}_θ ,

$$\begin{aligned}\theta^* &= \arg \max_{\theta} \mathbb{E}_{\mathbf{X}_0} \{ \text{ELBO}(\mathbf{X}_0; \theta, \phi) \} \\ &= \arg \min_{\theta} \mathbb{E}_{\mathbf{X}_0, \mathbf{Z}, i \sim \mathcal{U}(1, T)} \left\{ \lambda_i \left\| \hat{z}_\theta(\mathbf{X}_0 + \sigma_i \mathbf{Z}) - \mathbf{Z} \right\|^2 \right\},\end{aligned}$$

for $i = T, T-1, \dots, 1$ do

$$\mathbf{X}_{i-1} = \mathbf{X}_i - \frac{\sigma_i^2 - \sigma_{i-1}^2}{\sigma_i} \hat{z}_{\theta^*}(\mathbf{X}_i) + \tau_i \mathbf{Z}, \quad \mathbf{Z} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}),$$

starting with $\mathbf{X}_T \sim \mathcal{N}(\mathbf{0}, \sigma_T^2 \mathbf{I})$.

To be continued...

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical
solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion &
ML

Continuous
case
DDPM
SMLD

- Conditioning (called *guidance* in diffusion networks literature);
- Boundary conditions (Maarten's question);
- (Schrödinger) bridges and manifolds...

Further reading...

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion & ML

Continuous
case
DDPM
SMLD

Stochastic differential equations:

- A. Friedman: *Stochastic Differential Equations and Applications*, vol. I & II, Academic Press (1975);
- P.E. Kloeden, E. Platen: *Numerical Solution of Stochastic Differential Equations*, 3rd ed., Springer (1999);
- J.-F. Le Gall: *Mouvement Brownien, Martingales et Calcul Stochastique*, Springer (2013);
- B.K. Øksendal: *Stochastic Differential Equations: An Introduction with Applications*, 6th ed., Springer (2003);
- S. Särkkä, A. Solin: *Applied Stochastic Differential Equations*, Cambridge University Press (2019);
- C. Soize: *The Fokker-Planck Equation for Stochastic Dynamical Systems and its Explicit Steady State Solutions*, World Scientific (1994);
- D. Talay: Simulation of stochastic differential systems. In *Probabilistic Methods in Applied Physics* (P. Krée & W. Wedig, eds.), pp. 54-96. Lecture Notes in Physics **451**, Springer (1995).

Further reading...

SDE

W. Genuist,
É. Savin

SDE

First-order
systems
Stochastic
integrals
Diffusion
processes

Numerical solutions

Stochastic
modeling
Numerical
schemes

GAN

VAE

Diffusion & ML

Continuous
case
DDPM
SMLD

Variational diffusion models:

- S. H. Chan. Tutorial on diffusion models for imaging and vision. [arXiv:2403.18103](#) (2024);
- V. de Bortoli. Generative modeling. https://vdebortoli.github.io/project/generative_modeling/ (2023);
- F. de Souza Ribeiro, B. Glocker. Demystifying variational diffusion models. [arXiv:2401.06281](#) (2024);
- T. Duan. Diffusion models from scratch. <http://www.tonyduan.com/diffusion/> (2023);
- C. Luo. Understanding diffusion models: A unified perspective. [arXiv:2208.11970](#) (2022);
- G. Peyré. Denoising diffusion models. Mathematical Tours (2023); [jupyter notebook](#);
- Y. Song, J. Sohl-Dickstein, D. P. Kingma, A. Kumar, S. Ermon, B. Poole. Score-based generative modeling through stochastic differential equations. [arXiv:2011.13456](#) (2020);
- R. E. Turner, C.-D. Diaconu, S. Markou, A. Shysheya, A. Y. K. Foong, B. Mlodozieniec. Denoising diffusion probabilistic models in six simple steps. [arXiv:2402.04384](#) (2024);
- L. Yang, Z. Zhang, Y. Song, S. Hong, R. Xu, Y. Zhao, W. Zhang, B. Cui, M.-H. Yang. Diffusion models: A comprehensive survey of methods and applications. [arXiv:2209.00796](#) (2022).