

# Econometrics I - TA section

Eric Schulman

The University of Texas at Austin

September 19, 2020



# Homework 1

## 1.5 HIE

If  $P(A) = 1/2$  and  $P(B) = 2/3$  can A and B be disjoint? Explain.

# Homework 1

## 1.5 HIE

If  $P(A) = 1/2$  and  $P(B) = 2/3$  can A and B be disjoint? Explain.

- No,  $P(A) + P(B) > 1$
- Disjoint events sum to at most 1.

# Homework 1

## 1.12 HIE

1. Drawing a King with one card.

# Homework 1

## 1.12 HIE

1. Drawing a King with one card. 4/52

# Homework 1

## 1.12 HIE

1. Drawing a King with one card.  $4/52$
2. Drawing a King on the second card, conditional on a King on the first card.

# Homework 1

## 1.12 HIE

1. Drawing a King with one card.  $4/52$
2. Drawing a King on the second card, conditional on a King on the first card.  $3/51$
3. Drawing two Kings with two cards.

# Homework 1

## 1.12 HIE

1. Drawing a King with one card.  $4/52$
2. Drawing a King on the second card, conditional on a King on the first card.  $3/51$
3. Drawing two Kings with two cards.  $4/52 \times 3/51$



# Homework 1

## 1.12 HIE

1. Drawing a King with one card.  $4/52$
2. Drawing a King on the second card, conditional on a King on the first card.  $3/51$
3. Drawing two Kings with two cards.  $4/52 \times 3/51$
4. Drawing a King on the second card, conditional on the first card is not a King.

# Homework 1

## 1.12 HIE

1. Drawing a King with one card.  $4/52$
2. Drawing a King on the second card, conditional on a King on the first card.  $3/51$
3. Drawing two Kings with two cards.  $4/52 \times 3/51$
4. Drawing a King on the second card, conditional on the first card is not a King.  $4/51$
5. Drawing a King on the second card, when the first card is placed face down (so is unknown).

# Homework 1

## 1.12 HIE

1. Drawing a King with one card.  $4/52$
2. Drawing a King on the second card, conditional on a King on the first card.  $3/51$
3. Drawing two Kings with two cards.  $4/52 \times 3/51$
4. Drawing a King on the second card, conditional on the first card is not a King.  $4/51$
5. Drawing a King on the second card, when the first card is placed face down (so is unknown).  
 $3/51 \times 4/52 + 4/51 \times 48/52$ 
  - Does it make a difference?

## 1.12 HIE

1. Drawing a King with one card.  $4/52$
2. Drawing a King on the second card, conditional on a King on the first card.  $3/51$
3. Drawing two Kings with two cards.  $4/52 \times 3/51$
4. Drawing a King on the second card, conditional on the first card is not a King.  $4/51$
5. Drawing a King on the second card, when the first card is placed face down (so is unknown).  
 $3/51 \times 4/52 + 4/51 \times 48/52$ 
  - Does it make a difference?
  - As expected, drawing a card face down doesn't effect the probability of drawing a king...

# Homework 1

## 1.14 HIE

1. Getting three heads in a row from three coin flips.

# Homework 1

## 1.14 HIE

1. Getting three heads in a row from three coin flips.  
 $(1/2)^3$

# Homework 1

## 1.14 HIE

1. Getting three heads in a row from three coin flips.  
 $(1/2)^3$
2. Getting a heads given that the previous coin was a tails.

# Homework 1

## 1.14 HIE

1. Getting three heads in a row from three coin flips.  
 $(1/2)^3$
2. Getting a heads given that the previous coin was a tails.  
 $1/2$



## 1.14 HIE

1. Getting three heads in a row from three coin flips.  
 $(1/2)^3$
2. Getting a heads given that the previous coin was a tails.  
 $1/2$
3. From two coin flips getting two heads given that at least one coin is a heads.

## 1.14 HIE

1. Getting three heads in a row from three coin flips.  
 $(1/2)^3$
2. Getting a heads given that the previous coin was a tails.  
 $1/2$
3. From two coin flips getting two heads given that at least one coin is a heads.  
 $1/3$  i.e. 4 possible outcomes, 1 is “ruled out” b/c only tails

# Homework 1

## 1.14 HIE

1. Getting three heads in a row from three coin flips.  
 $(1/2)^3$
2. Getting a heads given that the previous coin was a tails.  
 $1/2$
3. From two coin flips getting two heads given that at least one coin is a heads.  
 $1/3$  i.e. 4 possible outcomes, 1 is “ruled out” b/c only tails
4. Rolling a six from a pair of dice.

# Homework 1

## 1.14 HIE

1. Getting three heads in a row from three coin flips.  
 $(1/2)^3$
2. Getting a heads given that the previous coin was a tails.  
 $1/2$
3. From two coin flips getting two heads given that at least one coin is a heads.  
 $1/3$  i.e. 4 possible outcomes, 1 is “ruled out” b/c only tails
4. Rolling a six from a pair of dice.  
 $11/36$  i.e. 36 outcomes, don't double count (6,6)

# Homework 1

## 1.14 HIE

1. Getting three heads in a row from three coin flips.  
 $(1/2)^3$
2. Getting a heads given that the previous coin was a tails.  
 $1/2$
3. From two coin flips getting two heads given that at least one coin is a heads.  
 $1/3$  i.e. 4 possible outcomes, 1 is “ruled out” b/c only tails
4. Rolling a six from a pair of dice.  
 $11/36$  i.e. 36 outcomes, don't double count (6,6)
5. Rolling “snakes eyes” from a pair of dice. (Getting a pair of ones.)

# Homework 1

## 1.14 HIE

1. Getting three heads in a row from three coin flips.  
 $(1/2)^3$
2. Getting a heads given that the previous coin was a tails.  
 $1/2$
3. From two coin flips getting two heads given that at least one coin is a heads.  
 $1/3$  i.e. 4 possible outcomes, 1 is “ruled out” b/c only tails
4. Rolling a six from a pair of dice.  
 $11/36$  i.e. 36 outcomes, don't double count (6,6)
5. Rolling “snakes eyes” from a pair of dice. (Getting a pair of ones.)  
 $1/36$

# Homework 1

## 1.14 HIE

1. Getting three heads in a row from three coin flips.  
 $(1/2)^3$
2. Getting a heads given that the previous coin was a tails.  
 $1/2$
3. From two coin flips getting two heads given that at least one coin is a heads.  
 $1/3$  i.e. 4 possible outcomes, 1 is “ruled out” b/c only tails
4. Rolling a six from a pair of dice.  
 $11/36$  i.e. 36 outcomes, don't double count (6,6)
5. Rolling “snakes eyes” from a pair of dice. (Getting a pair of ones.)  
 $1/36$

## 1.19 HIE

1. “When I selected door A the probability that it has the prize was  $1/3$ . No information was revealed. So the probability that Door A has the prize remains  $1/3$ .”



## 1.19 HIE

1. “When I selected door A the probability that it has the prize was  $1/3$ . No information was revealed. So the probability that Door A has the prize remains  $1/3$ .” False

## 1.19 HIE

1. “When I selected door A the probability that it has the prize was  $1/3$ . No information was revealed. So the probability that Door A has the prize remains  $1/3$ .” False
2. “The original probability was  $1/3$  on each door. Now that door B is eliminated, doors A and C each have each probability of  $1/2$ . It does not matter if I stay with A or switch to C.”

## 1.19 HIE

1. “When I selected door A the probability that it has the prize was  $1/3$ . No information was revealed. So the probability that Door A has the prize remains  $1/3$ .” False
2. “The original probability was  $1/3$  on each door. Now that door B is eliminated, doors A and C each have each probability of  $1/2$ . It does not matter if I stay with A or switch to C.” False

## 1.19 HIE

1. “When I selected door A the probability that it has the prize was  $1/3$ . No information was revealed. So the probability that Door A has the prize remains  $1/3$ .” False
2. “The original probability was  $1/3$  on each door. Now that door B is eliminated, doors A and C each have each probability of  $1/2$ . It does not matter if I stay with A or switch to C.” False

## 1.19 HIE

1. “The host inadvertently revealed information. If door C had the prize, he was forced to open door B. If door B had the prize he would have been forced to open door C. Thus it is quite likely that door C has the prize.”

## 1.19 HIE

1. “The host inadvertently revealed information. If door C had the prize, he was forced to open door B. If door B had the prize he would have been forced to open door C. Thus it is quite likely that door C has the prize.” True

## 1.19 HIE

1. “The host inadvertently revealed information. If door C had the prize, he was forced to open door B. If door B had the prize he would have been forced to open door C. Thus it is quite likely that door C has the prize.” True
2. Assume a prior probability for each door of  $1/3$ . Calculate the posterior probability that door A and door C have the prize. What choice do you recommend for the contestant?

## 1.19 HIE

1. “The host inadvertently revealed information. If door C had the prize, he was forced to open door B. If door B had the prize he would have been forced to open door C. Thus it is quite likely that door C has the prize.” True
2. Assume a prior probability for each door of  $1/3$ . Calculate the posterior probability that door A and door C have the prize. What choice do you recommend for the contestant?  
 $1/3$  chance A.  $2/3$  chance B or C. We know C does not have the prize. Thus  $2/3$  chance  $B$  has the prize.



# Homework 1

## 1.21 HIE

1. A straight

## 1.21 HIE

### 1. A straight

This is five cards in a sequence (e.g., 4,5,6,7,8), with aces allowed to be either 1 or 13 (low or high) and with the cards allowed to be of the same suit (e.g., all hearts) or from some different suits. The number of such hands is  $10 \times (4\text{choose}1)^5$

### 2. A flush

## 1.21 HIE

### 1. A straight

This is five cards in a sequence (e.g., 4,5,6,7,8), with aces allowed to be either 1 or 13 (low or high) and with the cards allowed to be of the same suit (e.g., all hearts) or from some different suits. The number of such hands is  $10 \times (4\text{choose}1)^5$

### 2. A flush

Here all 5 cards are from the same suit (they may also be a straight). The number of such hands is  $(4\text{-choose-}1)^* (13\text{-choose-}5)$ .

### 3. A full house

## 1.21 HIE

### 1. A straight

This is five cards in a sequence (e.g., 4,5,6,7,8), with aces allowed to be either 1 or 13 (low or high) and with the cards allowed to be of the same suit (e.g., all hearts) or from some different suits. The number of such hands is  $10 \times (4\text{choose}1)^5$

### 2. A flush

Here all 5 cards are from the same suit (they may also be a straight). The number of such hands is  $(4\text{-choose-}1)^* (13\text{-choose-}5)$ .

### 3. A full house

This hand has the pattern AAABB where A and B are from distinct kinds. The number of such hands is  $(13\text{-choose-}1)(4\text{-choose-}3)(12\text{-choose-}1)(4\text{-choose-}2)$

# Homework 1

## 1.21 HIE

1. A straight

# Homework 1

## 1.21 HIE

### 3. A straight

- This is five cards in a sequence (e.g., 4,5,6,7,8)
- with aces allowed to be either 1 or 13 (low or high) i.e. 10 such sequences, can't start on J,Q,K
- the cards allowed to be of the same suit (e.g., all hearts) or from some different suits i.e.  $(4 - \textit{choose} - 1)$
- number of such hands is  $10 \times (4 - \textit{choose} - 1)^5$
- Denominator is  $(52 - \textit{choose} - 5)$

### 2. A flush

## 1.21 HIE

### 3. A straight

- This is five cards in a sequence (e.g., 4,5,6,7,8)
- with aces allowed to be either 1 or 13 (low or high) i.e. 10 such sequences, can't start on J,Q,K
- the cards allowed to be of the same suit (e.g., all hearts) or from some different suits i.e.  $(4 - \textit{choose} - 1)$
- number of such hands is  $10 \times (4 - \textit{choose} - 1)^5$
- Denominator is  $(52 - \textit{choose} - 5)$

### 2. A flush

- Here all 5 cards are from the same suit (they may also be a straight). The number of such hands is  $(4 - \textit{choose} - 1) \times (13 - \textit{choose} - 5)$
- Denominator is  $(52 - \textit{choose} - 5)$

## 1.21 HIE

### 3. A full house

- This hand has the pattern AAABB where A and B are from distinct kinds.
- The number of such hands is  $(13 - \text{choose} - 1)(4 - \text{choose} - 3)(12 - \text{choose} - 1)(4 - \text{choose} - 2)$
- Denominator is  $(52 - \text{choose} - 5)$



# Homework 2 Hints

## 2.1 HIE

Let  $X \sim U[0, 1]$ . Find the PDF of  $Y = X^2$

# Homework 2 Hints

## 2.1 HIE

Let  $X \sim U[0, 1]$ . Find the PDF of  $Y = X^2$

- $Y = g(X)$  so,  $X = g^{-1}(Y) = Y^{1/2}$

# Homework 2 Hints

## 2.1 HIE

Let  $X \sim U[0, 1]$ . Find the PDF of  $Y = X^2$

- $Y = g(X)$  so,  $X = g^{-1}(Y) = Y^{1/2}$
- $Pr(Y < y) = F_y(y) = F_x(g^{-1}(y)) = y^{1/2}$

# Homework 2 Hints

## 2.1 HIE

Let  $X \sim U[0, 1]$ . Find the PDF of  $Y = X^2$

- $Y = g(X)$  so,  $X = g^{-1}(Y) = Y^{1/2}$
- $Pr(Y < y) = F_y(y) = F_x(g^{-1}(y)) = y^{1/2}$
- If CDF is continuous and differentiable, density is first derivative.
- $dF_y(y)/dy = f_y(y) = 1/2y^{-1/2}$

## 2.1 HIE

Let  $X \sim U[0, 1]$ . Find the PDF of  $Y = X^2$

- $Y = g(X)$  so,  $X = g^{-1}(Y) = Y^{1/2}$
- $Pr(Y < y) = F_y(y) = F_x(g^{-1}(y)) = y^{1/2}$
- If CDF is continuous and differentiable, density is first derivative.
- $dF_y(y)/dy = f_y(y) = 1/2y^{-1/2}$  where  $0 \leq Y \leq 1$
- $f_y(y) = 0$  otherwise
- Note the bounds are important. This is a PDF, density must integrate to 1.

## 2.8 HIE

Show that if the density satisfies  $f(x) = f(-x)$  for  $x \in \mathcal{R}$  then the distribution function satisfies  $F(-x) = 1 - F(x)$

## 2.8 HIE

Show that if the density satisfies  $f(x) = f(-x)$  for  $x \in \mathcal{R}$  then the distribution function satisfies  $F(-x) = 1 - F(x)$

- We know:  $f(t) = f(-t)$
- Also we know  $F(x) = \int_{-\infty}^x f(t)dt$  for a continuous RV
- $F(-x) = \int_{-\infty}^{-x} f(t)dt$
- $= \int_{-\infty}^{-x} f(-t)dt$  from what we are given
- letting  $s = -t$  i.e. change of variable
- $= \int_x^{\infty} f(s)ds$
- $= \int_{\infty}^{-\infty} f(s)ds - \int_x^{-\infty} f(s)ds$
- $= 1 - F(x)$

## 2.13 HIE

Find  $a$  which minimizes  $E[(X - a)^2]$ . Your answer should be an expectation or moment of  $X$



## 2.13 HIE

Find  $a$  which minimizes  $E[(X - a)^2]$ . Your answer should be an expectation or moment of  $X$

- $E((X - a)^2) = a^2 - 2E(X)a + E(X^2) = (a - E(X))^2 + E(X^2) - E(X)^2.$
- Considering this as a quadratic function on  $a$
- it is clear that the minimum is acquired when  $a = E(X).$

## 3.1 HIE

For the Bernoulli distribution show...

1.  $\sum_{x=0}^1 \pi(x; p) = 1$

## 3.1 HIE

For the Bernoulli distribution show...

1.  $\sum_{x=0}^1 \pi(x; p) = 1$ 
  - $(1 - p) + p = 1$

## 3.1 HIE

For the Bernoulli distribution show...

1.  $\sum_{x=0}^1 \pi(x; p) = 1$ 
  - $(1 - p) + p = 1$
  - This should be clear from the definition of the density function for discrete RV
2.  $E(x) = p$

## 3.1 HIE

For the Bernoulli distribution show...

1.  $\sum_{x=0}^1 \pi(x; p) = 1$ 
  - $(1 - p) + p = 1$
  - This should be clear from the definition of the density function for discrete RV
2.  $E(x) = p$ 
  - $E(x) = \sum_{x=0}^1 x \times \pi(x; p)$
  - $0 \times (1 - p) + 1 \times p = p$

## 3.1 HIE

For the Bernoulli distribution show...

1.  $\sum_{x=0}^1 \pi(x; p) = 1$ 
  - $(1 - p) + p = 1$
  - This should be clear from the definition of the density function for discrete RV
2.  $E(x) = p$ 
  - $E(x) = \sum_{x=0}^1 x \times \pi(x; p)$
  - $0 \times (1 - p) + 1 \times p = p$
3.  $Var(x) = p(1 - p)$

## 3.1 HIE

For the Bernoulli distribution show...

1.  $\sum_{x=0}^1 \pi(x; p) = 1$ 
  - $(1 - p) + p = 1$
  - This should be clear from the definition of the density function for discrete RV
2.  $E(x) = p$ 
  - $E(x) = \sum_{x=0}^1 x \times \pi(x; p)$
  - $0 \times (1 - p) + 1 \times p = p$
3.  $Var(x) = p(1 - p)$ 
  - $Var(x) = E(x^2) - E(x)^2$
  - $[(0)^2(1 - p) + (1)^2p] - p^2 = p - p^2 = p(1 - p)$

## 3.1 HIE

For the Bernoulli distribution show...

1.  $\sum_{x=0}^1 \pi(x; p) = 1$ 
  - $(1 - p) + p = 1$
  - This should be clear from the definition of the density function for discrete RV
2.  $E(x) = p$ 
  - $E(x) = \sum_{x=0}^1 x \times \pi(x; p)$
  - $0 \times (1 - p) + 1 \times p = p$
3.  $Var(x) = p(1 - p)$ 
  - $Var(x) = E(x^2) - E(x)^2$
  - $[(0)^2(1 - p) + (1)^2p] - p^2 = p - p^2 = p(1 - p)$