

Econometrics I - TA section

Eric Schulman

The University of Texas at Austin

November 10, 2020



TEXAS

The University of Texas at Austin

Gauss-Markov (Theorem 4.4) “OLS” is “BLUE” i.e. best linear unbiased estimator

Gauss-Markov (Theorem 4.4) “OLS” is “BLUE” i.e. best linear unbiased estimator

- (1) unbiased - expected value is “true”

$$E(\hat{\beta}|X) = \beta$$

- (2) best - minimum variance

$$\text{var}(\tilde{\beta}|X) \geq \text{var}(\hat{\beta}|X)$$

- Does it matter if there is 1 or 1000 obs for efficiency and unbiased-ness?

- Does it matter if there is 1 or 1000 obs for efficiency and unbiased-ness?
 - No
- Trade-off between bias and variance i.e. is a constant efficient and unbiased

- Does it matter if there is 1 or 1000 obs for efficiency and unbiased-ness?
 - No
- Trade-off between bias and variance i.e. is a constant efficient and unbiased
 - Variance is 0, but super biased
 - There are estimators that have less variance than OLS, but these are biased – Ridge regression

4.1

For some integer k set $\mu_k = E(y^k)$

- Construct an estimator $\hat{\mu}_k$ for μ_k

4.1

For some integer k set $\mu_k = E(y^k)$

- Construct an estimator $\hat{\mu}_k$ for μ_k

$$\hat{\mu}_k = \frac{1}{n} \sum_{i=1}^n y_i^k$$

4.1

For some integer k set $\mu_k = E(y^k)$

- Construct an estimator $\hat{\mu}_k$ for μ_k

$$\hat{\mu}_k = \frac{1}{n} \sum_{i=1}^n y_i^k$$

- Show $\hat{\mu}_k$ is unbiased for μ_k

4.1

For some integer k set $\mu_k = E(y^k)$

- Construct an estimator $\hat{\mu}_k$ for μ_k

$$\hat{\mu}_k = \frac{1}{n} \sum_{i=1}^k y_i^k$$

- Show $\hat{\mu}_k$ is unbiased for μ_k

$$E(\hat{\mu}_k) = E\left(\frac{1}{n} \sum_{i=1}^n y_i^k\right) = \frac{1}{n} \sum_{i=1}^n E(y_i^k) = E(y_i^k)$$

4.1

For some integer k set $\mu_k = E(y^k)$

- Construct an estimator $\hat{\mu}_k$ for μ_k

$$\hat{\mu}_k = \frac{1}{n} \sum_{i=1}^k y_i^k$$

- Show $\hat{\mu}_k$ is unbiased for μ_k

$$E(\hat{\mu}_k) = E\left(\frac{1}{n} \sum_{i=1}^n y_i^k\right) = \frac{1}{n} \sum_{i=1}^n E(y_i^k) = E(y_i^k)$$

- Calculate the variance of $\hat{\mu}_k$ say $\text{var}(\hat{\mu}_k)$. What assumption is needed for $\text{var}(\hat{\mu}_k)$ to be finite?

- Calculate the variance of $\hat{\mu}_k$ say $var(\hat{\mu}_k)$. What assumption is needed for $var(\hat{\mu}_k)$ to be finite?

$$var(\hat{\mu}_k) = var\left(\frac{1}{n} \sum_{i=1}^n y_i^k\right) = \frac{1}{n^2} \sum_{i=1}^n var(y_i^k) = \frac{1}{n} var(y_i^k)$$

- Calculate the variance of $\hat{\mu}_k$ say $var(\hat{\mu}_k)$. What assumption is needed for $var(\hat{\mu}_k)$ to be finite?

$$var(\hat{\mu}_k) = var\left(\frac{1}{n} \sum_{i=1}^n y_i^k\right) = \frac{1}{n^2} \sum_{i=1}^n var(y_i^k) = \frac{1}{n} var(y_i^k)$$

We need $var(y_i^k)$ to be finite

- Calculate the variance of $\hat{\mu}_k$ say $var(\hat{\mu}_k)$. What assumption is needed for $var(\hat{\mu}_k)$ to be finite?

$$var(\hat{\mu}_k) = var\left(\frac{1}{n} \sum_{i=1}^n y_i^k\right) = \frac{1}{n^2} \sum_{i=1}^n var(y_i^k) = \frac{1}{n} var(y_i^k)$$

We need $var(y_i^k)$ to be finite

- Propose an estimator of $var(\hat{\mu}_k)$

- Calculate the variance of $\hat{\mu}_k$ say $\text{var}(\hat{\mu}_k)$. What assumption is needed for $\text{var}(\hat{\mu}_k)$ to be finite?

$$\text{var}(\hat{\mu}_k) = \text{var}\left(\frac{1}{n} \sum_{i=1}^n y_i^k\right) = \frac{1}{n^2} \sum_{i=1}^n \text{var}(y_i^k) = \frac{1}{n} \text{var}(y_i^k)$$

We need $\text{var}(y_i^k)$ to be finite

- Propose an estimator of $\text{var}(\hat{\mu}_k)$

$$\frac{1}{n^2} \left(\sum_{i=1}^n (y_i^k - \hat{\mu}_k)^2 \right)$$

- Calculate the variance of $\hat{\mu}_k$ say $var(\hat{\mu}_k)$. What assumption is needed for $var(\hat{\mu}_k)$ to be finite?

$$var(\hat{\mu}_k) = var\left(\frac{1}{n} \sum_{i=1}^n y_i^k\right) = \frac{1}{n^2} \sum_{i=1}^n var(y_i^k) = \frac{1}{n} var(y_i^k)$$

We need $var(y_i^k)$ to be finite

- Propose an estimator of $var(\hat{\mu}_k)$

$$\frac{1}{n^2} \left(\sum_{i=1}^n (y_i^k - \hat{\mu}_k)^2 \right)$$

4.7

Let $\tilde{\beta}$ be the GLS estimator. Assume that $\Omega = c^2\Sigma$ with Σ known and c^2 unknown. Define the residual vector $\tilde{e} = y - X\tilde{\beta}$ and

$$\tilde{c}^2 = \frac{1}{n - k} \tilde{e}'\Sigma^{-1}\tilde{e}$$

4.7

Let $\tilde{\beta}$ be the GLS estimator. Assume that $\Omega = c^2\Sigma$ with Σ known and c^2 unknown. Define the residual vector $\tilde{e} = y - X\tilde{\beta}$ and

$$\tilde{c}^2 = \frac{1}{n - k} \tilde{e}' \Sigma^{-1} \tilde{e}$$

- Show (4.18) $E(\tilde{\beta}|X) = \beta$

4.7

Let $\tilde{\beta}$ be the GLS estimator. Assume that $\Omega = c^2\Sigma$ with Σ known and c^2 unknown. Define the residual vector $\tilde{e} = y - X\tilde{\beta}$ and

$$\tilde{c}^2 = \frac{1}{n - k} \tilde{e}'\Sigma^{-1}\tilde{e}$$

- Show (4.18) $E(\tilde{\beta}|X) = \beta$

$$\begin{aligned} E(\tilde{\beta}|X) &= E((X'\Sigma^{-1}X)^{-1}(X'\Sigma^{-1}y)|X) \\ &= E((X'\Sigma^{-1}X)^{-1}(X'\Sigma^{-1}(X\beta + e))|X) \\ &= \beta + E((X'\Sigma^{-1}X)^{-1}X'\Sigma^{-1}e|X) = \beta \end{aligned}$$

4.7

- Show (4.19) $\text{var}(\tilde{\beta}|X) = (X'\Omega^{-1}X)^{-1}$

4.7

- Show (4.19) $\text{var}(\tilde{\beta}|X) = (X'\Omega^{-1}X)^{-1}$

$$\begin{aligned}\text{var}(\tilde{\beta}|X) &= \text{var}((X'\Sigma^{-1}X)^{-1}(X'\Sigma^{-1}y)|X) \\ &= \text{var}((X'\Sigma^{-1}X)^{-1}(X'\Sigma^{-1}(X\beta + e))|X) \\ &= \text{var}((X'\Sigma^{-1}X)^{-1}X'\Sigma^{-1}e|X) \\ &= (X'\Sigma^{-1}X)^{-1}X'\Sigma^{-1}\Omega(X'\Sigma^{-1})'((X'\Sigma^{-1}X)^{-1})' \\ &= (X'\Omega^{-1}X)^{-1}\end{aligned}$$

- Note

$$(A^{-1})' = (A')^{-1}$$

$$(AB)' = B'A'$$

4.23 Take the linear regression model with $E(y|X) = X\beta$. Define the ridge regression estimator $\hat{\beta} = (X'X + I_k\lambda)^{-1}X'y$ where λ is a fixed constant. Find $E(\hat{\beta}|X)$. Is $\hat{\beta}$ biased for β ?

4.23 Take the linear regression model with $E(y|X) = X\beta$. Define the ridge regression estimator $\hat{\beta} = (X'X + I_k\lambda)^{-1}X'y$ where λ is a fixed constant. Find $E(\hat{\beta}|X)$. Is $\hat{\beta}$ biased for β ?

$$\begin{aligned} E(X'X + I_k\lambda)^{-1}X'y|X) &= E((X'X)^{-1}X'y + I_k\lambda^{-1}X'y|X) \\ &= E((X'X)^{-1}X'X\beta + I_k\lambda^{-1}X'X\beta|X) = \beta + \lambda I_k X'X\beta \end{aligned}$$

Clearly it is biased.

- Asymptotic analysis. Now n comes into play.
- Consistency - kind of like being unbiased.
- Asymptotic normality - Normal “approximates” the true distribution.
- Allows us to make statements about how likely our estimator is.

Consistency/LLN

- Consistency

$$\hat{\theta} \rightarrow \theta$$

- Weak LLN (1) x_i are i.i.d. (2) and $E|x_i| < \infty$. Then as $n \rightarrow \infty$

$$\frac{1}{n} \sum_{i=1}^n x_i \xrightarrow{p} \mu$$

- As the sample gets large, variance gets small...
- Then apply Markov's inequality

Asymptotic Normality/CLT Lindbergh-Levy CLT: (1) x_i i.i.d.
and (2) $E(x_i^2) < \infty$ then

$$\sqrt{n}(\bar{x} - \mu) \xrightarrow{d} \mathcal{N}(0, \sigma^2)$$

Asymptotic Normality/CLT Lindbergh-Levy CLT: (1) x_i i.i.d.
and (2) $E(x_i^2) < \infty$ then

$$\sqrt{n}(\bar{x} - \mu) \xrightarrow{d} \mathcal{N}(0, \sigma^2)$$

- Do we know μ ? i.e. the population mean?

Asymptotic Normality/CLT Lindbergh-Levy CLT: (1) x_i i.i.d.
and (2) $E(x_i^2) < \infty$ then

$$\sqrt{n}(\bar{x} - \mu) \xrightarrow{d} \mathcal{N}(0, \sigma^2)$$

- Do we know μ ? i.e. the population mean?
- No! We usually need to choose it

Summarizing - “4 properties” you’ll be asked about

- Does not depend on n
 - Unbiased
 - Efficient
- Depends on n
 - Consistent
 - Asymptotically normal

7.9 Take the model

$$y_i = x_i\beta + e_i$$

$$E(e_i|x_i) = 0$$

Consider two estimators

$$\hat{\beta} = \sum_{i=1}^n \frac{\sum_{i=1}^n x_i y_i}{\sum_{i=1}^n x_i^2}$$

$$\tilde{\beta} = \frac{1}{n} \sum_{i=1}^n \frac{y_i}{x_i}$$

7.9

- Under stated assumptions, are both estimators consistent for β ?

7.9

- Under stated assumptions, are both estimators consistent for β ?

First compute $E(\hat{\beta}|X) = E\left(\frac{\sum_{i=1}^n x_i y_i}{\sum_{i=1}^n x_i^2} \mid X\right) =$

$$E\left(\frac{\sum_{i=1}^n x_i (\beta x_i + e_i)}{\sum_{i=1}^n x_i^2} \mid X\right) = E\left(\beta \frac{\sum_{i=1}^n x_i^2}{\sum_{i=1}^n x_i^2} \mid X\right) = \beta$$

This establishes unbiasedness i.e.

$E(\hat{\beta}) = E(E(\hat{\beta}|X)) = E(\beta) = \beta$. Now can apply the weak LLN.

7.9

- Under stated assumptions, are both estimators consistent for β ?

7.9

- Under stated assumptions, are both estimators consistent for β ?

$$E(\tilde{\beta}|X) = E\left(\frac{1}{n} \sum_{i=1}^n \frac{y_i}{x_i} \mid X\right) =$$

$$E\left(\frac{1}{n} \sum_{i=1}^n \frac{x_i\beta + e_i}{x_i} \mid X\right) = \beta$$

Similarly, can apply the weak LLN.

7.9

- Under stated assumptions, are both estimators consistent for β ?

$$E(\tilde{\beta}|X) = E\left(\frac{1}{n} \sum_{i=1}^n \frac{y_i}{x_i} \mid X\right) =$$

$$E\left(\frac{1}{n} \sum_{i=1}^n \frac{x_i\beta + e_i}{x_i} \mid X\right) = \beta$$

Similarly, can apply the weak LLN.

- Are there conditions under which either estimator is efficient?
Yes Gauss-Markov, (1) is the OLS estimator

7.14

Take the model

$$y_i = x_{1i}\beta_1 + x_{2i}\beta_2 + e_i$$

$$E(x_i e_i) = 0$$

Where β_1 and β_2 are scalars. Define $\theta = \beta_1\beta_2$

7.14

- What is the appropriate estimator $\hat{\theta}$ for θ

7.14

- What is the appropriate estimator $\hat{\theta}$ for $\theta \begin{bmatrix} \hat{\beta}_1 \\ \hat{\beta}_2 \end{bmatrix} = (X'X)^{-1}X'y$

Then

$\hat{\beta}_1\hat{\beta}_2 = \hat{\theta}$. Can show consistency using Slutsky's theorem i.e. $\hat{\beta}_1$ and $\hat{\beta}_2$ converge to a constant in probability. Their product also converges.

7.14

- What is the appropriate estimator $\hat{\theta}$ for $\theta \begin{bmatrix} \hat{\beta}_1 \\ \hat{\beta}_2 \end{bmatrix} = (X'X)^{-1}X'y$

Then

$\hat{\beta}_1\hat{\beta}_2 = \hat{\theta}$. Can show consistency using Slutsky's theorem i.e. $\hat{\beta}_1$ and $\hat{\beta}_2$ converge to a constant in probability. Their product also converges.

- Note, convergence in probability is stronger than convergence in distribution. As a result, the conditions for Slutsky's theorem (as written in the textbook) apply.

7.14

- Find the asymptotic distribution of $\hat{\theta}$ under standard regularity conditions.

7.14

- Find the asymptotic distribution of $\hat{\theta}$ under standard regularity conditions.

$$\begin{bmatrix} \hat{\beta}_1 \\ \hat{\beta}_2 \end{bmatrix} \rightarrow \mathcal{N}(0, \Sigma)$$

Need to compute a change of variable for $\hat{\theta} = \hat{\beta}_1 \hat{\beta}_2$. Similar to 4.8 on previous homework.

7.14

- Find the asymptotic distribution of $\hat{\theta}$ under standard regularity conditions.

$$\begin{bmatrix} \hat{\beta}_1 \\ \hat{\beta}_2 \end{bmatrix} \rightarrow \mathcal{N}(0, \Sigma)$$

Need to compute a change of variable for $\hat{\theta} = \hat{\beta}_1 \hat{\beta}_2$. Similar to 4.8 on previous homework.

- Show how to calculate an asymptotic 95% confidence interval for θ

7.14

- Find the asymptotic distribution of $\hat{\theta}$ under standard regularity conditions.

$$\begin{bmatrix} \hat{\beta}_1 \\ \hat{\beta}_2 \end{bmatrix} \rightarrow \mathcal{N}(0, \Sigma)$$

Need to compute a change of variable for $\hat{\theta} = \hat{\beta}_1 \hat{\beta}_2$. Similar to 4.8 on previous homework.

- Show how to calculate an asymptotic 95% confidence interval for θ

Need to solve for θ where $Pr(|\hat{\beta}_1 \hat{\beta}_2| \leq \theta) = .95$

7.14

- Find the asymptotic distribution of $\hat{\theta}$ under standard regularity conditions.

$$\begin{bmatrix} \hat{\beta}_1 \\ \hat{\beta}_2 \end{bmatrix} \rightarrow \mathcal{N}(0, \Sigma)$$

Need to compute a change of variable for $\hat{\theta} = \hat{\beta}_1 \hat{\beta}_2$. Similar to 4.8 on previous homework.

- Show how to calculate an asymptotic 95% confidence interval for θ

Need to solve for θ where $Pr(|\hat{\beta}_1 \hat{\beta}_2| \leq \theta) = .95$