## Lecture 12: Inference for Two Proportions STAT 630, Fall 2021

Let  $X_1, X_2, \dots, X_{n_1}$  be independent random variables from Bern $(p_1)$ , and  $Y_1, Y_2, \dots, Y_{n_2}$  be independent random variables from Bern $(p_2)$ . Assume that the two samples are independent of each other.

The sample proportions are defined as:

$$\hat{p}_1 = \frac{1}{n_1} \sum_{i=1}^{n_1} X_i$$
 and  $\hat{p}_2 = \frac{1}{n_2} \sum_{i=1}^{n_2} Y_i$ 

Find the expectation and variance for the difference between two proportions:

According to the central limit theorem, the sampling distribution for the difference between two proportions can be approximated by a normal distribution:

$$\hat{p}_1 - \hat{p}_2 \sim N \left( p_1 - p_2, \sqrt{\frac{p_1(1-p_1)}{n_1} + \frac{p_2(1-p_2)}{n_2}} \right)$$

The approximation should only be used if  $n_i p_i \ge 10$  and  $n_i (1 - p_i) \ge 10$  for i = 1, 2. That is, there are at least that 10 "successes" and "failures" for each group.

 $1-\alpha$  confidence interval for  $p_1-p_2$ 

$$\hat{p}_1 - \hat{p}_2 \pm z_{\alpha/2} \sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n_1} + \frac{\hat{p}_2(1-\hat{p}_2)}{n_2}}$$

Conditions:

- $n_i \hat{p}_i \ge 10$  and  $n_i (1 \hat{p}_i) \ge 10$  for i = 1, 2
- The data are independent within and between groups. Generally this is satisfied if the data come from two independent random samples, or if the data come from a randomized experiment.

Ex1: A 5-year study was conducted to evaluate the effectiveness of fish oils on reducing cardiovascular events, where each subject was randomized into one of two treatment groups. We'll consider heart attack outcomes in these patients:

|          | heart attack | no event | Total |
|----------|--------------|----------|-------|
| fish oil | 145          | 12788    | 12933 |
| placebo  | 200          | 12738    | 12938 |

- (a) Is this study an experiment or an observational study?
- (b) Are the conditions for inference satisfied?

(c) Calculate a 95% confidence interval for the difference in heart attack rates of patients that take fish oils and patients that take a placebo. Let's use R to do this:

```
n_fish <- 12933
phat_fish <- 145 / 12933; phat_fish

## [1] 0.01121163

n_placebo <- 12938
phat_placebo <- 200 / 12938; phat_placebo

## [1] 0.01545834

z_crit <- qnorm(0.975); z_crit

## [1] 1.959964

SE <- sqrt(phat_fish*(1-phat_fish)/n_fish + phat_placebo*(1-phat_placebo)/n_placebo)
ci_lower <- phat_fish - phat_placebo - z_crit * SE
ci_upper <- phat_fish - phat_placebo + z_crit * SE
round(c(ci_lower, ci_upper), 4)

## [1] -0.0070 -0.0015</pre>
```

| (d) Interpret the confidence interval in the context of | the data. |
|---|-----------|
|   |           |
|   |           |
|   |           |

## Hypothesis test for the difference between two proportions

Null and alternative hypothesis:

 $H_0$ :  $p_1 = p_2$  (the two population proportions are the same)

 $H_A$ :  $p_1 > p_2$  or  $p_1 < p_2$  or  $p_1 \neq p_2$ 

Test statistic: Assuming  $H_0$  is true, the two proportions are equal:  $p_1 = p_2 = p$ . Under this assumption, the variance is given by

We estimate p with the pooled proportion:

$$\hat{p} = \frac{n_1 \hat{p}_1 + n_2 \hat{p}_2}{n_1 + n_2}$$

Conclusively, the z-test statistic is given by

$$z = \frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\hat{p}(1-\hat{p})(1/n_1 + 1/n_2)}}$$

Once the z-test statistic is computed, the p-value can be computed to decide whether to reject or not reject the null hypothesis.

The conditions are the same as for confidence intervals except we use the pooled proportion  $\hat{p}$  when checking the number successes and failures (i.e.,  $n_i \hat{p} \geq 10$  and  $n_i (1 - \hat{p}) \geq 10$  for i = 1, 2).

**Ex2**: Are young people more concerned with environmental issues than older people? The Pew Research Center for the People and the Press conducted a survey in 2009 asking a random sample of adults whether or not there is solid evidence of global warming. Of the 197 people aged 18–29 years old who responded, 126 said yes, compared to 223 out of 406 people aged 30–39 years old. Does this indicate that a higher proportion of younger people believe there is evidence of global warming? <sup>1</sup>

<sup>&</sup>lt;sup>1</sup>http://www.pewresearch.org/2009/10/22/fewer-americans-see-solid-evidence-of-global-warming/