

Lecture 10: Properties of Least Squares Estimates

STAT 632, Spring 2020

Preliminaries: Let a and b be constants, and X a random variable.

$$\begin{aligned} E(aX + b) &= aE(X) + b \\ \text{Var}(aX + b) &= a^2 \text{Var}(X) \end{aligned}$$

Random Vectors

Let X_1, X_2, \dots, X_n be random variables. An $n \times 1$ random vector is given by¹

$$\mathbf{X} = \begin{pmatrix} X_1 \\ X_2 \\ \vdots \\ X_n \end{pmatrix}$$

The **variance-covariance** matrix of \mathbf{X} is the $n \times n$ symmetric matrix

$$\text{Var}(\mathbf{X}) = \begin{pmatrix} \text{Var}(X_1) & \text{Cov}(X_1, X_2) & \cdots & \text{Cov}(X_1, X_n) \\ \text{Cov}(X_2, X_1) & \text{Var}(X_2) & \cdots & \text{Cov}(X_2, X_n) \\ \vdots & \vdots & \ddots & \vdots \\ \text{Cov}(X_n, X_1) & \text{Cov}(X_n, X_2) & \cdots & \text{Var}(X_n) \end{pmatrix}$$

Note that $\text{Cov}(X_i, X_i) = \text{Var}(X_i)$.

Properties of Expectation and Variance for Random Vectors

Let \mathbf{A} be an $m \times n$ matrix of constants, \mathbf{b} an $m \times 1$ vector of constants, and \mathbf{X} an $n \times 1$ random vector, then

$$\begin{aligned} E(\mathbf{AX} + \mathbf{b}) &= \mathbf{A}E(\mathbf{X}) + \mathbf{b} \\ \text{Var}(\mathbf{AX} + \mathbf{b}) &= \mathbf{A}\text{Var}(\mathbf{X})\mathbf{A}' \end{aligned}$$

$$E(\mathbf{X}) = \begin{pmatrix} E(X_1) \\ E(X_2) \\ \vdots \\ E(X_n) \end{pmatrix}$$

$$\begin{aligned} \text{Cov}(X, Y) &= E[(X - E(X))(Y - E(Y))] \\ &= E(XY) - E(X)E(Y) \\ \text{Sample Covariance} &= \frac{\sum (x_i - \bar{x})(y_i - \bar{y})}{n-1} \end{aligned}$$

$$\begin{aligned} R: \text{Cov}(x, y) &= \frac{s_{xy}}{s_x s_y} \\ \text{Cor}(x, y) &= \frac{s_{xy}}{s_x s_y} \end{aligned}$$

¹Note that the \mathbf{X} here is different than the $n \times (p+1)$ model matrix in regression.

Properties of the Multiple Linear Regression Model

$$Y = X\beta + e$$

- Y is an $n \times 1$ response vector
- X is an $n \times (p+1)$ design matrix for the predictors
- β is a $(p+1) \times 1$ vector of regression parameters
- e is an $n \times 1$ vector of random errors; assuming $e_i \sim N(0, \sigma^2)$, independently

The variance-covariance matrix for the random errors e is given by

$$\text{Var}(e) = \begin{pmatrix} \sigma^2 & 0 & \cdots & 0 \\ 0 & \sigma^2 & \cdots & 0 \\ \vdots & & & \\ 0 & 0 & \cdots & \sigma^2 \end{pmatrix} = \sigma^2 I_n$$

which follows since $\text{Var}(e_i) = \sigma^2$, and $\text{Cov}(e_i, e_j) = 0$ when $i \neq j$ (by independence).

Use the properties of expectation and variance for random vectors to find $E(Y)$, $\text{Var}(Y)$, and the distribution of Y .

$$E(Y) = E(\underbrace{X\beta}_{\text{constant}} + \underbrace{e}_0) = X\beta + E(e) = X\beta$$

$$\text{Var}(Y) = \text{Var}(X\beta + e) = \text{Var}(e) = \sigma^2 I_n$$

$$Y \sim N(X\beta, \sigma^2 I_n)$$

Properties of Least Squares Estimates

Recall that the least squares estimates of the population parameters β is given by the $(p+1) \times 1$ vector

$$\hat{\beta} = (X'X)^{-1}X'Y$$

The estimator $\hat{\beta}$ is a random vector since it varies from sample to sample.

Show that $E(\hat{\beta}) = \beta$, the least squares estimates are unbiased.

$$\begin{aligned}
 E(\hat{\beta}) &= E[(X'X)^{-1}X'\epsilon] \\
 &= (X'X)^{-1}X'E(\epsilon) \\
 &= (X'X)^{-1}X'X\beta \\
 &= \beta
 \end{aligned}
 \quad \text{Since } A^{-1}A = I$$

Show that the variance-covariance matrix for the least squares estimates is given by $\text{Var}(\hat{\beta}) = \sigma^2(X'X)^{-1}$

$$\begin{aligned}
 \text{Var}(\hat{\beta}) &= \text{Var}[(X'X)^{-1}X'\epsilon] \\
 &= (X'X)^{-1}X' \text{Var}(\epsilon) [(X'X)^{-1}X']' \\
 &= (X'X)^{-1}X'(\sigma^2 I_n)X(X'X)^{-1} \\
 &= \sigma^2 (X'X)^{-1}X'X(X'X)^{-1} \quad \text{since } (A^{-1})' = (A')^{-1} \\
 &= \sigma^2 (X'X)^{-1} \quad \begin{aligned} &[(X'X)^{-1}]' \\ &= [(X'X)']^{-1} \\ &= (X'X)^{-1} \end{aligned}
 \end{aligned}$$

The variance of a particular estimate is given by the diagonal entry:

$$\text{Var}(\hat{\beta}_j) = \sigma^2 (X'X)^{-1}_{j+1,j+1} \quad \text{for } j = 0, \dots, p$$

This is how standard errors are calculated when constructing confidence intervals for the parameters. For example, a $1 - \alpha$ confidence interval for $\hat{\beta}_j$ is given by

$$\hat{\beta}_j \pm t_{\alpha/2; n-p-1} \text{se}(\hat{\beta}_j)$$

where $\text{se}(\hat{\beta}_j) = \hat{\sigma} \sqrt{(X'X)^{-1}_{j+1,j+1}}$, and $\hat{\sigma} = \sqrt{\text{RSS}/(n-p-1)}$

Using the menu pricing data set (lecture 7), the code below demonstrates how to manually compute the least squares estimates, variance-covariance matrix, and standard errors. The results are compared to the output from `lm()`. Note that it is always better to use `lm()` than to do the computations manually. I am just showing this to verify all the formulas.

```
nyc <- read.csv("https://ericwfox.github.io/data/nyc.csv")

# response vector
Y <- matrix(nyc$Price, ncol=1)

# design matrix
X <- cbind(Intercept = 1, nyc[,c('Food', 'Decor', 'East')])
X <- as.matrix(X)
rownames(X) <- nyc$Restaurant
X[1:5,]

##               Intercept Food Decor East
## Daniella Ristorante      1   22   18   0
## Tello's Ristorante       1   20   19   0
## Biricchino               1   21   13   0
## Bottino                  1   20   20   0
## Da Umberto               1   24   19   0

# manually calculate least squares estimates
betaHat <- solve(t(X) %*% X) %*% t(X) %*% Y
betaHat

##               [,1]
## Intercept -24.026880
## Food      1.536346
## Decor     1.909373
## East      2.067013

# compare with lm()
lm1 <- lm(Price ~ Food + Decor + East, data=nyc)
coef(lm1)

## (Intercept)      Food      Decor      East
## -24.026880   1.536346   1.909373   2.067013
```

```

# manually calculate standard errors for least squares estimates
n <- nrow(nyc)
p <- 3
resid <- as.numeric(Y - X %*% betaHat)
sigmaHat2 <- sum(resid^2) / (n-p-1)
covBetaHat <- sigmaHat2 * solve(t(X) %*% X)
covBetaHat

##           Intercept           Food           Decor           East
## Intercept 21.8344883 -0.94964667 -0.12475347  0.19880619
## Food      -0.9496467  0.06926176 -0.02530816 -0.04612459
## Decor     -0.1247535 -0.02530816  0.03610588  0.01149201
## East       0.1988062 -0.04612459  0.01149201  0.86827717

seBetaHat <- sqrt(diag(covBetaHat))
seBetaHat

## Intercept      Food      Decor      East
## 4.6727388 0.2631763 0.1900155 0.9318139

# compare with lm()
summary(lm1)$coef

##           Estimate Std. Error  t value    Pr(>|t|)
## (Intercept) -24.026880  4.6727388 -5.141926 7.669997e-07
## Food         1.536346  0.2631763  5.837705 2.758775e-08
## Decor        1.909373  0.1900155 10.048513 8.201940e-19
## East         2.067013  0.9318139  2.218268 2.791009e-02

vcov(lm1)

##           (Intercept)           Food           Decor           East
## (Intercept) 21.8344883 -0.94964667 -0.12475347  0.19880619
## Food      -0.9496467  0.06926176 -0.02530816 -0.04612459
## Decor     -0.1247535 -0.02530816  0.03610588  0.01149201
## East       0.1988062 -0.04612459  0.01149201  0.86827717

```