

# FROM LAGRANGE TO FEYNMAN

A different approach

---

Eric Yeung

November 24, 2014

PHYC54H3

# LAGRANGE

---

$$\mathcal{L} = T - V$$

- In PHYC54's section on variational calculus, we calculated the action  $\mathcal{S} = \int_{t_1}^{t_2} \mathcal{L}[x, \dot{x}; t] dt$
- This  $\mathcal{S}$  is fundamental in Feynman's path integral

SCHRÖDINGER

---

$$\left( \frac{-\hbar^2}{2m} \nabla^2 + V \right) \Psi = i\hbar \frac{\partial \Psi}{\partial t}$$

- $V$  is just the associated potential

$$\left( \frac{-\hbar^2}{2m} \nabla^2 + V \right) \Psi = i\hbar \frac{\partial \Psi}{\partial t}$$

- $V$  is just the associated potential
- $\nabla^2$  is the spatial Laplacian

$$\left( \frac{-\hbar^2}{2m} \nabla^2 + V \right) \Psi = i\hbar \frac{\partial \Psi}{\partial t}$$

- $V$  is just the associated potential
- $\nabla^2$  is the spatial Laplacian
- $\hbar$  is the reduced Planck constant

# TIME-DEPENDENT SCHRÖDINGER EQUATION

$$\left( \frac{-\hbar^2}{2m} \nabla^2 + V \right) \Psi = i\hbar \frac{\partial \Psi}{\partial t}$$

- $V$  is just the associated potential
- $\nabla^2$  is the spatial Laplacian
- $\hbar$  is the reduced Planck constant
- $\Psi$  is the position-space wave-function



- Recall that we solve Schrödinger's equation by finding the energy eigenvalues
- We dealt in the quantity called the wave-function
- In the path integral approach, we deal in the quantity known as the kernel

# FEYNMAN PATH INTEGRAL

---

## AN ANALOGY: BLAST TO THE PAST!




Figure: Two slits



⋮


Figure: Infinite slits


One can think of this approach as an infinite slit experiment.

- The double slit experiment was to demonstrate that light and matter could behave like classical particles AND waves.
- The event of an electron going from a source to a destination has one amplitude while passing through one of the two slits and has another amplitude for passing through the other slit
- The total amplitude of this event is then the superposition of the two amplitudes from the two respective slits.

## CONTRIBUTING PATHS

- In quantum mechanics (unlike classical mechanics), there is not just a single path (of least action) contributing to the amplitude.
- In addition to the classical path, there are paths that fluctuate around it.
- This is known as quantum fluctuation  $x(t) = x_c(t) + \delta x(t)$
- This differentiates classical and quantum particles, a classical particle cannot escape a potential barrier, but a quantum particle can tunnel through





**Figure:** Case A includes only the classical trajectory while case B includes all possible paths.

- The contributions from every path is equal in magnitude, but occur at different times

- The contributions from every path is equal in magnitude, but occur at different times

·

$$\phi[x(t)] = \text{constant } e^{i\mathcal{S}[x(t)]/\hbar} \quad (1)$$



- The contributions from every path is equal in magnitude, but occur at different times

·

$$\phi[x(t)] = \text{constant } e^{i\mathcal{S}[x(t)]/\hbar} \quad (1)$$

·

$$K(b, a) = \sum_{\text{paths from } a \text{ to } b} \phi[x(t)] \quad (2)$$

$$K(b, a) = \int \mathcal{D}x(t) e^{i S[b, a] / \hbar}$$

- $K(b, a)$  is the kernel, the propagator, or the total amplitude

$$K(b, a) = \int \mathcal{D}x(t) e^{i S[b, a] / \hbar}$$

- $K(b, a)$  is the kernel, the propagator, or the total amplitude
- $x_b, x_a$  are the initial and final positions

$$K(b, a) = \int \mathcal{D}x(t) e^{i S[b, a] / \hbar}$$

- $K(b, a)$  is the kernel, the propagator, or the total amplitude
- $x_b, x_a$  are the initial and final positions
- $t_b, t_a$  are the initial and final times

$$K(b, a) = \int \mathcal{D}x(t) e^{i S[b, a] / \hbar}$$

- $K(b, a)$  is the kernel, the propagator, or the total amplitude
- $x_b, x_a$  are the initial and final positions
- $t_b, t_a$  are the initial and final times
- $S$  is the classical action found from  $\mathcal{L}$

$$K(b, a) = \int \mathcal{D}x(t) e^{i\mathcal{S}[b,a]/\hbar}$$

- $K(b, a)$  is the kernel, the propagator, or the total amplitude
- $x_b, x_a$  are the initial and final positions
- $t_b, t_a$  are the initial and final times
- $\mathcal{S}$  is the classical action found from  $\mathcal{L}$
- $\mathcal{D}$  is the weight of all the possible paths

- As  $\hbar \rightarrow 0$ , the phase starts to oscillate

- As  $\hbar \rightarrow 0$ , the phase starts to oscillate
- As a result, most of the paths are cancelled out.



- As  $\hbar \rightarrow 0$ , the phase starts to oscillate
- As a result, most of the paths are cancelled out.
- This is because of the small  $\hbar$  limit; neighbouring paths will have slightly varying actions at different phases- thus constructively interfering.

## THINGS TO NOTE

- As  $\hbar \rightarrow 0$ , the phase starts to oscillate
- As a result, most of the paths are cancelled out.
- This is because of the small  $\hbar$  limit; neighbouring paths will have slightly varying actions at different phases- thus constructively interfering.
- The path that does survive makes the action stationary or  $\delta \mathcal{S} = 0$ , it has the biggest contribution in the small  $\hbar$  limit

## THINGS TO NOTE


- As  $\hbar \rightarrow 0$ , the phase starts to oscillate
- As a result, most of the paths are cancelled out.
- This is because of the small  $\hbar$  limit; neighbouring paths will have slightly varying actions at different phases- thus constructively interfering.
- The path that does survive makes the action stationary or  $\delta \mathcal{S} = 0$ , it has the biggest contribution in the small  $\hbar$  limit

$$K(b, a) = Fe^{i\mathcal{S}_{cl}/\hbar} \quad (3)$$

## PATH INTEGRAL FOR MORE THAN ONE EVENT

- The amplitude of a path with the mapping  $a \rightarrow c \rightarrow b$  is the sum of the sum of the amplitudes from  $a \rightarrow c$  and the one from  $c \rightarrow b$ .
- The action for multiple events is

$$\mathcal{S}[b, a] = \mathcal{S}[b, c] + \mathcal{S}[c, a]$$



**Figure:** A path with multiple events.

- $K(b, a) = \int_a^b \mathcal{D}x(t) e^{i(S[b,c] + S[c,a])/\hbar}$
- $K(b, a) = \int_{-\infty}^{\infty} \int_c^b \mathcal{D}x(t) K(c, a) e^{iS[b,c]/\hbar} dx_c$
- $K(b, a) = \int_{-\infty}^{\infty} K(b, c) K(c, a) dx_c$

## FAMILIAR EXAMPLES

---

- The Lagrangian is trivially  $\mathcal{L} = m/2(\dot{x}^2)$

- The Lagrangian is trivially  $\mathcal{L} = m/2(\dot{x}^2)$

•

$$x_c(t) = x_a + \frac{x_b - x_a}{t_b - t_a}(t - t_a) \quad (4)$$



- The Lagrangian is trivially  $\mathcal{L} = m/2(\dot{x}^2)$

•

$$x_c(t) = x_a + \frac{x_b - x_a}{t_b - t_a}(t - t_a) \quad (4)$$

•

$$S = \frac{m}{2} \frac{(x_b - x_a)^2}{t_b - t_a} + \frac{m}{4} \sum_{n=1}^{\infty} \frac{(n\pi)^2}{t_b - t_a} a_n^2 \quad (5)$$

- The Lagrangian is trivially  $\mathcal{L} = m/2(\dot{x}^2)$


- $$x_c(t) = x_a + \frac{x_b - x_a}{t_b - t_a}(t - t_a) \quad (4)$$

- $$\mathcal{S} = \frac{m}{2} \frac{(x_b - x_a)^2}{t_b - t_a} + \frac{m}{4} \sum_{n=1}^{\infty} \frac{(n\pi)^2}{t_b - t_a} a_n^2 \quad (5)$$

- $$K_0(b, a) = \left( \frac{m}{2\pi i \hbar (t_b - t_a)} \right)^{1/2} \exp \left[ \frac{im(x_b - x_a)^2}{2\hbar(t_b - t_a)} \right] \quad (6)$$

## FINDING THE CLASSICAL MOMENTUM

- Fix time, vary distance




**Figure:** The plot of the real part of  $\sqrt{i}$  times our kernel with varying distance.

$$2\pi = \frac{m(x + \lambda)^2}{2\hbar t} - \frac{mx^2}{2\hbar t} = \frac{mx\lambda}{\hbar t} + \frac{m\lambda^2}{2\hbar t} \quad (7)$$

$$\lambda = \frac{2\pi\hbar}{mxt^{-1}} = \frac{h}{p} \quad (8)$$

## FINDING THE CLASSICAL ENERGY

- Fix distance, vary time



**Figure:** The plot of the real part of  $\sqrt{i}$  times our kernel with varying time.

$$2\pi = \frac{mx^2}{2\hbar t} - \frac{mx^2}{2\hbar(t+T)} = \frac{mx^2}{2\hbar t^2} \left( \frac{T}{1+T/t} \right) \quad (9)$$

$$\omega \approx \frac{m}{2\hbar} \left( \frac{x}{t} \right)^2 = E/\hbar \quad (10)$$

Schrödinger's equation Have to find energy eigenvalues


Path Integral Have only to find action

BEYOND!

---


- A Wiener process is a stochastic (random) time-continuous process.
- It is also known as Brownian Motion.
- It is important to note that it forms a basis to the path integral through the Feynman-Kac formula.
- Using python, we can simulate a Wiener process

# 1 PATH







# 10 PATHS



# 50 PATHS



# 100 PATHS



## A quick summary of the Feynman approach

- Find the Lagrangian of the system

## A quick summary of the Feynman approach

- Find the Lagrangian of the system
- Find the classical path  $x_c$

## A quick summary of the Feynman approach

- Find the Lagrangian of the system
- Find the classical path  $x_c$
- Find the Action from the Lagrangian from  $\mathcal{S}[b, a] = \int_a^b \mathcal{L} dt$

## A quick summary of the Feynman approach

- Find the Lagrangian of the system
- Find the classical path  $x_c$
- Find the Action from the Lagrangian from  $\mathcal{S}[b, a] = \int_a^b \mathcal{L} dt$
- Substitute the Action into the equation for the kernel

## A quick summary of the Feynman approach

- Find the Lagrangian of the system
- Find the classical path  $x_c$
- Find the Action from the Lagrangian from  $\mathcal{S}[b, a] = \int_a^b \mathcal{L} dt$
- Substitute the Action into the equation for the kernel
- From the kernel/propagator, find the classical momentum and energy by varying distance and time respectively



## A quick summary of the Feynman approach

- Find the Lagrangian of the system
- Find the classical path  $x_c$
- Find the Action from the Lagrangian from  $\mathcal{S}[b, a] = \int_a^b \mathcal{L} dt$
- Substitute the Action into the equation for the kernel
- From the kernel/propagator, find the classical momentum and energy by varying distance and time respectively
- Done

QUESTIONS?