

A Candidate-context outcomes (n = 47)

Moderate = 0

Strong • 1 Weak / limited • 3



One unit = one displayed gene × broad-network context; repeated genes can contribute more than one unit.

B Direct candidate-level findings

APOE Astrocytes **STRONG GENE-LEVEL SUPPORT**

rs429358 • AD fine-map inclusion 1.0 • $P \approx 1.88 \times 10^{-155}$

Fallback brain evidence; not an exact astrocyte colocalization

COX7C Astrocytes + inhibitory neurons **WEAK / LIMITED**

rs2010322 • bulk sQTL CL5 • AD $P \approx 2.64 \times 10^{-6}$

One source result shown in two network contexts; not two replications

SELENOW Excitatory neurons **WEAK / LIMITED**

Reported in the source TWAS gene list

Model statistic and exact excitatory context unavailable in the public table

C Unresolved candidates

NO DIRECT MAPPING IN THE REGISTERED FILTERED SOURCE
16 nuclear genes • 23 contexts

ANKRD11	ATP6V1F	COX4I1	COX6B1
DYNLT1	FTL	LAMTOR5	LAPTM4A
NCOA1	RPL11	RPL15	RPL38
RPLP1	RPS13	RPS15	UQCR10

NOT ASSESSABLE WITH THE AVAILABLE SOURCE
6 mtDNA genes • 20 contexts

NA · MT-ATP6	NA · MT-CO2	NA · MT-CO3
NA · MT-CYB	NA · MT-ND4	NA · MT-ND5

Requires mtDNA-specific heteroplasmy, haplogroup, copy-number + NUMT controls.

No direct map ≠ no genetic role • Not assessable ≠ negative
Evidence categories summarize source coverage and direct mapping; they are not causal probabilities.