

Cultivar-dependent transcriptional responses of lychee to *Peronophythora litchii*: a registered genome-wide analysis

This single supplemental PDF contains Figures S1 to S3 with captions. Complete machine-readable Tables S1 to S18, figure source data, the registered protocol, and analysis code are openly archived at:

<https://doi.org/10.5281/zenodo.22436625>

Figure S1. Replicate-level normalized counts for the two cross-context-supported genes, twelve Tier B genes, and two legacy highlights. Points are individual deposited libraries; bars show cultivar-treatment medians.

Figure S2. Parametric detection probability for cultivar-by-infection effects under genome-wide discovery and candidate-family external adjustment. Curves show the overall result and mean-expression quartiles; the dashed line marks 80% power.

Figure S3. Quarantined PRJNA1090613 signed-signature estimate, retained as exploratory rather than confirmatory evidence.

Figure S1





Figure S1. Replicate-level normalized counts for the two cross-context-supported genes, twelve Tier B genes, and two legacy highlights. Points are individual deposited libraries; bars show cultivar-treatment medians.

Figure S2



Dashed line: 80% power. Dotted line: registered $|\log_2FC|$ threshold. Parametric power is conditional on fitted means, dispersions, deposited-library independence, and the simulated non-null prevalence.

Figure S2. Parametric detection probability for cultivar-by-infection effects under genome-wide discovery and candidate-family external adjustment. Curves show the overall result and mean-expression quartiles; the dashed line marks 80% power.

Figure S3




Figure S3. Quarantined PRJNA1090613 signed-signature estimate, retained as exploratory rather than confirmatory evidence.