

MT4531/5731: (Advanced) Bayesian Inference

Introduction to Bayesian Computation (Monte Carlo sampling)

Nicolò Margaritella

School of Mathematics and Statistics, University of St Andrews



University
of
St Andrews

Outline

- 1 Introduction
- 2 Monte Carlo Integration

Outline

1 Introduction

2 Monte Carlo Integration

Bayesian inference - univariate case

- Data $\mathbf{x} = (x_1, \dots, x_n)'$ with likelihood $f(\mathbf{x}|\theta)$.
- Prior $p(\theta)$ for the single parameter θ .
- Posterior

$$\pi(\theta|\mathbf{x}) = \frac{f(\mathbf{x}|\theta)p(\theta)}{f(\mathbf{x})}$$

where $f(\mathbf{x}) = \int f(\mathbf{x}|\theta)p(\theta)d\theta$

Bayesian inference - multivariate case (1)

- Data $\mathbf{x} = (x_1, \dots, x_n)'$ with likelihood $f(\mathbf{x}|\boldsymbol{\theta})$.
- Prior $p(\boldsymbol{\theta})$, now for a vector $\boldsymbol{\theta}$ of parameters.
- Posterior

$$\pi(\boldsymbol{\theta}|\mathbf{x}) = \frac{f(\mathbf{x}|\boldsymbol{\theta})p(\boldsymbol{\theta})}{f(\mathbf{x})}$$

where $f(\mathbf{x}) = \int \dots \int f(\mathbf{x}|\boldsymbol{\theta})p(\boldsymbol{\theta})d\boldsymbol{\theta}$

Bayesian inference - multivariate case (2)

- Often the parameters $\boldsymbol{\theta} = \{\theta_1, \theta_2, \theta_3, \dots\}$ are assumed to be independent of each other, *a priori*, so that,

$$p(\boldsymbol{\theta}) = \prod_{i=1}^n p(\theta_i).$$

- It is also common to introduce prior dependence between the parameters, by introducing another hyper-parameter ϕ , so that,

$$p(\boldsymbol{\theta}|\phi) = \prod_{i=1}^n p(\theta_i|\phi), \quad p(\phi) = \dots$$

- In more than one dimensions the posterior distribution significantly increases in complexity.

Intractable posteriors

- Data: x_1, \dots, x_n with $x_i \sim N(\mu, \sigma^2)$. μ and σ^2 unknown.
- Priors: $\mu \sim N(\phi, \tau^2)$ and $\sigma^2 \sim \Gamma^{-1}(\alpha, \beta)$
-



$$\pi(\mu, \sigma^2 | \mathbf{x}) \propto (\sigma^2)^{-(n/2 + \alpha + 1)} \exp\left(-\frac{\sum_{i=1}^n (x_i - \mu)^2}{2\sigma^2}\right) \\ \exp\left(-\frac{\beta}{\sigma^2}\right) \exp\left(-\frac{(\mu - \phi)^2}{2\tau^2}\right)$$

-

$$f(\mathbf{x}) = \int_{\mu=-\infty}^{\infty} \int_{\sigma^2=0}^{\infty} (\sigma^2)^{-(n/2 + \alpha + 1)} \exp\left(-\frac{\sum_{i=1}^n (x_i - \mu)^2}{2\sigma^2}\right) \\ \exp\left(-\frac{\beta}{\sigma^2}\right) \exp\left(-\frac{(\mu - \phi)^2}{2\tau^2}\right) d\mu d\sigma^2$$



Bayesian computation

- Q: How do we do Bayesian analysis when we have posterior distributions of non-standard form, possibly of high dimension?
- A: Use a computer to generate random samples from the posterior.
- Use these samples to provide empirical estimates of the quantities we want





Bayesian computation

- Q: How do we do Bayesian analysis when we have posterior distributions of non-standard form, possibly of high dimension?
- A: Use a computer to generate random samples from the posterior.
- Use these samples to provide empirical estimates of the quantities we want



Bayesian computation


- Q: How do we do Bayesian analysis when we have posterior distributions of non-standard form, possibly of high dimension?
- A: Use a computer to generate random samples from the posterior.
- Use these samples to provide empirical estimates of the quantities we want




Samples can be independent or dependent

- Two ways to generate the samples: independent or dependent (i.e., value of one sample depends on the value of the previous one)
- Both give the same distribution in the end, but with dependent samples:
 - Start point matters for a while – i.e., it takes a while to “burn-in”
 - You need more samples for the same level of accuracy


Independent samples, 1000 points



Dependent samples, 1000 points




Dependent samples, 3000 points, first 50 removed




Samples can be independent or dependent

- Two ways to generate the samples: independent or dependent (i.e., value of one sample depends on the value of the previous one)
- Both give the same distribution in the end, but with dependent samples:
 - Start point matters for a while – i.e., it takes a while to “burn-in”
 - You need more samples for the same level of accuracy


Independent samples, 1000 points



Dependent samples, 1000 points




Dependent samples, 3000 points, first 50 removed




Samples can be independent or dependent

- Two ways to generate the samples: independent or dependent (i.e., value of one sample depends on the value of the previous one)
- Both give the same distribution in the end, but with dependent samples:
 - Start point matters for a while – i.e., it takes a while to “burn-in”
 - You need more samples for the same level of accuracy


Independent samples, 1000 points



Dependent samples, 1000 points



Dependent samples, 3000 points, first 50 removed



Outline

1 Introduction

2 Monte Carlo Integration

Monte Carlo Integration

- Example: Estimate the posterior (marginal) expectation of θ given \mathbf{x}

$$\mathbb{E}_{\pi}(\theta) = \int \theta \pi(\theta|\mathbf{x}) d\theta$$

- Say we have a sample $\theta^1, \dots, \theta^n$ from the $\pi(\theta|\mathbf{x})$
- We can estimate the expectation using

$$\hat{\mathbb{E}}_{\pi}(\theta) = \frac{1}{n} \sum_{i=1}^n \theta^i$$

- This idea extends to any function of θ , $f(\theta)$. For example, the posterior mean of $f(\theta)$:

$$\hat{\mathbb{E}}_{\pi}(f(\theta)) = \frac{1}{n} \sum_{i=1}^n f(\theta^i)$$

Monte Carlo Integration

- For 100 samples θ^i , the 5% quantile of $\pi(\theta|\mathbf{x})$, is obtained by considering the fifth sample $\theta^{(5)}$, after sorting the sample to $\theta^{(1)}, \theta^{(2)}, \theta^{(3)}, \dots$ in increasing order:

$$P_{\pi}(\theta \leq \theta^{(5)}|\mathbf{x}) \simeq 0.05.$$

- The posterior variance of $f(\theta)$?
- The probability $f(\theta)$ is more than some value c ?

See Exercise 2 in tutorial 4.

Example

- Suppose $\pi(\theta|\mathbf{x}) \sim N(0, 1)$.
- Hence $\mathbb{E}(\theta) = 0$ and $\text{sd}(\theta) = 1$.
- Simulate values from `rnorm` in R:

Samples n	1		2		3	
	Sample Mean	Sample SD	S. Mean	S. SD	S. Mean	S. SD
10	-0.39	1.26	-0.36	1.32	-0.09	0.87
100	0.36	0.88	-0.23	0.94	0.10	0.95
1000	-0.10	1.01	-0.01	1.01	0.02	0.98
10000	-0.01	1.00	0.00	0.99	0.00	1.00
100000	0.00	1.00	0.00	1.00	0.00	1.00

Monte Carlo Error

- Measures the between-simulation variability in estimates
- For n independent θ^i samples, for estimating the expected value of θ :

$$\bar{\theta} \sim N\left(\mathbb{E}_{\pi}(\theta), \frac{\sigma^2}{n}\right).$$

- σ^2/n is the Monte Carlo variance
- $\sqrt{\sigma^2/n}$ is the Monte Carlo error.
- See later in module for dependent samples...

Aside: Calculating any definite integral using MC integration

- MC integration can in fact be used to calculate any definite integral

$$\int_a^b f(x) dx$$

- you can introduce the Uniform $U(a, b)$ within the integral

$$(b - a) \int_a^b f(x) \frac{1}{b - a} dx$$

- so that the integral is calculated by approximating the expectation of $f(x)$ multiplied by $(b - a)$.

$$\int_a^b f(x) dx \simeq (b - a) \hat{\mathbb{E}}(f(x)) = \frac{(b - a)}{n} \sum_{i=1}^n f(x^i)$$

- where the x^i are sampled from the uniform $U(a, b)$.

Conclusion

- Bayesian problems frequently lead to intractable integrals
- If we can sample from the posterior, we can use Monte Carlo integration to obtain estimates of quantities of interest
- So, we have replaced an integration problem with a sampling problem
- How do we obtain samples from the posterior?
- ... see future lectures.