

Class14: RNA-Seq Mini Project

Yu (Ericsson) Cao (PID: A16421048)

Run a complete RNASeq analysis workflow from counts to enriched genesets...

Data import

```
counts <- read.csv("GSE37704_featurecounts.csv", row.names=1)
metadata <- read.csv("GSE37704_metadata.csv")
```

Data exploration

```
head(counts, 3)
```

	length	SRR493366	SRR493367	SRR493368	SRR493369	SRR493370
ENSG00000186092	918	0	0	0	0	0
ENSG00000279928	718	0	0	0	0	0
ENSG00000279457	1982	23	28	29	29	28
	SRR493371					
ENSG00000186092	0					
ENSG00000279928	0					
ENSG00000279457	46					

```
head(metadata)
```

	id	condition
1	SRR493366	control_sirna
2	SRR493367	control_sirna
3	SRR493368	control_sirna
4	SRR493369	hoxa1_kd

```
5 SRR493370      hoxa1_kd
6 SRR493371      hoxa1_kd
```

Check if the colData and countData match up

```
metadata$id
```

```
[1] "SRR493366" "SRR493367" "SRR493368" "SRR493369" "SRR493370" "SRR493371"
```

```
colnames(counts)
```

```
[1] "length"      "SRR493366" "SRR493367" "SRR493368" "SRR493369" "SRR493370"
[7] "SRR493371"
```

```
#RM first column
countData <- counts[,-1]
```

Now check if the metadata and count data columns match

```
all(colnames(countData) == metadata$id)
```

```
[1] TRUE
```

We need to remove all the zero count genes.

```
head(countData)
```

	SRR493366	SRR493367	SRR493368	SRR493369	SRR493370	SRR493371
ENSG00000186092	0	0	0	0	0	0
ENSG00000279928	0	0	0	0	0	0
ENSG00000279457	23	28	29	29	28	46
ENSG00000278566	0	0	0	0	0	0
ENSG00000273547	0	0	0	0	0	0
ENSG00000187634	124	123	205	207	212	258

To identify these zero count genes we can sum across the rows and check if the sum is more than zero

```
to.keep.inds <- rowSums(countData) > 0
nonzero.counts <- countData[to.keep.inds,]
```

DESeq setup and analysis

```
library(DESeq2)

dds <- DESeqDataSetFromMatrix(countData = nonzero.counts,
                              colData = metadata,
                              design = ~condition)
```

Warning in DESeqDataSet(se, design = design, ignoreRank): some variables in design formula are characters, converting to factors

```
dds <- DESeq(dds)
```

estimating size factors

estimating dispersions

gene-wise dispersion estimates

mean-dispersion relationship

final dispersion estimates

fitting model and testing

Result extraction

```
res <- results(dds)
head(res)
```

log2 fold change (MLE): condition hoxa1 kd vs control sirna

Wald test p-value: condition hoxa1 kd vs control sirna

DataFrame with 6 rows and 6 columns

	baseMean	log2FoldChange	lfcSE	stat	pvalue
	<numeric>	<numeric>	<numeric>	<numeric>	<numeric>
ENSG00000279457	29.9136	0.1792571	0.3248216	0.551863	5.81042e-01
ENSG00000187634	183.2296	0.4264571	0.1402658	3.040350	2.36304e-03
ENSG00000188976	1651.1881	-0.6927205	0.0548465	-12.630158	1.43989e-36
ENSG00000187961	209.6379	0.7297556	0.1318599	5.534326	3.12428e-08
ENSG00000187583	47.2551	0.0405765	0.2718928	0.149237	8.81366e-01
ENSG00000187642	11.9798	0.5428105	0.5215599	1.040744	2.97994e-01
	padj				
	<numeric>				
ENSG00000279457	6.86555e-01				
ENSG00000187634	5.15718e-03				
ENSG00000188976	1.76549e-35				
ENSG00000187961	1.13413e-07				
ENSG00000187583	9.19031e-01				
ENSG00000187642	4.03379e-01				

```
summary(res)
```

out of 15975 with nonzero total read count

adjusted p-value < 0.1

LFC > 0 (up) : 4349, 27%

LFC < 0 (down) : 4396, 28%

outliers [1] : 0, 0%

low counts [2] : 1237, 7.7%


(mean count < 0)

[1] see 'cooksCutoff' argument of ?results

[2] see 'independentFiltering' argument of ?results

Volcano Plot

```
plot(res$log2FoldChange, -log(res$padj))
```




Setup a wee color vector for this plot

```
mycols <- rep("gray", nrow(res) )
```

```
mycols[ abs(res$log2FoldChange) > 2 ] <- "red"
```

```
inds <- (res$padj < 0.01) & (abs(res$log2FoldChange) > 2 )
mycols[ inds ] <- "blue"
```

```
plot( res$log2FoldChange, -log(res$padj),col=mycols, xlab="Log2(FoldChange)", ylab="-Log(P
```



Gene annotation

```
library("AnnotationDbi")
library("org.Hs.eg.db")
```

```
columns(org.Hs.eg.db)
```

```
[1] "ACCNUM"      "ALIAS"       "ENSEMBL"     "ENSEMBLPROT" "ENSEMBLTRANS"
[6] "ENTREZID"    "ENZYME"      "EVIDENCE"    "EVIDENCEALL"  "GENENAME"
[11] "GENETYPE"    "GO"          "GOALL"       "IPI"          "MAP"
[16] "OMIM"        "ONTOLOGY"    "ONTOLOGYALL" "PATH"         "PFAM"
[21] "PMID"        "PROSITE"     "REFSEQ"      "SYMBOL"       "UCSCKG"
[26] "UNIPROT"
```

```
res$symbol <- mapIds(x=org.Hs.eg.db,
                     keys=rownames(nonzero.counts),
                     keytype="ENSEMBL",
```

```
column="SYMBOL")
```

'select()' returned 1:many mapping between keys and columns

```
res$entrez <- mapIds(x=org.Hs.eg.db,
                    keys=rownames(nonzero.counts),
                    keytype="ENSEMBL",
                    column="ENTREZID")
```

'select()' returned 1:many mapping between keys and columns

```
res$name <- mapIds(x=org.Hs.eg.db,
                  keys=rownames(nonzero.counts),
                  keytype="ENSEMBL",
                  column="GENENAME")
```

'select()' returned 1:many mapping between keys and columns

```
head(res)
```

log2 fold change (MLE): condition hoxa1 kd vs control sirna

Wald test p-value: condition hoxa1 kd vs control sirna

DataFrame with 6 rows and 9 columns

	baseMean	log2FoldChange	lfcSE	stat	pvalue
	<numeric>	<numeric>	<numeric>	<numeric>	<numeric>
ENSG00000279457	29.9136	0.1792571	0.3248216	0.551863	5.81042e-01
ENSG00000187634	183.2296	0.4264571	0.1402658	3.040350	2.36304e-03
ENSG00000188976	1651.1881	-0.6927205	0.0548465	-12.630158	1.43989e-36
ENSG00000187961	209.6379	0.7297556	0.1318599	5.534326	3.12428e-08
ENSG00000187583	47.2551	0.0405765	0.2718928	0.149237	8.81366e-01
ENSG00000187642	11.9798	0.5428105	0.5215599	1.040744	2.97994e-01
	padj	symbol	entrez	name	
	<numeric>	<character>	<character>	<character>	
ENSG00000279457	6.86555e-01	NA	NA	NA	
ENSG00000187634	5.15718e-03	SAMD11	148398	sterile alpha motif ..	
ENSG00000188976	1.76549e-35	NOC2L	26155	NOC2 like nucleolar ..	
ENSG00000187961	1.13413e-07	KLHL17	339451	kelch like family me..	
ENSG00000187583	9.19031e-01	PLEKHN1	84069	pleckstrin homology ..	
ENSG00000187642	4.03379e-01	PERM1	84808	PPARGC1 and ESRR ind..	

Pathway analysis

```
library(gage)
library(gageData)
library(pathview)
```

```
data(kegg.sets.hs)
data(sigmet.idx.hs)
```

```
kegg.sets.hs = kegg.sets.hs[sigmet.idx.hs]
```

```
head(kegg.sets.hs, 3)
```

```
$`hsa00232 Caffeine metabolism`
```

```
[1] "10" "1544" "1548" "1549" "1553" "7498" "9"
```

```
$`hsa00983 Drug metabolism - other enzymes`
```

```
[1] "10" "1066" "10720" "10941" "151531" "1548" "1549" "1551"
[9] "1553" "1576" "1577" "1806" "1807" "1890" "221223" "2990"
[17] "3251" "3614" "3615" "3704" "51733" "54490" "54575" "54576"
[25] "54577" "54578" "54579" "54600" "54657" "54658" "54659" "54963"
[33] "574537" "64816" "7083" "7084" "7172" "7363" "7364" "7365"
[41] "7366" "7367" "7371" "7372" "7378" "7498" "79799" "83549"
[49] "8824" "8833" "9" "978"
```

```
$`hsa00230 Purine metabolism`
```

```
[1] "100" "10201" "10606" "10621" "10622" "10623" "107" "10714"
[9] "108" "10846" "109" "111" "11128" "11164" "112" "113"
[17] "114" "115" "122481" "122622" "124583" "132" "158" "159"
[25] "1633" "171568" "1716" "196883" "203" "204" "205" "221823"
[33] "2272" "22978" "23649" "246721" "25885" "2618" "26289" "270"
[41] "271" "27115" "272" "2766" "2977" "2982" "2983" "2984"
[49] "2986" "2987" "29922" "3000" "30833" "30834" "318" "3251"
[57] "353" "3614" "3615" "3704" "377841" "471" "4830" "4831"
[65] "4832" "4833" "4860" "4881" "4882" "4907" "50484" "50940"
[73] "51082" "51251" "51292" "5136" "5137" "5138" "5139" "5140"
[81] "5141" "5142" "5143" "5144" "5145" "5146" "5147" "5148"
[89] "5149" "5150" "5151" "5152" "5153" "5158" "5167" "5169"
[97] "51728" "5198" "5236" "5313" "5315" "53343" "54107" "5422"
[105] "5424" "5425" "5426" "5427" "5430" "5431" "5432" "5433"
[113] "5434" "5435" "5436" "5437" "5438" "5439" "5440" "5441"
```



```
[121] "5471"    "548644" "55276"  "5557"   "5558"   "55703"  "55811"  "55821"
[129] "5631"    "5634"   "56655"  "56953"  "56985"  "57804"  "58497"  "6240"
[137] "6241"    "64425"  "646625" "654364" "661"    "7498"   "8382"   "84172"
[145] "84265"   "84284"  "84618"  "8622"   "8654"   "87178"  "8833"   "9060"
[153] "9061"    "93034"  "953"    "9533"   "954"    "955"    "956"    "957"
[161] "9583"    "9615"
```

```
foldchanges = res$log2FoldChange
names(foldchanges) = res$entrez
head(foldchanges)
```

```
<NA>      148398      26155      339451      84069      84808
0.17925708 0.42645712 -0.69272046 0.72975561 0.04057653 0.54281049
```

```
# Get the results
keggres = gage(foldchanges, gsets=kegg.sets.hs)
attributes(keggres)
```

```
$names
[1] "greater" "less"    "stats"
```

Now, let's look at one of these pathways: hsa04110

```
pathview(gene.data=foldchanges, pathway.id="hsa04110")
```

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/yucao515/Desktop/BIMM 143/Class 14

Info: Writing image file hsa04110.pathview.png

```
# A different PDF based output of the same data
pathview(gene.data=foldchanges, pathway.id="hsa04110", kegg.native=FALSE)
```

'select()' returned 1:1 mapping between keys and columns

Warning: reconcile groups sharing member nodes!




Figure 1: The KEGG Cell Cycle Pathway

```
      [,1] [,2]
[1,] "9"  "300"
[2,] "9"  "306"
```

Info: Working in directory /Users/yuca515/Desktop/BIMM 143/Class 14

Info: Writing image file hsa04110.pathview.pdf

```
## Focus on top 5 upregulated pathways here for demo purposes only
keggrespathways <- rownames(keggres$greater)[1:5]
```

```
# Extract the 8 character long IDs part of each string
keggresids = substr(keggrespathways, start=1, stop=8)
keggresids
```

```
[1] "hsa04640" "hsa04630" "hsa00140" "hsa04142" "hsa04330"
```

```
pathview(gene.data=foldchanges, pathway.id=keggresids, species="hsa")
```

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/yuca515/Desktop/BIMM 143/Class 14

Info: Writing image file hsa04640.pathview.png

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/yuca515/Desktop/BIMM 143/Class 14

Info: Writing image file hsa04630.pathview.png

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/yuca515/Desktop/BIMM 143/Class 14

Info: Writing image file hsa00140.pathview.png

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/yucao515/Desktop/BIMM 143/Class 14


Info: Writing image file hsa04142.pathview.png


Info: some node width is different from others, and hence adjusted!

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/yucao515/Desktop/BIMM 143/Class 14

Info: Writing image file hsa04330.pathview.png





Gene Ontology

```
data(go.sets.hs)
data(go.subs.hs)

# Focus on Biological Process subset of GO
gobpsets <- go.sets.hs[go.subs.hs$BP]

gobpres <- gage(foldchanges, gsets=gobpsets, same.dir=TRUE)

head(gobpres$less)
```

	p.geomean	stat.mean	p.val
GO:0048285 organelle fission	1.536227e-15	-8.063910	1.536227e-15
GO:0000280 nuclear division	4.286961e-15	-7.939217	4.286961e-15
GO:0007067 mitosis	4.286961e-15	-7.939217	4.286961e-15
GO:0000087 M phase of mitotic cell cycle	1.169934e-14	-7.797496	1.169934e-14
GO:0007059 chromosome segregation	2.028624e-11	-6.878340	2.028624e-11
GO:0000236 mitotic prometaphase	1.729553e-10	-6.695966	1.729553e-10

	q.val	set.size	expl
GO:0048285 organelle fission	5.843127e-12	376	1.536227e-15
GO:0000280 nuclear division	5.843127e-12	352	4.286961e-15
GO:0007067 mitosis	5.843127e-12	352	4.286961e-15
GO:0000087 M phase of mitotic cell cycle	1.195965e-11	362	1.169934e-14
GO:0007059 chromosome segregation	1.659009e-08	142	2.028624e-11
GO:0000236 mitotic prometaphase	1.178690e-07	84	1.729553e-10

Reactome Analysis

We can use reactome via an R package or use their relatively new website interface. Let's use the latter.

It wants a list of our most interesting (i.e. significant) genes in gene SYMBOL format.

```
sig_genes <- res[res$padj <= 0.05 & !is.na(res$padj), "symbol"]
print(paste("Total number of significant genes:", length(sig_genes)))
```

```
[1] "Total number of significant genes: 8147"
```

```
#sig_genes
```

We will write these out to a text file so we can use them on the website:

```
write.table(sig_genes, file="significant_genes.txt", row.names=FALSE, col.names=FALSE, quote=FALSE)
```




Figure 2: Example figure from reactome online - mitosis

Q: What pathway has the most significant “Entities p-value”? Do the most significant pathways listed match your previous KEGG results? What factors could cause differences between the two methods?

A: Cell Cycle has the most significant entities P-value. Yes, the most significant pathways listed do match my previous KEGG results. They analyze pathways from different perspectives.