

Standardization

Understanding Confounding and Population Structure

Erik Westlund

2025-10-20



Motivating Example

Delayed Seeking of Necessary Healthcare

People wait to get help until a health issue becomes serious or avoidable harm has already occurred

Why This Matters

- Skipping routine visits or screenings until symptoms worsen
- Delaying care due to cost, time, or caregiving demands
- Arriving at the health system only during a crisis

Reveals inequities in access, prevention, and confidence in the health system

Our Question

Who faces greater risk (e.g., men or women?) and how can we compare that risk fairly?

The Challenge

Crude rates can mislead us about *who* is at risk and *why*

Gender

- **Gender-based analysis** asks why – how roles, norms, and power shape health and behavior
- Those processes interact with age and the gender equality context

This lecture's focus: How causal structure and population composition determine what crude rates actually mean

Agenda

1. **The core idea:** Statistical expectation and population structure
2. **Example 1:** Age $\rightarrow$ Outcome (composition matters)
3. **Example 2:** Age + Gender $\rightarrow$ Outcome (Simpson's paradox)
4. **Example 3:** Age + Gender + Gender Opportunity $\rightarrow$ Outcome (limits of manual standardization)
5. **Summary:** Getting the expected value right

The Core Problem

- Every population has its own **mix of people** – by age, gender, opportunity, and access.
- That mix shapes what we see in the data: who gets sick, who shows up, and when.
- When we compare groups without accounting for those differences, we risk blaming behavior or biology for what is really structure.

Statistical Expectation

- The **expected rate** is what we'd see if we looked across everyone in a group, *given their mix of ages and circumstances*.
- If two groups have different mixes, their expected rates aren't directly comparable.
- We get the **wrong picture** because we're averaging over different kinds of people.

How Standardization Fixes It

- **Step 1:** Agree on a common mix – the same age, gender, or opportunity structure for everyone.
- **Step 2:** Recalculate each group's rate *as if they had that same mix*.
- **Result:** Differences now reflect real risk, not who happens to be older, poorer, or more constrained.

In short: Standardization answers, “What would these rates look like if all groups lived under the same conditions?”

Example 1: Age →
Outcome

The Simplest Case: Age Only

Research question: What's the delayed healthcare seeking rate in our population?

Our Simulated Data

- Registry data on preventable health crises
- E.g., diabetic emergencies, hypertensive crises, late-stage cancers
- **We define the risk by age: Young: 5%, Middle: 15%, Older: 40%**

The Question

What crude rate do we expect?

Key insight: Depends entirely on population age structure!

Causal Structure: Age → Outcome

Causal Structure: Age Determines Outcome




No confounding here. Just one cause: Age
But crude rate still depends on age.

Simulated Dataset 1: Age Distribution

Population Age Structure (Dataset 1)


Known From Simulation $P(\text{Age}) = (30\%, 35\%, 35\%)$



Age-Specific Rates Of Late Presentation

Late Presentation Age-Specific Rates

We set these from simulated data, so we know the truth/unbiased rates



Computing the Expected Value

$$E[Y] = \sum_{\text{age}} E[Y \mid \text{Age}] \times P(\text{Age})$$

```
# Our age-specific rates (known truth)
rates <- c(Young = 0.05, Middle = 0.15, Older = 0.40)

# Our population weights
weights <- c(Young = 0.30, Middle = 0.35, Older = 0.35)


# Expected value
expected_rate <- sum(rates * weights)
expected_rate

[1] 0.2075
```

Expected crude rate: 20.7%

This is what we should see in the data.

Observed vs Expected



Because we don't have confounding here, our actual rate matches our expected rate.

What If We Had a Different Population?

- Imagine Population A: Mostly young people → low crude rate
- Imagine Population B: Mostly older people → high crude rate
- **Same age-specific rates (5%, 15%, 40%)**
- **Different crude rates** because populations differ

Why Standardization Matters

To compare populations fairly, we need:

- Apply age-specific rates to a **standard population**
- Use the **same age weights** for both groups
- Remove the effect of different population structures

Example 2: Age + Gender

Adding Gender: Simpson's Paradox

Research question: Do men or women have higher late-presentation rates?


What We Know

- Now we have Gender and Age in our data
- **Known truth** (from simulation):
 - Men have higher risk within **every** age group
 - Women live longer → more older women

Question: What will crude rates show?

Causal Structure: Age + Gender → Outcome

Causal Structure: Gender → Age → Outcome, Gender → Outcome



Confounding By Age

This is confounding!


Gender affects Age (women live longer), and Age affects Outcome.

Crude comparison of men vs women is **biased** by age composition.


Simulated Dataset 2: Age Structure by Gender

Age Distribution by Gender (Dataset 2)

Women concentrated in older ages; men in younger ages



Crude Rates by Gender



Observation: Women have higher crude rate
But is this the truth?

Age-Specific Rates by Gender

```
age_gender_rates <- sim2_data |>
  group_by(gender, age_group) |>
  summarise(rate = mean(late_presentation), .groups = "drop")
```

age_gender_rates

```
# A tibble: 6 × 3
  gender age_group rate
<chr> <chr> <dbl>
1 Men Middle 0.162
2 Men Older 0.462
3 Men Young 0.0633
4 Women Middle 0.153
5 Women Older 0.399
6 Women Young 0.0508
```


Within every age group, men have higher rates!

This is Simpson's Paradox.

Visualizing Simpson's Paradox

Within Each Age Group, Men Have Higher Rates

Simpson's Paradox: Composition flips the overall pattern



What Went Wrong?

Remember the expectation formula:

$$E[Y \mid \text{Gender}] = \sum_{\text{age}} E[Y \mid \text{Gender}, \text{Age}] \times P(\text{Age} \mid \text{Gender})$$

The Problem

- $P(\text{Age} \mid \text{Man}) \neq P(\text{Age} \mid \text{Woman})$
- Women concentrated in **older ages** (high risk)
- Men concentrated in **younger ages** (low risk)

Why It Flips

- Men have higher risk within **every** age group
- But **different age weights** flip the crude pattern
- This is confounding in action

Indirect Standardization: The Goal

Estimate what each group's outcomes *should look like* if they experienced the same rates as a reference population

The Method

- Start with **known rates** from a large or “standard” population (for example, national age-specific rates)
- Apply those rates to your **study population’s structure** (how many people you have in each age group)
- This gives you the **expected number of events** *if your group had the same risks as the reference group*

Interpreting the Results

- Compare observed vs expected events → this ratio is the **Standardized Mortality (or Morbidity) Ratio (SMR)**
- $SMR = 1$ → outcomes match the reference population
- $SMR > 1$ → more events than expected
- $SMR < 1$ → fewer events than expected

Why “Indirect”?

We don't re-calculate each group's own rates

- We *borrow* the standard rates and apply them to our data
- Then compare what we **saw** to what we **would expect**

Step 1: Define Our “Ingredients”

```
# Calculate observed rates in our data
age_gender_rates <- sim2_data |>
  group_by(gender, age_group) |>
  summarise(rate = mean(late_presentation), .groups = "drop")

# Define standard rates (e.g., from national statistics)
standard_rates <- data.frame(
  age_group = factor(c("Young", "Middle", "Older"),
    levels = c("Young", "Middle", "Older")),
  std_rate = c(0.04, 0.14, 0.42) # Externally-determined rates; they differ a
)

# Get population counts by gender and age
pop_counts <- sim2_data |>
  group_by(gender, age_group) |>
  summarise(n = n(), .groups = "drop")
```

Step 2: Calculate SMR for Each Group


```
# For each gender, calculate observed and expected events
indirect_std <- pop_counts |>
  left_join(standard_rates, by = "age_group") |> # Merge in our rates
  left_join(age_gender_rates, by = c("gender", "age_group")) |>
  group_by(gender) |>
  summarise(
    observed = sum(n * rate),      # Actual events in our data
    expected = sum(n * std_rate),  # Expected if standard rates applied
    SMR = observed / expected
  )
```

The Results

```
indirect_std
```

```
# A tibble: 2 × 4
  gender observed expected SMR
<chr>      <dbl>      <dbl> <dbl>
1 Men      2085.      1805.  1.16
2 Women    2338      2355.  0.993
```


SMR: Standardized Morbidity Ratio



Interpretation: Men have 16% higher risk than expected from standard rates

Direct Standardization: The Goal

What would rates be if men and women had the same age distribution?

```
# Calculate observed rates in our data
age_gender_rates <- sim2_data |>
  group_by(gender, age_group) |>
  summarise(rate = mean(late_presentation), .groups = "drop")

# Use marginal P(Age) as standard weights
std_weights <- sim2_data |>
  count(age_group) |>
  mutate(weight = n / sum(n))
```

Calculate Standardized Rates

```
# Apply standard weights to gender-age specific rates
direct_std <- age_gender_rates |>
  left_join(std_weights, by = "age_group") |>
  group_by(gender) |>
  summarise(standardized_rate = sum(rate * weight))

direct_std

# A tibble: 2 × 2
  gender standardized_rate
<chr>      <dbl>
1 Men      0.238
2 Women    0.208
```

Crude vs Standardized Rates

Crude vs Direct Standardized Rates
Standardization reveals men have higher rates



Pattern reverses! Standardization reveals the truth: men have higher risk.

Key Insight: Confounding

- Crude rates get it wrong because Gender $\rightarrow$ Age and Age $\rightarrow$ Outcome
- Age is a confounder of the Gender-Outcome relationship
- Direct standardization fixes $P(\text{Age})$ to break the confounding

The Result

Direct standardization gives us the **correct comparison of gender differences**

Next: What happens when we have multiple confounders?

Example 3: Age + Gender + GOC

Adding Gender Opportunity Context

Research question: Same question, but now we also have GOC data

GOC = Gender Opportunity Context (gendered norms/power/access)

Known Truth from Simulation

- Age effect: Young=5%, Middle=15%, Older=40%
- Gender effect: Men higher within age groups
- GOC effect: Low/Mid higher than High

Multiple Confounding Pathways


- GOC → Age (low GOC = younger; more constraints/shorter healthy lifespan)
- GOC → Outcome (low GOC = higher risk; barriers to access/norms)
- Gender → Age (women live longer)
- Age → Outcome
- Gender → Outcome

The Complexity

All three variables affect both each other AND the outcome

Causal Structure: Multiple Confounders

Gender Opportunity Context introduces multiple pathways



This is complex!


To compare groups fairly, we need to adjust for Age AND GOC.

Manual standardization becomes unmanageable.

Simulated Dataset 3: Age by Gender and GOC

Age Distribution by Gender and GOC (Dataset 3)

Complex patterns: $P(\text{Age} \mid \text{Gender}, \text{GOC})$ varies, but marginal (i.e., overall) $P(\text{Age})$ same as D1/D2



Crude Rates by Gender

```
# By Gender
sim3_data |>
  group_by(gender) |>
  summarise(crude_rate = mean(late_presentation))
```

```
# A tibble: 2 × 2
  gender crude_rate
<chr>      <dbl>
1 Men      0.244
2 Women    0.258
```

Crude Rates by GOC

```
# By GOC
sim3_data |>
  group_by(goc) |>
  summarise(crude_rate = mean(late_presentation))
```

```
# A tibble: 3 × 2
  goc   crude_rate
<chr>      <dbl>
1 High      0.248
2 Low       0.247
3 Mid       0.258
```

All confounded! Low GOC appears to have every so slightly lower risk (wrong!) because low GOC contexts are younger.

Why Manual Standardization Fails

To standardize by Age and GOC:

- Need age-GOC-specific rates for each gender: $3 \text{ ages} \times 3 \text{ GOC} = 9 \text{ strata per gender}$
- With small cell counts, rates become unstable
- Adding more confounders (ethnicity, region, comorbidity)
→ exponential growth

The Solution

Regression models do multivariable standardization automatically

Poisson Regression

```
glm_fit <- glm(late_presentation ~ gender + age_group + goc,  
              family = poisson(link = "log"),  
              data = sim3_data)
```

What the Model Does

Fits a model:

$$\log(E[Y \mid \text{Gender, Age, GOC}]) = \beta_0 + \beta_{\text{Gender}} + \beta_{\text{Age}} + \beta_{\text{GOC}}$$

Why This Works

- Automatically adjusts for all confounders simultaneously
- **This is standardization!** Model weights by the data's covariate distribution

Adjusted Probabilities: Men vs Women

```
library(emmeans)
```

```
# Get marginal probabilities by gender (adjusted for age and GOC)  
emmeans(glm_fit, ~ gender, type = "response")
```

gender	rate	SE	df	asymp.LCL	asymp.UCL
Men	0.194	0.00466	Inf	0.185	0.204
Women	0.167	0.00421	Inf	0.159	0.175

Results are averaged over the levels of: age_group, goc
Confidence level used: 0.95
Intervals are back-transformed from the log scale

Gender Probability Differences

```
# Probability difference  
pairs(regrid(emmeans(glm_fit, ~ gender, type = "response"))), reverse = TRUE)
```

contrast	estimate	SE	df	z.ratio	p.value
Women - Men	-0.0275	0.00514	Inf	-5.342	<.0001

Results are averaged over the levels of: age_group, goc

Men have 3.2 percentage points higher risk than women
(after adjusting for age and GOC)

Adjusted Probabilities by GOC

```
# Get marginal probabilities by GOC (adjusted for age and gender)
emmeans(glm_fit, ~ goc, type = "response")
```

goc	rate	SE	df	asympt.LCL	asympt.UCL
High	0.162	0.00560	Inf	0.151	0.173
Low	0.203	0.00491	Inf	0.194	0.213
Mid	0.177	0.00490	Inf	0.168	0.187

Results are averaged over the levels of: gender, age_group
Confidence level used: 0.95
Intervals are back-transformed from the log scale

GOC Probability Differences

```
# Probability differences  
pairs(regrid(emmeans(glm_fit, ~ goc, type = "response")), reverse = TRUE)
```

contrast	estimate	SE	df	z.ratio	p.value
Low - High	0.0417	0.00666	Inf	6.259	<.0001
Mid - High	0.0154	0.00655	Inf	2.349	0.0493
Mid - Low	-0.0263	0.00604	Inf	-4.350	<.0001

Results are averaged over the levels of: gender, age_group
P value adjustment: tukey method for comparing a family of 3 estimates

Interpretation

- Low GOC contexts have 4.7 pp higher risk than High GOC (adjusted for age/gender)
- Mid GOC contexts have 1.7 pp higher risk than High GOC (adjusted for age/gender)

Model-based Approaches

Modeling lets us extend the logic of standardization to more complex causal structures gracefully

Summary

Recap

1. **Example 1 (Age only):** Crude rate depends on population structure
2. **Example 2 (Age + Gender):** Confounding creates Simpson's Paradox; direct standardization fixes it
3. **Example 3 (Age + Gender + GOC):** Multiple confounders: regression models generalize standardization

All Methods Compared

Getting the Expected Value Right Requires Understanding Causal Structure

Method	Example 2 (Gender)	Example 3 (GOC + Gender)
Crude	Women appear worse (incorrect)	Low GOC appears better (incorrect)
Direct Standardization	Men worse (correct)	Too many strata (unmanageable)
Poisson GLM	Men worse (correct)	Men worse, Low GOC worse (correct)

Key Takeaways

1. **Expectation** = weighted average of subgroup rates
2. **Population structure** determines which weights we use
3. **Confounding** occurs when groups have different structures
4. **Standardization** uses common weights to make fair comparisons
5. **Regression models** = automated multivariable standardization

Connecting to Policy

If we only looked at crude rates:

- **Example 2:** Crude data: women seem worse. Truth: men higher risk
- **Example 3:** Crude data: low-opportunity contexts seem safer. Truth: risk highest where gender equality is lowest

The Consequences

- **Wrong interventions:** Misallocated resources, ineffective programs
- **Root cause:** Ignored causal structure and population composition

Gendered health analysis requires understanding confounding, causal structure, and standardization

Summary

What you just learned:

- Causal structure and confounding
- Direct/indirect standardization for 1-2 confounders
- Why manual standardization fails with many confounders

Questions?

Email me at ewestlund@jhu.edu and I'm happy to talk!

Thank You

Erik Westlund

ewestlund@jhu.edu

Johns Hopkins Biostatistics Center

Johns Hopkins Bloomberg School of Public Health

Slides and code: github.com/erikwestlund/gdaw-standardization-presentation-2025