

Notes on LQ project I

Nicolas Mileo^{1,*} and Alejandro Szynkman^{1,†}

¹*IFLP, CONICET – Dpto. de Física, Universidad
Nacional de La Plata, C.C. 67, 1900 La Plata, Argentina*

(Dated: March 18, 2016)

* mileo@fisica.unlp.edu.ar

† szynkman@fisica.unlp.edu.ar

I. STANDARD MODEL CROSS SECTION

Within the Standard Model (SM) there are two types of interaction between the incoming neutrino and the nucleons in the detector that are the most relevant in order to understand the signal events collected in IceCube, namely the charged current process (C.C. process) in which the interaction between the neutrino and the nucleon proceeds via the exchange of a W boson and the neutral current process (N.C. process) in which a Z boson is exchanged [?].

Let us first compute the cross section for the C.C. process $\nu_\ell N \rightarrow \ell^- + \text{anything}$. The relevant dominant diagrams at parton level are shown in Fig. 1 where $\ell = e, \mu, \tau$, the index i denotes the quark flavor and $\tilde{Q} = p - q$.



FIG. 1: Dominant charged current process diagrams. The left diagram shows the scattering of a neutrino by a down-type quark whereas the right diagram depicts the scattering by an up-type antiquark.

In the case of the first diagram the amplitude is given by

$$\mathcal{M}(\nu_\ell d_i \rightarrow \ell^- u_i) = \left(-\frac{ig}{\sqrt{2}}\right)^2 [\bar{u}(q)\gamma_\mu \frac{(1-\gamma_5)}{2} u(p)] \frac{-i}{\tilde{Q}^2 - M_W^2} \left(g^{\mu\nu} - \frac{\tilde{Q}^\mu \tilde{Q}^\nu}{M_W^2}\right) [\bar{u}(r)\gamma_\nu \frac{(1-\gamma_5)}{2} u(k)]. \quad (1)$$

The contribution proportional to $\tilde{Q}^\mu \tilde{Q}^\nu / M_W^2$ can be written as

$$-\frac{ig^2}{8M_W^2} \frac{1}{\tilde{Q}^2 - M_W^2} [\bar{u}(q) \tilde{Q}(1-\gamma_5)u(p)] [\bar{u}(r) \tilde{Q}(1-\gamma_5)u(k)], \quad (2)$$

and by using $\tilde{Q} = \not{p} - \not{q}$ along with the Dirac equation for the spinors $\bar{u}(q)$ and $u(p)$, it can be shown that the first term in square brackets is given by

$$\bar{u}(q) \tilde{Q}(1-\gamma_5)u(p) = -m_\ell \bar{u}(q)(1-\gamma_5)u(p). \quad (3)$$

Similarly, the second term in square brackets is proportional to the mass of the quark u_i , so that the contribution in Eq. (2) is of order $m_\ell m_u / M_W^2$ and then we will neglect it here. Therefore we have

$$\frac{1}{2} \sum_{\substack{d_i, u_i, \ell^- \\ \text{spins}}} |\mathcal{M}(\nu_\ell d_i \rightarrow \ell^- u_i)|^2 = \frac{g^4}{64(\tilde{Q}^2 - M_W^2)^2} A^{\mu\nu} B_{\mu\nu}, \quad (4)$$

where

$$A^{\mu\nu} \equiv \sum_{\ell^- \text{ spin}} [\bar{u}(q)\gamma^\mu(1-\gamma_5)u(p)][\bar{u}(p)\gamma^\nu(1-\gamma_5)u(q)], \quad (5)$$

$$B_{\mu\nu} \equiv \frac{1}{2} \sum_{\substack{d_i u_i \\ \text{spins}}} [\bar{u}(r)\gamma_\mu(1-\gamma_5)u(k)][\bar{u}(k)\gamma_\nu(1-\gamma_5)u(r)], \quad (6)$$

and we have omitted the spin indices in the spinors for simplicity. By computing the sums in Eqs. (5) and (6) we obtain

$$A^{\mu\nu} = 8\{q^\mu p^\nu + q^\nu p^\mu - (q \cdot p)g^{\mu\nu} - i\epsilon^{\mu\nu\alpha\beta}q_\alpha p_\beta\} \quad (7)$$

$$B_{\mu\nu} = 4\{r_\mu k_\nu + r_\nu k_\mu - (r \cdot k)g_{\mu\nu} - i\epsilon_{\mu\nu\rho\sigma}r^\rho k^\sigma\}. \quad (8)$$

From the replacement of Eqs. (7) and (8) in Eq. (4) we have

$$|\mathcal{M}|_{\text{unpol.}}^2 = \frac{2g^4}{(\tilde{Q}^2 - M_W^2)^2}(r \cdot q)(p \cdot k). \quad (9)$$

Since the vector \tilde{Q} is space like, we will use instead the quantity $Q^2 \equiv -\tilde{Q}^2$. On the other hand, by assuming the involved particles as massless the dot products appearing in Eq. (9) can be expressed in terms of Mandelstam variables in the following manner

$$(p \cdot k) = \frac{(p+k)^2}{2} = \frac{\hat{s}}{2} \quad (10)$$

$$(r \cdot q) = \frac{(r+q)^2}{2} = \frac{\hat{s}}{2}. \quad (11)$$

Hence, the unpolarized squared amplitude can be written as

$$|\mathcal{M}(\nu_\ell d_i \rightarrow \ell^- u_i)|_{\text{unpol.}}^2 = \frac{g^4 \hat{s}^2}{2(Q^2 + M_W^2)^2}. \quad (12)$$

Let us now compute the Lorentz invariant phase space element. If we consider the center of mass system for simplicity, we have

$$dLIPS = \frac{1}{2} \frac{1}{4E^2} \frac{1}{2E_\ell} \frac{1}{2E_u} \frac{d^3q}{(2\pi)^3} \frac{d^3r}{(2\pi)^3} (2\pi)^4 \delta^{(4)}(r+q-p-k), \quad (13)$$

where E denotes the energy of the neutrino and the down type quark in the CM, and E_ℓ and E_u are the energies of the lepton and the up type quark in the same reference frame. By choosing spherical coordinates for the momentum q we have

$$dLIPS = \frac{1}{8E^2} \frac{1}{2E_\ell} \frac{1}{2E_u} \frac{1}{(2\pi)^2} |\vec{q}|^2 \sin \theta_{\text{CM}} d\theta_{\text{CM}} d\phi d|\vec{q}| d^3r \delta^{(4)}(r+q-p-k). \quad (14)$$

By integrating in d^3r and $d\phi$ we obtain

$$dLIPS = \frac{1}{8E^2} \frac{1}{4\tilde{E}^2} \frac{1}{2\pi} d\cos \theta_{\text{CM}} |\vec{q}|^2 \frac{d|\vec{q}|^2}{2} \delta(2\tilde{E} - 2E)$$

$$= \frac{1}{2\hat{s}} \frac{1}{16\pi} d \cos \theta_{\text{CM}}, \quad (15)$$

where in the last equality we have used that in the CM, $\hat{s} = (2E)^2 = 4E^2$. Moreover, in the CM the variable \hat{t} is given by

$$\hat{t} = (q - p)^2 = (0, E \sin \theta_{\text{CM}}, 0, E \cos \theta_{\text{CM}} - E)^2 = -2E^2(1 - \cos \theta_{\text{CM}}) = -\hat{s} \frac{(1 - \cos \theta_{\text{CM}})}{2}, \quad (16)$$

and then $d\hat{t} = (\hat{s}/2) d \cos \theta_{\text{CM}}$. Hence, Eq. (15) can be written as follows

$$dLIPS = \frac{d\hat{t}}{16\pi\hat{s}^2}. \quad (17)$$

By combining Eqs. (12) and (17) we finally obtain the unpolarized differential cross section for the process $\nu_\ell d_i \rightarrow \ell^- u_i$,

$$\frac{d\sigma}{d\hat{t}}(\nu_\ell d_i \rightarrow \ell^- u_i) = \frac{M_W^4}{(Q^2 + M_W^2)^2} \frac{G_F^2}{\pi}, \quad (18)$$

where the Fermi constant G_F is defined through the relation

$$\frac{G_F}{\sqrt{2}} = \frac{g^2}{8M_W^2}. \quad (19)$$

Let us turn now to the computation of the right diagram of Fig.1. By neglecting again the contribution proportional to $\tilde{Q}^\mu \tilde{Q}^\nu / M_W^2$ we have in this case

$$\frac{1}{2} \sum_{\substack{\bar{u}_i, \bar{d}_i, \ell^- \\ \text{spins}}} |\mathcal{M}(\nu_\ell \bar{u}_i \rightarrow \ell^- \bar{d}_i)|^2 = \frac{g^4}{64(\tilde{Q}^2 - M_W^2)^2} A^{\mu\nu} \tilde{B}_{\mu\nu}, \quad (20)$$

where the tensor A is the same already given in Eq. (7) and \tilde{B} is given by

$$\tilde{B}_{\mu\nu} \equiv \frac{1}{2} \sum_{\substack{\bar{u}_i, \bar{d}_i \\ \text{spins}}} [\bar{v}(k) \gamma_\mu (1 - \gamma_5) v(r)] [\bar{v}(r) \gamma_\nu (1 - \gamma_5) v(k)]. \quad (21)$$

The computation of the sum in Eq. (21) gives the tensor in Eq. (8) but with the momenta k and r interchanged,

$$\tilde{B}_{\mu\nu} = 4\{k_\mu r_\nu + k_\nu r_\mu - (r \cdot k) g_{\mu\nu} - i\epsilon_{\mu\nu\rho\sigma} k^\rho r^\sigma\}. \quad (22)$$

By inserting Eqs. (7) and (21) in Eq. (20) we obtain

$$|\mathcal{M}|_{\text{unpol.}}^2 = \frac{2g^4}{(\tilde{Q}^2 - M_W^2)^2} (k \cdot q)(p \cdot r). \quad (23)$$

Again, if we assume that the particles are massless we can write the dot products in Eq. (23) in terms of Mandelstam's variables,

$$(k \cdot q) = -\frac{(k - q)^2}{2} = -\frac{\hat{u}}{2} \quad (24)$$

$$(p \cdot r) = -\frac{(p-r)^2}{2} = -\frac{\hat{u}}{2}. \quad (25)$$

These relations lead to the following expression for the unpolarized squared amplitude

$$|\mathcal{M}(\nu_\ell \bar{u}_i \rightarrow \ell^- \bar{d}_i)|_{\text{unpol.}}^2 = \frac{g^4 \hat{u}^2}{2(Q^2 + M_W^2)^2}, \quad (26)$$

which combined with Eq. (17) gives the corresponding differential cross section,

$$\frac{d\sigma}{d\hat{t}}(\nu_\ell \bar{u}_i \rightarrow \ell^- \bar{d}_i) = \frac{M_W^4}{(Q^2 + M_W^2)^2} \frac{G_F^2}{\pi} \left(\frac{\hat{u}}{\hat{s}} \right)^2. \quad (27)$$

Now, it is convenient to express this result in terms of dimensionless combinations of kinematic variables defined in the context of the interaction with a nucleon. If we define the momentum of the nucleon to be P , the momentum of a constituent parton can be written as $k = \xi P$ where ξ is the corresponding longitudinal fraction. Similarly, the quantity \hat{s} is given by

$$\hat{s} = (p + k)^2 = 2p \cdot k = 2\xi P \cdot p = \xi s \quad (28)$$

with s the squared of the center of mass energy of the system νN . By assuming the involved partons to be massless, the longitudinal fraction can be obtained in terms of the momenta of the neutrino and the lepton since

$$0 \approx (k + \tilde{Q})^2 = 2k \cdot \tilde{Q} + \tilde{Q}^2 = 2\xi P \cdot \tilde{Q} - Q^2, \quad (29)$$

from which

$$\xi = x, \text{ with } x \equiv \frac{Q^2}{2P \cdot \tilde{Q}}. \quad (30)$$

Finally, we define the variable y which in the rest frame of the nucleon is the fraction of the neutrino's energy that is transferred to the hadronic system,

$$y \equiv \frac{2P \cdot \tilde{Q}}{2P \cdot p} = \frac{2k \cdot (p - q)}{2k \cdot p} = \frac{\hat{s} + \hat{u}}{\hat{s}} = 1 - \frac{\hat{u}}{\hat{s}}. \quad (31)$$

From Eq. (31) we obtain the following relation between the Mandelstam variables \hat{u} and \hat{s} of the partons and the dimensionless quantity y ,

$$\frac{\hat{u}^2}{\hat{s}^2} = (1 - y)^2. \quad (32)$$

Further, we note that $\hat{t} = -Q^2$. By using the relation in Eq. (32) we can write Eq. (27) as follows

$$\frac{d\sigma}{d\hat{t}}(\nu_\ell \bar{u}_i \rightarrow \ell^- \bar{d}_i) = \frac{M_W^4}{(Q^2 + M_W^2)^2} \frac{G_F^2}{\pi} (1 - y)^2. \quad (33)$$

In order to obtain the differential cross section for the process $\nu_\ell N \rightarrow \ell^- + \text{anything}$ we must weight the parton differential cross sections obtained in Eqs. (18) and (33) with appropriated parton distribution functions. Therefore we encounter

$$\frac{d^2\sigma}{dQ^2 dx} = \frac{M_W^2}{(Q^2 + M_W^2)^2} \frac{G_F^2}{\pi} \{f_d(x) + (1 - y)^2 f_{\bar{u}}(x)\}, \quad (34)$$

where $f_d(x)$ and $f_{\bar{u}}(x)$ denote the parton distribution functions for the down-type quarks and up-type antiquarks in N respectively. For convenience we will express the distribution in terms of dimensionless quantities by changing the variables from x, Q^2 to x, y . In order to do so we note that $Q^2 = xys$, and then $dx dQ^2 = (dQ^2/dy) dx dy = xs dx dy$. With these variables, we obtain

$$\frac{d^2\sigma^{(C.C.)}}{dx dy} = \frac{M_W^2}{(Q^2 + M_W^2)^2} \frac{G_F^2}{\pi} s \{x f_d(x) + x f_{\bar{u}}(x)(1-y)^2\}. \quad (35)$$

If we consider an isoscalar nucleon $N \equiv (n+p)/2$ and take into account the violation of Bjorken scaling, the parton distribution functions can be written as follows

$$f_d(x) \rightarrow \frac{u_v(x, Q^2) + d_v(x, Q^2)}{2} + \frac{u_s(x, Q^2) + d_s(x, Q^2)}{2} + s_s(x, Q^2) + b_s(x, Q^2), \quad (36)$$

$$f_u(x) \rightarrow \frac{u_v(x, Q^2) + d_v(x, Q^2)}{2} + \frac{u_s(x, Q^2) + d_s(x, Q^2)}{2} + c_s(x, Q^2) + t_s(x, Q^2), \quad (37)$$

$$f_{\bar{d}}(x) \rightarrow \frac{u_s(x, Q^2) + d_s(x, Q^2)}{2} + s_s(x, Q^2) + b_s(x, Q^2), \quad (38)$$

$$f_{\bar{u}}(x) \rightarrow \frac{u_s(x, Q^2) + d_s(x, Q^2)}{2} + c_s(x, Q^2) + t_s(x, Q^2), \quad (39)$$

where the subscripts v and s denote valence and sea contributions, while u, d, c, s, t, b label the distributions for various quark flavors in a proton. In the case of the expression in Eq. (35) we need to use the functions in Eqs. (36) and (39) which we will denote $q(x, Q^2)$ and $\bar{q}(x, Q^2)$ respectively. By using this notation and taking into account that in the laboratory frame $s = 2P \cdot p = 2E_\nu M$ where E_ν is the energy of the incoming neutrino and M the nucleon mass, we find

$$\frac{d^2\sigma^{(C.C.)}}{dx dy} = \frac{2M_W^2}{(Q^2 + M_W^2)^2} \frac{G_F^2}{\pi} M E_\nu \{x q(x, Q^2) + x \bar{q}(x, Q^2)(1-y)^2\}. \quad (40)$$

Let us now consider the neutral current cross section. The diagram at parton level are depicted in Fig.2 where again $\ell = e, \mu, \tau$ and q denotes a generic quark.



FIG. 2: Dominant neutral current process diagrams. The left diagram shows the scattering of a neutrino by a given quark q whereas the right diagram depicts the scattering by a given antiquark \bar{q} .

The amplitude corresponding to the first diagram is given by

$$\mathcal{M}(\nu_\ell q \rightarrow \nu_\ell q) = \left(\frac{-ig}{2 \cos \theta_W} \right)^2 [\bar{u}(q) \gamma_\mu \frac{(1 - \gamma_5)}{2} u(p)] \frac{-i}{\tilde{Q}^2 - M_Z^2} \left(g^{\mu\nu} - \frac{\tilde{Q}^\mu \tilde{Q}^\nu}{M_Z^2} \right) [\bar{u}(r) \gamma_\nu (g_V - g_A \gamma_5) u(k)], \quad (41)$$

where the parameters g_V and g_A are different for up and down quarks,

$$g_V^u = \frac{1}{2} - \frac{4}{3} \sin^2 \theta_W, \quad g_A^u = \frac{1}{2} \quad (42)$$

and

$$g_V^d = -\frac{1}{2} + \frac{2}{3} \sin^2 \theta_W, \quad g_A^d = -\frac{1}{2}. \quad (43)$$

By neglecting the term proportional to $\tilde{Q}^\mu \tilde{Q}^\nu$ we have

$$\frac{1}{2} \sum_{q \text{ spins}} |\mathcal{M}(\nu_\ell q \rightarrow \nu_\ell q)|^2 = \frac{g^4}{64 \cos^4 \theta_W (\tilde{Q}^2 - M_Z^2)^2} A^{\mu\nu} B_{\mu\nu}, \quad (44)$$

where the tensor $A^{\mu\nu}$ is the same already given in Eqs. (5) and (6) for the charged current case and $B_{\mu\nu}$ is written as follows

$$B_{\mu\nu} = \frac{1}{2} \sum_{q \text{ spins}} [\bar{u}(r) \gamma_\mu (g_V - g_A \gamma_5) u(k)] [\bar{u}(k) \gamma_\nu (g_V - g_A \gamma_5) u(r)]. \quad (45)$$

In Eqs. (44) and (45) we average over the spins of the quark in the initial state and sum over the spins of the quark in the final state. However, in order to simplify the expressions, we include only one sum symbol. By solving the sum in Eq. 45 and disregarding the contribution proportional to m_q^2 we obtain

$$B_{\mu\nu} = 2(g_V^2 + g_A^2)(r_\mu k_\nu + r_\nu k_\mu - (r \cdot k)g_{\mu\nu}) - 4ig_V g_A r^\alpha k^\beta \epsilon_{\alpha\beta\mu\nu}. \quad (46)$$

By replacing Eqs. (7) and (46) in Eq. (44) we obtain the unpolarized squared amplitude,

$$|\mathcal{M}|_{\text{unpol.}}^2 = \frac{g^4}{2 \cos^4 \theta_W (Q^2 + M_Z^2)^2} ((g_V + g_A)^2 (q \cdot r)(p \cdot k) + (g_V - g_A)^2 (q \cdot k)(p \cdot r)), \quad (47)$$

that in terms of Madelstam's variables can be written as

$$|\mathcal{M}|_{\text{unpol.}}^2 = \frac{g^4}{8 \cos^4 \theta_W (Q^2 + M_Z^2)^2} ((g_V + g_A)^2 \hat{s}^2 + (g_V - g_A)^2 \hat{u}^2), \quad (48)$$

where the relations given in Eqs. (10)-(11) and (24)-(25) have been used. By combining this result with the phase space element obtained in Eq. (17) we obtain the unpolarized differential cross section,

$$\frac{d\sigma}{d\hat{t}}(\nu_\ell q \rightarrow \nu_\ell q) = \frac{G_F^2}{4\pi} \frac{M_Z^4}{(Q^2 + M_Z^2)^2} ((g_V + g_A)^2 + (g_V - g_A)^2 (1 - y)^2), \quad (49)$$

where the relation in Eq. (32) have been used.

Let us now compute the right diagram of Fig. 2. In this case we have

$$\frac{1}{2} \sum_{q \text{ spins}} |\mathcal{M}(\nu_\ell \bar{q} \rightarrow \nu_\ell \bar{q})|^2 = \frac{g^4}{64 \cos^4 \theta_W (Q^2 + M_Z^2)^2} A^{\mu\nu} \tilde{B}_{\mu\nu}, \quad (50)$$

where A is given in Eq. (7) and \tilde{B} is defined as

$$\tilde{B}_{\mu\nu} = \frac{1}{2} \sum_{q \text{ spins}} [\bar{v}(k) \gamma_\mu (g_V - g_A \gamma_5) v(r)] [\bar{v}(r) \gamma_\nu (g_V - g_A \gamma_5) v(k)], \quad (51)$$

where we have again neglected the contribution proportional to $\tilde{Q}^\mu \tilde{Q}^\nu$. By computing the corresponding sum we obtain

$$\tilde{B}_{\mu\nu} = 2(g_V^2 + g_A^2)^2 (r_\mu k_\nu + r_\nu k_\mu - (r \cdot k) g_{\mu\nu}) + 4i g_V g_A r^\alpha k^\beta \epsilon_{\alpha\beta\mu\nu}. \quad (52)$$

By using Eqs. (7) and (51) in Eq. (50) we obtain the unpolarized squared amplitude,

$$|\mathcal{M}|_{\text{unpol.}}^2 = \frac{g^4}{8 \cos^4 \theta_W (Q^2 + M_Z^2)^2} ((g_V - g_A)^2 \hat{s}^2 + (g_V + g_A)^2 \hat{u}^2), \quad (53)$$

from which the unpolarized differential cross section is obtained,

$$\frac{d\sigma}{dt}(\nu_\ell \bar{q} \rightarrow \nu_\ell \bar{q}) = \frac{G_F^2}{4\pi} \frac{M_Z^4}{(Q^2 + M_Z^2)^2} ((g_V - g_A)^2 + (g_V + g_A)^2 (1 - y)^2). \quad (54)$$

By using Eqs. (35) and (54) along with appropriate parton distribution functions the differential cross section of the process $\nu_\ell N \rightarrow \nu_\ell + \text{anything}$ is computed,

$$\begin{aligned} \frac{d^2\sigma^{(N.C.)}}{dx dy} &= \frac{G_F^2}{4\pi} \frac{M_Z^4}{(Q^2 + M_Z^2)^2} s x \left\{ (f_u(x) L_u^2 + f_{\bar{u}}(x) R_u^2 + f_d(x) L_d^2 + f_{\bar{d}}(x) R_d^2) + \right. \\ &\quad \left. + (1 - y)^2 (f_{\bar{u}}(x) L_u^2 + f_u(x) R_u^2 + f_{\bar{d}}(x) L_d^2 + f_d(x) R_d^2) \right\}, \end{aligned} \quad (55)$$

where

$$L_u \equiv g_V^u + g_A^u = 1 - \frac{4}{3} \sin^2 \theta_W \quad (56)$$

$$L_d \equiv g_V^d + g_A^d = -1 + \frac{2}{3} \sin^2 \theta_W \quad (57)$$

$$R_u \equiv g_V^u - g_A^u = -\frac{4}{3} \sin^2 \theta_W \quad (58)$$

$$R_d \equiv g_V^d - g_A^d = \frac{2}{3} \sin^2 \theta_W. \quad (59)$$

We will use the following definitions,

$$q^0(x) \equiv f_u(x) L_u^2 + f_{\bar{u}}(x) R_u^2 + f_d(x) L_d^2 + f_{\bar{d}}(x) R_d^2, \quad (60)$$

$$\bar{q}^0(x) \equiv f_{\bar{u}}(x) L_u^2 + f_u(x) R_u^2 + f_{\bar{d}}(x) L_d^2 + f_d(x) R_d^2. \quad (61)$$

If we consider an isoscalar nucleon and take into account the violation of Bjorken scaling we can insert the replacements given in Eqs. (36)-(39) into Eqs. (60) and (61) in order to obtain the functions $q^0(x, Q^2)$ and $\bar{q}^0(x, Q^2)$ respectively. Therefore, the differential cross section for the process $\nu_\ell N \rightarrow \nu_\ell + \text{anything}$ with $N = (n + p)/2$ is given by

$$\frac{d^2\sigma^{(N.C.)}}{dx dy} = \frac{G_F^2}{2\pi} \frac{M_Z^4}{(Q^2 + M_Z^2)^2} M E_\nu \{x q^0(x, Q^2) + x \bar{q}^0(x, Q^2) (1 - y)^2\}. \quad (62)$$

II. COMPUTATION OF THE CROSS SECTION - SINGLET LQ SCENARIO

Let us consider a simple scenario in which a LQ with $SU(3)_C \times SU(2)_L \times U(1)_Y$ quantum numbers $(\bar{\mathbf{3}}, \mathbf{1}, \mathbf{1}/3)$ couples to lepton and quarks giving rise to vertices of the type $LQ - l - q$. We will compute the cross section for the s-channel which is the resonant case under the assumption that the LQ couples only to a given family of leptons and a given family of quarks. This assumption goes in the line of [?] in which the LQ is assumed to connect the first family of quarks with the third family of leptons (see Eq. (A13)). Again, we distinguish between a *neutral* (N) and a *charged* (C) case. In the former case there is a neutrino in the final state whereas in the latter there is a charged lepton. At parton level we have then

$$\text{N case: } \nu_\ell + D \rightarrow \nu_\ell + D$$

$$\text{C case: } \nu_\ell + D \rightarrow \ell^- + U,$$

where $\ell = e, \mu, \tau$, $U = u, c, t$ and $D = d, s, b$. The corresponding diagrams are depicted in Fig. 3.




FIG. 3: Relevant s-channel diagrams. The left diagram shows the *charged* process while the right one shows the *neutral* process.

Let us first compute the cross section corresponding to the C process. The lagrangian is given in Eq. (A18) while the Feynman rules can be derived by using the expansions given in Eqs. (B1)-(B2) and (B17)-(B18). We denote the amplitude corresponding to left-handed (right-handed) particles in the final state as $\mathcal{M}_L(\mathcal{M}_R)$. With this notation the total amplitude is given by

$$\mathcal{M} = \mathcal{M}_L + \mathcal{M}_R, \quad (63)$$

with

$$\mathcal{M}_L = g_L^2 \frac{[\bar{u}_{DL}^c(k) u_{\nu L}(p)] [\bar{u}_{\ell L}(q) u_{UL}^c(r)]}{(\hat{s} - M^2 - iM\Gamma)}, \quad (64)$$

$$\mathcal{M}_R = g_L g_R \frac{[\bar{u}_{DL}^c(k) u_{\nu L}(p)] [\bar{u}_{\ell R}(q) u_{UR}^c(r)]}{(\hat{s} - M^2 - iM\Gamma)}, \quad (65)$$

where $\hat{s} = (p + k)^2$ and M is the LQ mass. The left-handed contribution can be written as

$$|\mathcal{M}_L|_{\text{unpol.}}^2 \equiv \frac{1}{2} \sum_{\substack{D, U, \ell \\ \text{spins}}} \frac{g_L^4 |a|^2 |b|^2}{(\hat{s} - M^2)^2 + M^2 \Gamma^2}, \quad (66)$$

where we have used the following definitions

$$a \equiv \bar{u}_{DL}^c(k) u_{\nu L}(p) = \bar{v}_{DR}(k) u_{\nu L}(p), \quad (67)$$

$$b \equiv \bar{u}_{\ell L}(q) u_{UL}^c(r) = \bar{u}_{\ell L}(q) v_{UR}(r). \quad (68)$$

In the second equality of Eqs. (67) and (68) we have used the relations in Eqs. (B15) and (B16). From Eq. (67) we have

$$|a|^2 = \text{Tr} [v_D(k) \bar{v}_D(k) u_{\nu L}(p) \bar{u}_{\nu L}], \quad (69)$$

and then

$$\sum_{D \text{ spins}} |a|^2 = \text{Tr} [(\not{k} - m_D) P_L \not{p}] = 2(k \cdot p). \quad (70)$$

On the other hand,

$$|b|^2 = \text{Tr} [u_{\ell}(q) \bar{u}_{\ell}(q) P_R v_U(r) \bar{v}_U(r) P_L], \quad (71)$$

and hence

$$\sum_{U, \ell \text{ spins}} |b|^2 = \text{Tr} [\not{q}(\not{r} - m_U) P_L] = 2(q \cdot r). \quad (72)$$

We can write these expressions in terms of Mandelstam's variables by taking into account that

$$\hat{s} = (q + r)^2 = m_{\ell}^2 + m_U^2 + 2(q \cdot r) \sim m_U^2 + 2(q \cdot r),$$

from which we have the relation

$$2(q \cdot r) = \hat{s} - m_U^2. \quad (73)$$

Similarly,

$$\hat{s} = (p + k)^2 = m_{\nu}^2 + m_D^2 + 2(k \cdot p),$$

and neglecting the neutrino and D quark masses we find

$$2(k \cdot p) = \hat{s}. \quad (74)$$

By using the relations in Eqs. (73) and (74) in Eqs. (72) and (70) respectively we obtain

$$\sum_{D \text{ spins}} |a|^2 = \hat{s} \quad , \quad \sum_{U, \ell \text{ spins}} |b|^2 = \hat{s} - m_U^2, \quad (75)$$

and replacing these expressions into Eq. (66) we obtain the following expression

$$|\mathcal{M}_L|_{\text{unpol.}}^2 = \frac{g_L^4}{2} \frac{\hat{s}(\hat{s} - m_U^2)}{(\hat{s} - M^2)^2 + M^2 \Gamma^2}. \quad (76)$$

Let us turn now to the computation of the right-handed contribution,

$$|\mathcal{M}_R|_{\text{unpol.}}^2 \equiv \frac{1}{2} \sum_{\substack{D, U, \ell \\ \text{spins}}} \frac{g_L^2 g_R^2 |a|^2 |\tilde{b}|^2}{(\hat{s} - M^2)^2 + M^2 \Gamma^2}, \quad (77)$$

where a was defined in Eq. (67) and

$$\tilde{b} \equiv \bar{u}_{\ell R}(q) u_{UR}^c(r) = \bar{u}_{\ell R}(q) v_{UL}(r). \quad (78)$$

From Eq. (78) we have

$$\sum_{U, \ell \text{ spins}} |\tilde{b}|^2 = \text{Tr} [\not{q}(\not{\psi} - m_U) P_R] = 2(q \cdot r). \quad (79)$$

By replacing Eqs. (70) and (79) in Eq. (77) we obtain

$$|\mathcal{M}_R|_{\text{unpol.}}^2 = \frac{g_L^2 g_R^2}{2} \frac{\hat{s}(\hat{s} - m_U^2)}{(\hat{s} - M^2)^2 + M^2 \Gamma^2}. \quad (80)$$

The complete unpolarized squared amplitude is given by

$$|\mathcal{M}|_{\text{unpol.}}^2 = |\mathcal{M}_L + \mathcal{M}_R|_{\text{unpol.}}^2 = |\mathcal{M}_L|_{\text{unpol.}}^2 + |\mathcal{M}_R|_{\text{unpol.}}^2 + 2\text{Re}(\mathcal{M}_L \mathcal{M}_R^*)_{\text{unpol.}}. \quad (81)$$

Now we will show that the last term can be neglected and in this sense there is no interference between the left- and right-handed parts of the amplitude. That term is proportional to $|a|^2 \tilde{b} \tilde{b}^*$ with

$$\tilde{b} \tilde{b}^* = \text{Tr} [u_\ell(q) \bar{u}_\ell(q) P_R v_U(r) \bar{v}_U(r) P_R], \quad (82)$$

and hence

$$\frac{1}{2} \sum_{\substack{U, \ell \\ \text{spins}}} \tilde{b} \tilde{b}^* = (1/2) \text{Tr} [(q + m_\ell) P_R (\not{\psi} - m_U) P_R] = -m_\ell m_U. \quad (83)$$

We then see that the contribution of the interference term being proportional to the lepton mass is negligible as compared to the other two terms appearing in Eq. (81).

In order to complete the computation of the cross section it only remains to obtain the LQ width Γ . We will assume as in [?] that the width is dominated by $LQ \rightarrow \ell^+ \bar{U}$ and $LQ \rightarrow \bar{\nu} \bar{D}$. Let us call the widths corresponding to these decays as Γ_I and Γ_{II} respectively. In the case of Γ_I we can again compute the left- and right-handed parts separately,

$$\mathcal{M}_L^I = i g_L [\bar{v}_{UL}^c(p_2) v_{\ell L}(p_1)] \quad (84)$$

$$\mathcal{M}_R^I = i g_R [\bar{v}_{UR}^c(p_2) v_{\ell R}(p_1)]. \quad (85)$$

From Eqs. (84) and (85) we obtain

$$|\mathcal{M}_L^I|_{\text{unpol.}}^2 = 2g_L^2(p_1 \cdot p_2) = g_L^2(M^2 - m_U^2) \quad (86)$$

$$|\mathcal{M}_R^I|_{\text{unpol.}}^2 = 2g_R^2(p_1 \cdot p_2) = g_R^2(M^2 - m_U^2). \quad (87)$$

The differential width is given by

$$d\Gamma_I = \frac{1}{32\pi^2} |\mathcal{M}^I|^2 \frac{|\vec{p}_1|}{M^2} d\Omega_1 = \frac{1}{32\pi^2} (g_L^2 + g_R^2) \frac{M^2 - m_U^2}{M^2} \frac{M^2 - m_U^2}{2M} d\phi_1 d(\cos \theta_1), \quad (88)$$

where we have used that $|\vec{p}_1| = (M^2 - m_U^2)/2M$ if the lepton mass is neglected. By performing the integrations we obtain

$$\Gamma_1 = \frac{M}{16\pi} (g_L^2 + g_R^2) \left(1 - \frac{m_U^2}{M^2}\right)^2. \quad (89)$$

Let us now compute Γ_2 . The corresponding amplitude is given by

$$\mathcal{M}^{II} = -ig_L [\bar{v}_{DL}^c(p_2) v_{\nu L}(p_1)], \quad (90)$$

from which we obtain

$$|\mathcal{M}^{II}|_{\text{unpol.}}^2 = 2g_L^2(p_1 \cdot p_2) = g_L^2 M^2. \quad (91)$$

Therefore, the differential width is

$$d\Gamma_{II} = \frac{1}{32\pi^2} (g_L^2 + g_R^2) \frac{(M^2 - m_U^2)}{M^2} \frac{(M^2 - m_U^2)}{2M} d\phi_1 d(\cos \theta_1), \quad (92)$$

where we have use that $|\vec{p}_1| = M/2$ under the assumption $m_\nu^2, m_\ell^2 \simeq 0$. By integrating Eq. (92) over the angles we have

$$\Gamma_{II} = \frac{M}{16\pi} g_L^2. \quad (93)$$

From Eqs. (89) and (93) we see that the LQ width is given by

$$\Gamma = \frac{M}{16\pi} \{g_L^2[(1 - \lambda_U)^2 + 1] + g_R^2(1 - \lambda_U)^2\}, \quad (94)$$

where $\lambda_U \equiv m_U^2/M^2$.

By using Eq. (94) and under the narrow-width approximation,

$$\frac{1}{(\hat{s} - M^2)^2 + M^2\Gamma^2} \approx \frac{\pi}{M\Gamma} \delta(\hat{s} - M^2), \quad (95)$$

we obtain from Eqs. (76) and (80) that

$$|\mathcal{M}(\nu D \rightarrow \ell^- U)|_{\text{unpol.}}^2 = \frac{8\pi^2 g_L^2 (g_L^2 + g_R^2) \hat{s} (\hat{s} - m_U^2)}{M^2 [(g_L^2 + g_R^2)(1 - \lambda_U)^2 + g_L^2]} \delta(\hat{s} - M^2). \quad (96)$$

From Eq. (96) we can write the corresponding unpolarized differential cross section for the C case:

$$\frac{d\sigma^C}{d\hat{t}} = \frac{\pi}{2\hat{s}} \frac{g_L^2 (g_L^2 + g_R^2) (\hat{s} - m_U^2)}{M^2 [(g_L^2 + g_R^2)(1 - \lambda_U)^2 + g_L^2]} \delta(\hat{s} - M^2). \quad (97)$$

Now we can write the differential cross section for the process $\nu N \rightarrow \ell^- + \text{anything}$:

$$\frac{d\sigma^C}{dQ^2 dx} = \frac{\pi}{2xs} \frac{g_L^2 (g_L^2 + g_R^2) (xs - m_U^2)}{M^2 [(g_L^2 + g_R^2)(1 - \lambda_U)^2 + g_L^2]} f_D(x) \delta(xs - M^2), \quad (98)$$

where $f_D(x)$ is the parton distribution function corresponding to the down-type quark D in the nucleon N . In order to use the inelasticity parameter we have to obtain the relation between Q^2 and x in the case in which the mass of the quark in the final state is not neglected and then it is still valid in the case of considering the top quark. The mass of the up quark can be related to the longitudinal fraction ξ as follows

$$m_U^2 = r^2 = (p + k - q)^2 = (\tilde{Q} + k)^2 = \tilde{Q}^2 + 2\tilde{Q} \cdot k = -Q^2 + 2\xi P \cdot \tilde{Q}, \quad (99)$$

from which

$$\xi = x, \text{ with } x \equiv \frac{Q^2 + m_U^2}{2P \cdot \tilde{Q}}. \quad (100)$$

The inelasticity parameter can be written in turn in the following manner

$$y = \frac{2P \cdot \tilde{Q}}{2P \cdot p} = \frac{2P \cdot \tilde{Q}}{2M_N E_\nu}. \quad (101)$$

From Eqs. (100) and (101),

$$Q^2 = xy(2M_N E_\nu) - m_U^2, \text{ and } dx dQ^2 = x(2M_N E_\nu) dx dy. \quad (102)$$

By using Eqs. (98) and (102) and integrating in x we finally obtain

$$\frac{d\sigma^C}{dy} = \frac{\pi}{2} \frac{g_L^2(g_L^2 + g_R^2)(1 - \lambda_U)}{(g_L^2 + g_R^2)(1 - \lambda_U)^2 + g_L^2} \frac{f_D(M^2/s)}{s}, \quad (103)$$

where the $1/s$ factor arises from the integration by using the delta function $\delta(xs - M^2)$. We also note that the delta function imposes $x = M^2/s$, so that

$$Q^2 = M^2 y - m_U^2, \quad (104)$$

and hence $\lambda_U \leq y \lesssim 1$. Let us now to compute the cross section for the neutral case. The amplitude is given by

$$\mathcal{M} = (ig_L)^2 \frac{[\bar{u}_{DL}^c(k) u_{\nu L}(p)][\bar{u}_{\nu L}(q) u_{DL}^c(r)]}{\hat{s} - M^2 - iM\Gamma}. \quad (105)$$

The unpolarized squared amplitude is

$$|\mathcal{M}|_{\text{unpol.}}^2 \equiv \frac{1}{2} \sum_{D \text{ spins}} \frac{g_L^4 |a|^2 |c|^2}{(\hat{s} - M^2)^2 + M^2 \Gamma^2}, \quad (106)$$

where a was already defined and

$$c \equiv \bar{u}_{\nu L}(q) v_{DR}(r). \quad (107)$$

From Eq. (107) we have

$$\sum_{D \text{ spins}} |c|^2 = \text{Tr}[(\not{q} + m_\nu) P_R (\not{r} - m_D) P_L] = 2(q \cdot r) = \hat{s}. \quad (108)$$

By using this expression we obtain

$$|\mathcal{M}|_{\text{unpol.}}^2 = \frac{\pi g_L^4}{M\Gamma} \frac{\hat{s}^2}{2} \delta(\hat{s} - M^2), \quad (109)$$

where we have used again the narrow-width approximation. The differential cross section is then given by

$$\frac{d\sigma^N}{d\hat{t}} = \frac{\pi g_L^4 \delta(\hat{s} - M^2)}{2M^2[(g_L^2 + g_R^2)(1 - \lambda_U)^2 + g_L^2]}. \quad (110)$$

From this equation we can derive the differential cross section for the process $\nu N \rightarrow \nu + \text{anything}$,

$$\frac{d\sigma^N}{dy} = \frac{\pi}{2} \frac{g_L^4}{(g_L^2 + g_R^2)(1 - \lambda_U)^2 + g_L^2} \frac{f_D(M^2/s)}{s}, \quad (111)$$

where we have performed the integration over x . We note that in this case $Q^2 = xys$ and since the delta function imposes $x = M^2/s$, $Q^2 = yM^2$, restoring the range of the inelasticity parameter, $0 \leq y \leq 1$.

III. COMPUTATION OF THE CROSS SECTION - TRIPLET LQ SCENARIO

The amplitudes corresponding to the neutral diagrams depicted in Fig. are given by

$$\mathcal{M}_1 = -\lambda_\ell^{i*} \lambda_{\ell'}^j \mathbb{P}_{\chi_1}(p+k) [\bar{v}_R(\nu_\ell; p) u_L(U_i; k)] [\bar{u}_L(U_j; r) v_R(\nu_{\ell'}; q)] \quad (112)$$

$$\mathcal{M}_2 = -\frac{\lambda_\ell^{i*} \lambda_{\ell'}^j \mathbb{P}_{\chi_2}(p+k)}{2} [\bar{v}_R(\nu_\ell; p) u_L(D_i; k)] [\bar{u}_L(D_j; r) v_R(\nu_{\ell'}; q)] \quad (113)$$

$$\mathcal{M}_3 = -\lambda_\ell^{j*} \lambda_{\ell'}^i \mathbb{P}_{\chi_1}(p-r) [\bar{v}_R(\nu_\ell; p) v_L(\bar{U}_i; r)] [\bar{v}_L(\bar{U}_j; k) v_R(\nu_{\ell'}; q)] \quad (114)$$

$$\mathcal{M}_4 = -\frac{\lambda_\ell^{j*} \lambda_{\ell'}^i \mathbb{P}_{\chi_2}(p-r)}{2} [\bar{v}_R(\nu_\ell; p) v_L(\bar{D}_j; r)] [\bar{v}_L(\bar{D}_i; k) v_R(\nu_{\ell'}; q)], \quad (115)$$

and the associated unpolarized squared amplitudes are

$$|\mathcal{M}_1|_{\text{unpol.}}^2 = |\lambda_\ell^i \lambda_{\ell'}^j|^2 |\mathbb{P}_{\chi_1}(p+k)|^2 \frac{\hat{s}(\hat{s} - m_{U_j}^2)}{2} \quad (116)$$

$$|\mathcal{M}_2|_{\text{unpol.}}^2 = |\lambda_\ell^i \lambda_{\ell'}^j|^2 |\mathbb{P}_{\chi_2}(p+k)|^2 \frac{\hat{s}^2}{8} \quad (117)$$

$$|\mathcal{M}_3|_{\text{unpol.}}^2 = |\lambda_\ell^j \lambda_{\ell'}^i|^2 |\mathbb{P}_{\chi_1}(p-r)|^2 \frac{\hat{u}(\hat{u} - m_{\bar{U}_j}^2)}{2} \quad (118)$$

$$|\mathcal{M}_4|_{\text{unpol.}}^2 = |\lambda_\ell^j \lambda_{\ell'}^i|^2 |\mathbb{P}_{\chi_2}(p-r)|^2 \frac{\hat{u}^2}{8}, \quad (119)$$

where $\mathbb{P}_x(p) \equiv (p^2 - m_x^2 - im_x \Gamma_x)^{-1}$. For the charged cases shown in Fig. we have

$$\mathcal{M}_I = -\frac{\lambda_\ell^{i*} \lambda_{\ell'}^j \mathbb{P}_{\chi_2}(p+k)}{2} [\bar{v}_R(\nu_\ell; p) u_L(D_i; k)] [\bar{u}_L(U_j; r) v_R(\nu_{\ell'}; q)] \quad (120)$$

$$\mathcal{M}_{II} = -\frac{\lambda_\ell^{j*} \lambda_{\ell'}^i \mathbb{P}_{\chi_2}(p-r)}{2} [\bar{v}_R(\nu_\ell; p) v_L(\bar{D}_j; r)] [\bar{v}_L(\bar{U}_i; k) v_R(\nu_{\ell'}; q)], \quad (121)$$

and the corresponding unpolarized squared amplitudes,

$$|\mathcal{M}_I|_{\text{unpol.}}^2 = |\lambda_\ell^i \lambda_{\ell'}^j|^2 |\mathbb{P}_{\chi_2}(p+k)|^2 \frac{\hat{s}(\hat{s} - m_{U_j}^2)}{8} \quad (122)$$

$$|\mathcal{M}_{II}|_{\text{unpol.}}^2 = |\lambda_\ell^j \lambda_{\ell'}^i|^2 |\mathbb{P}_{\chi_2}(p-r)|^2 \frac{\hat{u}^2}{8}. \quad (123)$$

Note that $|\mathcal{M}_{II}|_{\text{unpol.}}^2 = |\mathcal{M}_4|_{\text{unpol.}}^2$ and if the up-quark masses are neglected, $|\mathcal{M}_I|_{\text{unpol.}}^2 = |\mathcal{M}_2|_{\text{unpol.}}^2$. Now, let us focus on the resonant diagrams and use the narrow-width approximation for the LQs χ_1 and χ_2 . By neglecting the up-quark masses in Eqs. (116), (118) and (123) we obtain for the process $\nu_\ell N \rightarrow \nu_{\ell'} + X$:

$$\frac{d\sigma}{dx dy} = \sum_{i,j} |\lambda_\ell^i \lambda_{\ell'}^j|^2 \left\{ \frac{f_{u_i}(x, Q^2)}{32 m_{\chi_1} \Gamma_{\chi_1}} \delta(xs - m_{\chi_1}^2) + \frac{f_{d_i}(x, Q^2)}{128 m_{\chi_2} \Gamma_{\chi_2}} \delta(xs - m_{\chi_2}^2) \right\} \quad (124)$$

By performing the integration over x , we encounter

$$\frac{d\sigma}{dy}(\nu_\ell N \rightarrow \nu_{\ell'} + X) = \sum_{i,j} \frac{|\lambda_\ell^i \lambda_{\ell'}^j|^2}{32s} \left\{ \frac{m_{\chi_1}}{\Gamma_{\chi_1}} f_{u_i}(m_{\chi_1}^2/s, m_{\chi_1}^2 y) + \frac{m_{\chi_2}}{4\Gamma_{\chi_2}} f_{d_i}(m_{\chi_2}^2/s, m_{\chi_2}^2 y) \right\}. \quad (125)$$

By a similar computation we can obtain the differential cross section for the charged case in which only χ_2 is involved,

$$\frac{d\sigma}{dy}(\nu_\ell N \rightarrow \ell' + X) = \sum_{i,j} \frac{|\lambda_\ell^i \lambda_{\ell'}^j|^2}{128s} \frac{m_{\chi_2}}{\Gamma_{\chi_2}} f_{d_i}(m_{\chi_2}^2/s, m_{\chi_2}^2 y). \quad (126)$$

Under the assumption that the interaction of the LQs with fermions proceeds only between the first family of quarks and the first and second families of leptons, the above expressions can be simplified as follows,

$$\frac{d\sigma}{dy}(\nu_e N \rightarrow \nu_e X) = \frac{|\lambda_1^1|^4}{32s} F(y) \quad (127)$$

$$\frac{d\sigma}{dy}(\nu_\mu N \rightarrow \nu_\mu X) = \frac{|\lambda_2^1|^4}{32s} F(y) \quad (128)$$

$$\frac{d\sigma}{dy}(\nu_\mu N \rightarrow \nu_e X) = \frac{|\lambda_1^1|^2 |\lambda_2^1|^2}{32s} F(y) \quad (129)$$

$$\frac{d\sigma}{dy}(\nu_e N \rightarrow \nu_\mu X) = \frac{d\sigma}{dy}(\nu_\mu N \rightarrow \nu_e X) \quad (130)$$

for the neutral processes, where we have used the definition

$$F(y) \equiv \frac{m_{\chi_1}}{\Gamma_{\chi_1}} f_u(m_{\chi_1}^2/s, m_{\chi_1}^2 y) + \frac{m_{\chi_2}}{4\Gamma_{\chi_2}} f_d(m_{\chi_2}^2/s, m_{\chi_2}^2 y), \quad (131)$$

and

$$\frac{d\sigma}{dy}(\nu_e N \rightarrow e^- X) = \frac{|\lambda_1^1|^4}{128s} \frac{m_{\chi_2}}{\Gamma_{\chi_2}} f_d(m_{\chi_2}^2/s, m_{\chi_2}^2 y) \quad (132)$$

$$\frac{d\sigma}{dy}(\nu_\mu N \rightarrow \mu^- X) = \frac{|\lambda_2^1|^4}{128s} \frac{m_{\chi_2}}{\Gamma_{\chi_2}} f_d(m_{\chi_2}^2/s, m_{\chi_2}^2 y) \quad (133)$$

$$\frac{d\sigma}{dy}(\nu_e N \rightarrow \mu^- X) = \frac{|\lambda_1^1|^2 |\lambda_2^1|^2}{128s} \frac{m_{\chi_2}}{\Gamma_{\chi_2}} f_d(m_{\chi_2}^2/s, m_{\chi_2}^2 y) \quad (134)$$

$$\frac{d\sigma}{dy}(\nu_\mu N \rightarrow e^- X) = \frac{d\sigma}{dy}(\nu_e N \rightarrow \mu^- X) \quad (135)$$

for the charged processes. Under the assumptions that Γ_{χ_1} is dominated by the decay $\chi_1 \rightarrow \nu_\ell u_j$ and Γ_{χ_2} is dominated by $\chi_2 \rightarrow \nu_\ell d_j$ and $\chi_2 \rightarrow \ell^- u_j$, the width can be written in terms of the couplings as follows,

$$\Gamma_{\chi_1} = \frac{m_{\chi_1}}{16\pi} (|\lambda_1^1|^2 + |\lambda_2^1|^2) \quad (136)$$

$$\Gamma_{\chi_2} = \frac{m_{\chi_2}}{16\pi} (|\lambda_1^1|^2 + |\lambda_2^1|^2) \quad (137)$$

Appendix A: Electroweak-singlet LQ Lagrangian

Let us assume the case of a leptoquark S whose $SU(3)_C \times SU(2)_L \times U(1)_Y$ quantum numbers are $(\bar{\mathbf{3}}, \mathbf{1}, 1/3)$. In order to construct a gauge and Lorentz invariant lagrangian that describe the interaction of this LQ with quark and leptons, the transformation properties of the involved fields must be taken into account. Here, we will focus on the gauge group $SU(2)_L \times U(1)_Y$ and the Lorentz transformation. The fields along with their number quantumms are given in the following equation

$$Q_L = \begin{pmatrix} u_L \\ d_L \end{pmatrix} \rightarrow (\mathbf{2}, \mathbf{1/6}), \quad L_L = \begin{pmatrix} \nu_L \\ \ell_L \end{pmatrix} \rightarrow (\mathbf{2}, -\mathbf{1/2}),$$

$$u_R \rightarrow (\mathbf{1}, \mathbf{2/3}), \quad d_R \rightarrow (\mathbf{1}, -\mathbf{1/3}), \quad \ell_R \rightarrow (\mathbf{1}, -\mathbf{1}). \quad (\text{A1})$$

If the fields S and L_L are included in a given term, then one would say that the field \bar{Q}_L must also be included in order to satisfy $SU(2)_L$ gauge invariance. However, the net hypercharge in the term $\bar{Q}_L L_L S$ would not be zero: $-1/6 - 1/2 + 1/3 \neq 0$. Hence, we need a doublet that transforms as \bar{Q}_L under $SU(2)_L$ and as Q_L under $U(1)_Y$. The requirement on Y is easily satisfied by taking \bar{Q}_L^\dagger . But unfortunately this field does not transform as \bar{Q}_L under $SU(2)_L$. Hence, we will use \bar{Q}_L^\dagger but adjusting it to obtain the appropriate transformation property. We will show now that the field

$$\tilde{\bar{Q}}_L \equiv -i(\tau_2 \bar{Q}_L^\dagger)^T$$

transforms as \bar{Q}_L under $SU(2)_L$. Under an infinitesimal transformation we have

$$\delta \bar{Q}_L = -\frac{i}{2} g \omega_j \bar{Q}_L \tau_j, \quad (\text{A2})$$

from which

$$\delta \bar{Q}_L^\dagger = \frac{i}{2} g \omega_j \tau_j \bar{Q}_L^\dagger. \quad (\text{A3})$$

From Eq. (A3) and the definition of $\tilde{\bar{Q}}_L$ we can write

$$\begin{aligned} \delta \tilde{\bar{Q}}_L &= -i \left(\tau_2 \frac{i}{2} g \omega_j \tau_j \bar{Q}_L^\dagger \right)^T \\ &= -i \left(-\frac{i}{2} g \omega_j \tau_j^T \tau_2 \bar{Q}_L^\dagger \right)^T \\ &= -\frac{i}{2} g \omega_j (-i \tau_2 \bar{Q}_L^\dagger)^T \tau_j \\ &= -\frac{i}{2} g \omega_j \tilde{\bar{Q}}_L \tau_j, \end{aligned} \quad (\text{A4})$$

where we have used that $\tau_2 \tau_j = -\tau_j^T \tau_2$. By comparing Eqs. (A2) and (A4) we see that $\tilde{\bar{Q}}_L = -i(\tau_2 \bar{Q}_L^\dagger)^T = \bar{Q}_L^* i \tau_2$ transforms as \bar{Q}_L under $SU(2)_L$. Also, $\tilde{\bar{Q}}_L$ transforms as Q_L under $U(1)_Y$ as can be easily checked from its definition. However, if we were to write the interaction as $\tilde{\bar{Q}}_L L_L S = \bar{Q}_L^* i \tau_2 L_L S$ then the couplings between the fields with $s = 1/2$ would be of the form

$\bar{\psi}^* \psi$ which is not Lorentz invariant in contrast to $\bar{\psi} \psi$. The Lorentz invariance can be restored if the charge conjugate field is used since $\bar{\psi}^c \psi$ with $\bar{\psi}^c \equiv -i\psi^T \gamma^2 \gamma^0$ is invariant as we will show in the following.

Under an infinitesimal Lorentz transformation,

$$\psi \rightarrow \Lambda_{1/2} \psi = \left(1 - \frac{i}{2} \omega_{\mu\nu} S^{\mu\nu}\right) \psi, \quad (\text{A5})$$

where $S^{\mu\nu} = i[\gamma^\mu, \gamma^\nu]/2$. On the other hand, the charge conjugate field transforms as follows

$$\bar{\psi}^c = -i\psi^T \gamma^2 \gamma^0 \rightarrow -i \left[\left(1 - \frac{i}{2} \omega_{\mu\nu} S^{\mu\nu}\right) \psi \right]^T \gamma^2 \gamma^0 = -i\psi^T \left(1 - \frac{i}{2} \omega_{\mu\nu} S^{\mu\nu T}\right) \gamma^2 \gamma^0. \quad (\text{A6})$$

From the definition of $S^{\mu\nu}$ it can be shown that $S^{0i T} = S^{0i}$ if $i = 1, 3$ and $S^{02 T} = -S^{02}$ whereas $S^{ij T} = S^{ij}$ if $i = 2$ or $j = 2$ and $S^{ij T} = -S^{ij}$ otherwise. Moreover, it is easy to show that

- $[S^{\mu\nu}, \gamma^2] = 0$ if $\mu \neq 2$ and $\nu \neq 2$ and $\{S^{\mu\nu}, \gamma^2\} = 0$ otherwise.
- $[S^{\mu\nu}, \gamma^0] = 0$ if $\mu \neq 0$ and $\nu \neq 0$ and $\{S^{\mu\nu}, \gamma^0\} = 0$ otherwise.

From these properties we see that $S^{\mu\nu T} \gamma^2 \gamma^0 = -\gamma^2 \gamma^0 S^{\mu\nu}$ and finally arrive to

$$\bar{\psi}^c \rightarrow -i\psi^T \gamma^2 \gamma^0 \left(1 + \frac{i}{2} \omega_{\mu\nu} S^{\mu\nu}\right) = \bar{\psi}^c \Lambda_{1/2}^{-1}. \quad (\text{A7})$$

By combining Eqs. (A5) and (A7) we arrive to the result that $\bar{\psi}^c \psi$ is invariant under Lorentz transformations. In Eq. (A4) we have proved that the transformation law under $SU(2)_L$ of $\bar{Q}_L^* i\tau_2$ and \bar{Q}_L is the same but we have also shown that in order to get Lorentz invariant interactions the field $\bar{Q}_L^c \equiv -iQ_L^T \gamma^2 \gamma^0$ must be used instead of \bar{Q}_L^* . Hence, we should prove that $\bar{Q}_L^c i\tau_2$ also transforms as \bar{Q}_L under $SU(2)_L$.

First we note that in general $\bar{\psi}^* = (\psi^\dagger \gamma^0)^* = \psi^T \gamma^0$ and then

$$\bar{\psi}^c = -i\psi^T \gamma^2 \gamma^0 = -i\bar{\psi}^* \gamma^0 \gamma^2 \gamma^0 = \bar{\psi}^* i\gamma^2.$$

In this way we can obtain \bar{Q}_L^c from \bar{Q}_L^* as follows

$$\bar{Q}_L^c = \bar{Q}_L^* (i\gamma^2 \mathbb{1}_{SU(2)_L}) \quad (\text{A8})$$

and then

$$\bar{Q}_L^c i\tau_2 = \bar{Q}_L^* i\gamma^2 \mathbb{1}_{SU(2)_L} i\tau_2 = \bar{Q}_L^* i\tau_2 i\gamma^2 \mathbb{1}_{SU(2)_L} = \tilde{\bar{Q}}_L i\gamma^2 \mathbb{1}_{SU(2)_L}. \quad (\text{A9})$$

From Eqs. (A9) and (A4) we can derive the required transformation law,

$$\begin{aligned} \delta(\bar{Q}_L^c i\tau_2) &= \delta \tilde{\bar{Q}}_L i\gamma^2 \mathbb{1}_{SU(2)_L} \\ &= -\frac{i}{2} g \omega_j \tilde{\bar{Q}}_L \tau_j i\gamma^2 \mathbb{1}_{SU(2)_L} \\ &= -\frac{i}{2} g \omega_j (\bar{Q}_L^c i\tau_2) \tau_j. \end{aligned} \quad (\text{A10})$$

From Eqs. (A10) and (A2) we see that $\bar{Q}_L^c i\tau_2$ transforms as \bar{Q}_L under $SU(2)_L$. Also, from Eq. (A8) it transforms as Q_L under $U(1)_Y$. Therefore, we conclude that the interaction $\bar{Q}_L^c i\tau_2 L_L S$ is $SU(2)_L \times U(1)_Y$ invariant and also Lorentz invariant.

The case of right-handed fields is easier since they are all $SU(2)_L$ singlets and hence we need to be careful only with the $U(1)_Y$ and Lorentz invariance. By taking into account the discussion given above, we can conclude that the suitable interaction is $\bar{u}_R^c \ell_R S$. On one hand, the hypercharge of this term is $2/3 - 1 + 1/3 = 0$ and on the other hand, the presence of the charge conjugate field \bar{u}_R^c guarantees the Lorentz invariance. We are now ready to write down the terms of the LQ lagrangian that describe the interaction between quarks and leptons. If we consider only one LQ mediating the lepton-quark interaction, these terms are written as follows,

$$\mathcal{L}_{LQ} \supset g_{ij}^L \bar{Q}_{iL}^c i\tau_2 L_{jL} S + g_{ij}^R \bar{u}_{iR}^c \ell_{jR} S + \text{h.c.}, \quad (\text{A11})$$

where the indices i and j denote the quark and lepton families respectively and the sum over them is implicit. This form is adopted for example in [?]. Conversely, in the case of assuming no mixing between the families, universal couplings and one LQ by family, we would have

$$\mathcal{L}_{LQ} \supset (g^L \bar{Q}_{iL}^c i\tau_2 L_{iL} + g^R \bar{u}_{iR}^c \ell_{iR}) S_i + \text{h.c.} \quad (\text{A12})$$

This form is used for example in [?]. Finally, in the case assumed in [?], in which the LQ couples only to the first family of quarks and the third family of leptons, we have

$$\mathcal{L}_{LQ} \supset (g^L \bar{Q}_{1L}^c i\tau_2 L_{3L} + g^R \bar{u}_{1R}^c \ell_{3R}) S + \text{h.c.}, \quad (\text{A13})$$

where $Q_{1L} = (u, d)^T$, $L_{3L} = (\nu_{\tau L}, \tau_L)^T$, $u_{1R} = u_R$ and $\ell_{3R} = \tau_R$. It is instructive to expand a generic term and compute the corresponding hermitian conjugate,

$$\bar{Q}_L^c i\tau_2 L_L S = (\bar{u}_L^c \ell_L - \bar{d}_L^c \nu_L) S, \quad (\text{A14})$$

$$(\bar{u}_L^c \ell_L S)^\dagger = S^\dagger \ell_L^\dagger (\bar{u}_L^c)^\dagger = S^\dagger \bar{\ell}_L \gamma^0 ((-i\gamma^2 u_L)^T \gamma^0)^\dagger = S^\dagger \bar{\ell}_L (-i\gamma^2 u_L)^* = S^\dagger \bar{\ell}_L (-i\gamma^2 u_L^*) = S^\dagger \bar{\ell}_L u_L^c. \quad (\text{A15})$$

Similarly, it can be shown that

$$(\bar{d}_L^c \nu_L S)^\dagger = S^\dagger \bar{\nu}_L d_L^c, \quad (\text{A16})$$

$$(\bar{u}_R^c \ell_R S)^\dagger = S^\dagger \bar{\ell}_R u_R^c. \quad (\text{A17})$$

By using Eqs. (A14)-(A17) we can write the lagrangian that describes the interaction between a given family of quarks and a given family of leptons in the following manner,

$$\mathcal{L} = g_L \{ (\bar{u}_L^c \ell_L - \bar{d}_L^c \nu_L) S + S^\dagger (\bar{\ell}_L u_L^c - \bar{\nu}_L d_L^c) \} + g_R \{ \bar{u}_R^c \ell_R S + S^\dagger \bar{\ell}_R u_R^c \}, \quad (\text{A18})$$

where we have assumed that the couplings $g_{L,R}$ are real. This lagrangian is the one that must be taken into account in order to compute a generic s-channel diagram with the LQ as mediator.

Appendix B: How to use Conjugate Fields

The fields describing a $s = 1/2$ fermion can be written as follows

$$\psi(x) = \int \frac{d^3 p}{(2\pi)^3} \frac{1}{\sqrt{2E_{\mathbf{p}}}} \sum_s (a_{\mathbf{p}}^s u^s(p) e^{-ipx} + b_{\mathbf{p}}^{s\dagger} v^s(p) e^{ipx}), \quad (\text{B1})$$

$$\bar{\psi}(x) = \int \frac{d^3p}{(2\pi)^3} \frac{1}{\sqrt{2E_{\mathbf{p}}}} \sum_s (b_{\mathbf{p}}^s \bar{v}^s(p) e^{-ipx} + a_{\mathbf{p}}^{s\dagger} \bar{u}^s(p) e^{ipx}). \quad (\text{B2})$$

Under the action of the charge conjugation operator C , we have

$$C\psi(x)C = \int \frac{d^3p}{(2\pi)^3} \frac{1}{\sqrt{2E_{\mathbf{p}}}} \sum_s (b_{\mathbf{p}}^s u^s(p) e^{-ipx} + a_{\mathbf{p}}^{s\dagger} v^s(p) e^{ipx}), \quad (\text{B3})$$

$$C\bar{\psi}(x)C = \int \frac{d^3p}{(2\pi)^3} \frac{1}{\sqrt{2E_{\mathbf{p}}}} \sum_s (a_{\mathbf{p}}^s \bar{v}^s(p) e^{-ipx} + b_{\mathbf{p}}^{s\dagger} \bar{u}^s(p) e^{ipx}). \quad (\text{B4})$$

From Eqs. (B3) and (B4) it can be shown that

$$C\psi(x)C = -i\gamma^2\psi^* \equiv \psi^c, \quad (\text{B5})$$

$$C\bar{\psi}(x)C = (-i\gamma^2\psi)^T\gamma^0 \equiv \bar{\psi}^c. \quad (\text{B6})$$

Let us extend these definitions to the spinors, so that we have

$$u^s(p)^c \equiv -i\gamma^2 u^s(p)^* = v^s(p), \quad (\text{B7})$$

$$v^s(p)^c \equiv -i\gamma^2 v^s(p)^* = u^s(p), \quad (\text{B8})$$

$$\bar{u}^s(p)^c \equiv (-i\gamma^2 u(p))^T\gamma^0 = [(-i\gamma^2 u(p)^*)^*]^T\gamma^0 = \bar{v}^s(p), \quad (\text{B9})$$

$$\bar{v}^s(p)^c \equiv (-i\gamma^2 v(p))^T\gamma^0 = \bar{u}^s(p). \quad (\text{B10})$$

By using these definitions along with Eqs. (B3)-(B6) we can write,

$$\psi^c = \int \frac{d^3p}{(2\pi)^3} \frac{1}{\sqrt{2E_{\mathbf{p}}}} \sum_s (b_{\mathbf{p}}^s v^s(p)^c e^{-ipx} + a_{\mathbf{p}}^{s\dagger} u^s(p)^c e^{ipx}), \quad (\text{B11})$$

$$\bar{\psi}^c = \int \frac{d^3p}{(2\pi)^3} \frac{1}{\sqrt{2E_{\mathbf{p}}}} \sum_s (a_{\mathbf{p}}^s \bar{u}^s(p)^c e^{-ipx} + b_{\mathbf{p}}^{s\dagger} \bar{v}^s(p)^c e^{ipx}). \quad (\text{B12})$$

Regarding the left- and right-handed fields, we will use the same definitions given in Eqs. (B5)-(B6),

$$\psi_{L,R}^c = -i\gamma^2(\psi_{L,R})^* = P_{R,L}\psi^c, \quad (\text{B13})$$

$$\bar{\psi}_{L,R}^c = (-i\gamma^2\psi_{L,R})^T\gamma^0 = \bar{\psi}^c P_{L,R}, \quad (\text{B14})$$

where $P_{R,L} = (1 \pm \gamma^5)/2$. Similarly, by applying the definitions in Eqs. (B7)-(B10) to the left- and right-handed spinors we obtain

$$u_{L,R}^c = v_{R,L}, \quad \bar{u}_{L,R}^c = \bar{v}_{R,L}, \quad (\text{B15})$$

$$v_{L,R}^c = u_{R,L}, \quad \bar{v}_{L,R}^c = \bar{u}_{R,L}. \quad (\text{B16})$$

By combining the expansions in Eqs. (B3) and (B4) with Eqs. (B13) and (B14) and the relations in Eqs. (B15) and (B16) we arrive to the following expansions

$$\psi_{L,R}^c = \int \frac{d^3p}{(2\pi)^3} \frac{1}{\sqrt{2E_{\mathbf{p}}}} \sum_s (b_{\mathbf{p}}^s v_{L,R}^s(p)^c e^{-ipx} + a_{\mathbf{p}}^{s\dagger} u_{L,R}^s(p)^c e^{ipx}), \quad (\text{B17})$$

$$\bar{\psi}_{L,R}^c = \int \frac{d^3p}{(2\pi)^3} \frac{1}{\sqrt{2E_{\mathbf{p}}}} \sum_s (a_{\mathbf{p}}^s \bar{u}_{L,R}^s(p)^c e^{-ipx} + b_{\mathbf{p}}^{s\dagger} \bar{v}_{L,R}^s(p)^c e^{ipx}). \quad (\text{B18})$$

In the case of the fields ψ^c and $\bar{\psi}^c$ the expansions in Eqs. (B11) and (B12) must be used and the relations in Eqs. (B7)-(B10) allow to replace the charge conjugate spinors by the *usual* spinors. On the other hand, in the case of the fields $\psi_{L,R}^c$ and $\bar{\psi}_{L,R}^c$ we must take the expansions in Eqs. (B17) and (B18) along with the relations in Eqs. (B15) and (B16).

Appendix C: Electroweak-triplet LQ Lagrangian

Let us review how to represent a $SU(2)$ triplet in terms of a 2×2 matrix χ which transforms under the group as follows,

$$\chi \rightarrow U \chi U^\dagger, \quad (\text{C1})$$

where $U = \exp(i\omega_j \tau_j/2)$ and τ_j is the j -th Pauli matrix. In the spin one representation of $SU(2)$, the basis of the representation space is given by eigenvectors of both J^2 and J_3 . Hence, in that basis the generator is diagonal with eigenvalues $-1, 0, 1$. The so called triplet is given by a column vector,

$$\chi = \begin{pmatrix} \chi_1 \\ \chi_2 \\ \chi_3 \end{pmatrix}, \quad (\text{C2})$$

where χ_1 have J_3 quantum number $+1$, χ_2 has 0 and χ_3 has -1 . In order to express this triplet as a matrix we need to represent the state χ_i by a matrix M_i that is the analog of an eigenstate of J_3 . This condition is written as follows,

$$[J_3, M_i] = \lambda_i M_i, \quad (\text{C3})$$

where $\lambda_i = 1, 0, -1$ for $i = 1, 2, 3$ respectively. On the other hand, the transformation law given in Eq. (C1) is obviously valid in the case of the representation $\mathbf{2} \otimes \mathbf{2}$. By using tensorial notation, this representation is written as T_j^i , which can be splitted into two tensors that are invariant under the transformation law in Eq. (C1), one is traceless and the other is proportional to the identity. This is nothing but the decomposition of the direct product representation into a direct sum, $\mathbf{2} \otimes \mathbf{2} = \mathbf{3} \oplus \mathbf{1}$ or in tensorial notation $T_j^i = A_j^i + B \delta_j^i$, where A_j^i is traceless. This tensor can then be used to represent the triplet. We need to consider 2×2 traceless matrices that fulfill Eq. (C3) with $J_3 = \tau_2/2$. The Pauli matrices are in fact traceless and can be chosen as a basis of the space of 2×2 traceless matrices,

$$M_i = \sum_{j=1}^3 c_{ij} J_j, \quad (\text{C4})$$

where $J_j = \tau_j/2$. By replacing Eq. (C4) in Eq. (C3), we obtain

$$M_1 = c_1(J_1 + iJ_2), \quad M_2 = c_2 J_3, \quad M_3 = c_3(J_1 - iJ_2), \quad (\text{C5})$$

and hence the triplet can be written as follows,

$$\chi = \chi_1 M_1 + \chi_2 M_2 + \chi_3 M_3. \quad (\text{C6})$$

By choosing the normalization $c_1 = c_3 = 1$ and $c_2 = 1/\sqrt{2}$ we get

$$\chi = \begin{pmatrix} \chi_2/\sqrt{2} & \chi_1 \\ \chi_3 & -\chi_2/\sqrt{2} \end{pmatrix}. \quad (\text{C7})$$

We note that if we assign the hypercharge $-1/3$ to the triplet, the electric charges of the scalars $\chi_{1,2,3}$ are respectively $2/3$, $-1/3$ and $-4/3$. Now that we have the matrix representation (Eq. (C7)) of the triplet and the corresponding transformation law (Eq.(C1) we can construct their $SU(2)_L \times U(1)_Y$ invariant interaction with left-handed quarks and leptons,

$$\mathcal{L}_{LQ} \supset \sum_{\ell,j} \lambda_\ell^j \bar{Q}_{jL} \chi i\tau_2 L_{\ell L}^c + \text{h.c.} \quad (\text{C8})$$

By expanding Eq. (C8) we finally obtain

$$\begin{aligned} \mathcal{L}_{LQ} &\supset \lambda_\ell^i \left[\bar{u}_{iL} \left(-\chi_1 \nu_{\ell L}^c + \frac{\chi_2}{\sqrt{2}} \ell_L^c \right) + \bar{d}_{iL} \left(\frac{\chi_2}{\sqrt{2}} \nu_{\ell L}^c + \chi_3 \ell_L^c \right) \right] + \text{h.c.} = \\ &= \lambda_\ell^i \left[\bar{u}_i P_R \left(-\chi_1 \nu_\ell^c + \frac{\chi_2}{\sqrt{2}} \ell^c \right) + \bar{d}_i P_R \left(\frac{\chi_2}{\sqrt{2}} \nu_\ell^c + \chi_3 \ell^c \right) \right] + \text{h.c.} \end{aligned} \quad (\text{C9})$$

REFERENCES