Phase space mixing in the presence of self-gravity: spherically symmetric Vlasov-Poisson evolutions from initial data out of equilibrium

Erik Jiménez-Vázquez[1] and Olivier Sarbach[2]

[1] [(Erik) Affiliation to be completed.]

[2] *Instituto de Física y Matemáticas, Universidad Michoacana de San Nicolás de Hidalgo, Edificio C-3, Ciudad Universitaria, 58040 Morelia, Michoacán, México*

(Dated: November 8, 2026)

[(Erik) Provisional abstract; to be rewritten when section VI is final.] These notes are devoted to the numerical study of a spherically symmetric, collisionless kinetic gas which is trapped in an external central potential, whose self-gravity is taken into account, and whose initial distribution function is concentrated about one orbital phase, far from any steady state. In the absence of self-gravity, such a gas relaxes in time to a stationary state through phase space mixing. We analyze how this relaxation is modified by the self-gravity of the gas, for the reduced model in which all the gas particles have the same magnitude of the angular momentum, by means of a particle-in-cell code whose particles are placed on a regular lattice in action-angle variables. The outcome is controlled by a parameter μ which compares the amplitude of the potential generated by the gas with the spread of its orbital frequencies. Below a critical value of μ the gas mixes, more slowly than in the absence of self-gravity, and the potential which it generates relaxes to a stationary profile, as in Landau damping. Above this value it collapses into a compact, self-bound clump which oscillates radially and does not mix. For clouds which are compact in angle the critical value lies between 0.43 and 0.48, both in the isochrone and in the Kepler potential.

PACS numbers: 05.20.Dd, 98.10.+z, 02.60.Cb

I. INTRODUCTION

Collisionless kinetic gases provide the standard description for systems consisting of a large number of particles which interact with each other only through the mean field they generate. Prominent examples are the stars in a galaxy or in a globular cluster, and the dark matter particles in a galactic halo [1]. The state of such a gas at time t is described by its one-particle distribution function (DF) $f(t, x, p)$, a nonnegative function on the one-particle phase space which determines the number of gas particles in a small phase space volume about the point (x, p), and whose evolution is governed by the Vlasov equation.[1] In the Newtonian gravitational case, the Vlasov equation is coupled to Poisson's equation for the gravitational potential generated by the gas, which yields the Vlasov-Poisson system, see Ref. [2] for a review of its mathematical properties. Its general relativistic counterpart is the Vlasov-Einstein system [3].

It is instructive to compare the dynamics of a self-gravitating collisionless gas with the one of a pressureless fluid (dust), which is obtained in the limit in which the velocity dispersion of the gas vanishes. The gravitational collapse of a spherical dust cloud leads to the formation of a singularity in finite time at which the density diverges, both in the Newtonian theory and in general relativity [4]; in the latter case, part of this singularity may even be visible to distant observers [5, 6]. In contrast to this, the velocity dispersion of a kinetic gas prevents the particles from focusing. In fact, the Vlasov-Poisson system admits global classical solutions for general initial data [7–9], while for the spherically symmetric Vlasov-Einstein system global solutions exist for small initial data [10] and sufficiently concentrated data collapse to a black hole [11]. Therefore, for a Newtonian collisionless gas the relevant question is not whether or not a singularity forms, but rather how the gas behaves for large times and, in particular, whether or not it relaxes to a stationary state.

A fundamental mechanism leading to such a relaxation is phase space mixing [12, 13]. To explain it, consider first a gas whose self-gravity is neglected and which is trapped in the potential well of an external potential. If the one-particle Hamiltonian is integrable, each gas particle is confined to an invariant torus, and in terms of action-angle variables (Q, J) the solution of the Vlasov equation is $f(t, Q, J) = F(Q - \Omega(J)t, J)$, where F describes the initial DF and $\Omega(J)$ denotes the frequencies associated with the torus labeled by J. Since these frequencies vary from one torus to the other, the DF is stretched along the angle directions and develops structures on finer and finer scales as time increases. As a consequence, f does not converge pointwise as $t \to \infty$. However, the macroscopic observables which

[1] The Vlasov equation is also often called the collisionless Boltzmann equation in the literature.

are obtained by integrating f against a test function do converge, and their limit is the one associated with the average of the initial DF over the angle variables, which depends only on integrals of motion. Note that this holds for any initial DF, not only for small perturbations of a steady state. A mathematically precise formulation of this statement and its proof for rotationally symmetric potentials in $d = 1, 2, 3$ space dimensions satisfying a suitable non-degeneracy condition were given by Rioseco and Sarbach [14] (see also Refs. [15, 16]). The results in Ref. [14] include the case of a spherically symmetric gas configuration in which all the gas particles have the same value of the total angular momentum, which behaves like an effective one-dimensional system, and they explain the relaxation to a virialized state which had been observed in the numerical simulations of Ref. [17]. The final state is thus determined by the initial DF through its average over the angle variables; for a gas in an axisymmetric potential with two potential wells along the symmetry axis, in which the motion is integrable, this prediction has recently been compared with N-particle simulations in Ref. [18]. For the analogous phenomenon in the exterior of a black hole we refer the reader to Refs. [19, 20], and for estimates on the rate at which the observables converge to Refs. [21–23].

The generalization to the self-gravitating case, which was left open in Ref. [14], is considerably more involved, since the potential now depends on the DF, such that a change of the latter induces a change of the potential which, in turn, acts back on the gas. In plasma physics, where the particles interact through their mean electric field, the corresponding collective effect has been known since the work of Landau [24]: for a gas of charged particles close to a spatially homogeneous equilibrium the electric field decays in time while the DF undergoes mixing. This phenomenon, known as Landau damping, has been established rigorously at the nonlinear level by Mouhot and Villani [25]. The gravitational case differs from the electromagnetic one in two important aspects. First, the interaction is attractive, such that the collective response of the gas tends to reinforce a perturbation instead of screening it. Second, a self-gravitating gas configuration of finite mass does not possess spatially homogeneous equilibria. Instead, its steady states are spatially inhomogeneous, the gas particles are trapped in the potential well, and the frequencies of their radial motion lie in an interval which, in the situations considered in these notes, is bounded and bounded away from zero. For this reason, a collective oscillation whose frequency lies below this interval is not in resonance with any gas particle and hence cannot be damped by mixing.

Two situations have been analyzed in the gravitational case. The first one concerns small perturbations of a self-consistent steady state. Hadžić, Rein and Straub [26] (see also Refs. [27, 28]) showed that the essential spectrum of the operator governing the linearized dynamics about a large class of spherically symmetric steady states is determined by the radial periods of the gas particles, and, using a version of the Birman-Schwinger principle which was first applied to this problem by Mathur [29], they derived a criterion for the existence of an eigenvalue below this spectrum, which corresponds to an oscillating mode that is not damped. Related criteria and estimates for the number of such modes have been given in Ref. [30]. For the reduced system in which all the gas particles have the same magnitude of the angular momentum and move around a central point mass, Hadžić, Rein, Schrecker and Straub [31] proved the following dichotomy for steady states of sufficiently small mass whose DF is proportional to $(E_0 - E)^k$ for energies E below the cutoff energy E_0: the solutions of the linearized system are damped if $k > 1$, whereas an oscillating mode exists if $1/2 < k \leq 1$. Quantitative decay estimates for the linearized system have been obtained in Refs. [23, 32], and particle-in-cell simulations show that, depending on the steady state, its perturbations either oscillate without damping or decay [33, 34]. Perturbations of steady states in the setting of these notes are analyzed in a companion article. [(Erik) Reference to the companion article on perturbations of steady states to be added.] The second situation concerns initial data which are not a perturbation of a steady state of the gas. For a self-gravitating gas around a point mass, Chaturvedi and Luk [35] proved that small, spherically symmetric initial data which are supported on bounded orbits, with angular momenta in an interval bounded away from zero, undergo phase space mixing up to a time of the order of $\epsilon^{-1}(\log \epsilon^{-1})^{-2}$, where ϵ measures the size of the data, while the global existence and the asymptotic behavior of these solutions remain open. Finally, when the initial data are far from any steady state and the self-gravity is strong, the gas undergoes a violent relaxation [13] toward a quasi-stationary state, and the prediction of this state from the initial data is understood only in special cases, see Ref. [36] for a review.

In these notes, we study numerically the relaxation through phase space mixing of a spherically symmetric collisionless gas which is trapped in an external central potential, whose self-gravity is taken into account, and whose initial DF is far from any steady state. We consider the reduced model in which all the gas particles have the same magnitude $L_0 > 0$ of the angular momentum, which has a two-dimensional phase space and is the setting considered in Refs. [14, 31]. This model goes back to the numerical experiments of Hénon with spherical shells (see Ref. [37] and references therein) and has been used in the N-body simulations of Ref. [38]. For the external potential we choose the isochrone potential, for which the action-angle variables are known explicitly, and the potential of a point mass, which is the one considered in Refs. [31, 35]. As initial data we consider a gas cloud whose particles are concentrated about one orbital phase, with a smooth distribution of their action variables. In the absence of self-gravity this cloud mixes, and we address two questions: up to which mass does the mixing persist when the self-gravity of the cloud is taken into account, and which state does the gas reach when it does not mix. Our analysis is based on a particle-in-cell (PIC) code which solves the nonlinear Vlasov-Poisson system, and on a measure of the mixing in terms of the action-angle

variables of the instantaneous potential of the gas.

The main results of these notes can be summarized as follows. First, in the absence of self-gravity the observables relax algebraically, with an exponent which is determined by the behavior of the initial DF at the boundary of its support, and the PIC code reproduces the exact solution of this problem to within 6×10^{-9} in relative terms. Second, when the self-gravity of the gas is taken into account, the observables which measure the mixing have to be computed with the action-angle variables of the total potential; those of the external potential leave a time-independent residue which could be mistaken for an incomplete mixing. Third, the evolution of the cloud is controlled by a parameter μ which compares the amplitude of the potential generated by the cloud with the spread of its orbital frequencies. Below a critical value μ_c the cloud mixes, although more slowly than in the absence of self-gravity, and the potential which it generates relaxes to a stationary profile while the DF continues to mix, which is the analogue of Landau damping. Above μ_c the cloud collapses into a compact, self-bound clump which oscillates radially about the circular orbit and does not mix during the time span of our simulations. For clouds which are compact in angle, μ_c lies between 0.43 and 0.48, independently of the width of the cloud in action and of the external potential, whereas a cloud which is wide in angle requires a larger value. In particular, in the Kepler potential, which is the setting of Ref. [35], a cloud of a given shape only mixes when its mass lies below a critical value.

Particle methods have a long tradition in the numerical study of spherically symmetric collisionless systems, both in the Newtonian case [33, 34] and in the relativistic one, where they have been used to analyze the collapse of star clusters and the critical behavior at the threshold of black hole formation [39–43]; see Refs. [44–46] for introductions to the PIC method, Ref. [47] for a higher-order implementation in the context of plasma physics, and Ref. [48] for a convergence proof of a PIC scheme for the spherically symmetric Vlasov-Einstein system. Measuring the relaxation of the observables with such a method is delicate, because the signal has to be separated from the noise which is due to the finite number of computational particles. In these notes we address this problem by placing the particles on a regular lattice in action-angle variables, such that the sums over the particles are quadrature rules for the corresponding integrals over phase space. A loading of the particles which reduces the initial fluctuations (a quiet start) was already used for systems of spherical shells in Ref. [37]. Since the gas does not relax with respect to a fixed potential, we compute the observables with the action-angle variables of the instantaneous potential.

The remainder of these notes is organized as follows. In section II we formulate the Vlasov-Poisson system with an external potential in spherical symmetry and the reduced model with fixed angular momentum, and we introduce the action-angle variables and the observables used throughout these notes. In section III we review the mixing in the absence of self-gravity. Our numerical methods are described in section IV, and their validation is presented in section V. Section VI is devoted to the evolution of the gas cloud. Conclusions and an outlook, including the extension to a distribution of angular momenta, are given in section VII. Technical details regarding the numerical computation of the action-angle variables are included in Appendix A.

II. THE VLASOV-POISSON SYSTEM IN SPHERICAL SYMMETRY

In this section we formulate the Vlasov-Poisson system with an external potential and reduce it to spherical symmetry, first for a gas whose particles have arbitrary angular momenta and then for the reduced model in which all of them have the same magnitude of the angular momentum. Next, we introduce the dimensionless variables used in our numerical computations, the action-angle variables associated with a time-independent potential, in terms of which we characterize the steady states, and, finally, the observables on which our analysis of the mixing phenomenon is based.

We consider a collisionless gas consisting of identical particles which are subject to a time-independent external potential Φ_{ext} and to the gravitational potential Φ_{gas} generated by the gas itself. Since all the gas particles have the same mass, it is convenient to normalize the DF such that $f(t, x, p) d^3x\, d^3p$ is the mass of the gas contained in the volume element $d^3x\, d^3p$ of phase space, where from now on p denotes the momentum per unit mass.[2] With these conventions, the Vlasov-Poisson system consists of the Vlasov equation

$$\frac{\partial f}{\partial t} + p \cdot \nabla_x f - (\nabla_x \Phi) \cdot \nabla_p f = 0, \qquad \Phi := \Phi_{ext} + \Phi_{gas}, \tag{1}$$

[2] This is equivalent to working in units for which the mass of each gas particle is one, as in Ref. [14].

and the Poisson equation

$$\Delta\Phi_{gas} = 4\pi G\rho, \qquad \rho(t,x) := \int\limits_{\mathbb{R}^3} f(t,x,p)d^3p, \tag{2}$$

where G denotes Newton's constant and ρ is the mass density of the gas. To obtain a unique solution of Eq. (2) we require that $\Phi_{gas}(t,x) \to 0$ for $|x| \to \infty$.

For the external potential we choose Hénon's isochrone potential [1, 49, 50]

$$\Phi_{ext}(x) = -\frac{GM}{b+\sqrt{b^2+r^2}}, \qquad r := |x|, \tag{3}$$

which is generated by a spherical matter distribution of total mass M. The parameter $b \geq 0$ is a length scale which describes the transition from a harmonic potential (when $r \ll b$) to the potential of a point particle of mass M (when $r \gg b$). For $b = 0$, Eq. (3) reduces to the Kepler potential $\Phi_{ext} = -GM/r$, which is the external potential considered in Refs. [31, 35]. The reason for this choice is that in this potential the period of the radial motion depends only on the energy of the particle [49, 50], and that the corresponding action-angle variables are known in closed form [1]. When the self-gravity of the gas is neglected, this provides exact solutions against which our numerical methods can be validated.

A. Reduction to spherical symmetry

A spherically symmetric gas configuration is described by a DF which is invariant with respect to simultaneous rotations of x and p. Such a DF depends on (x,p) only through the radius r, the radial component $p_r := x \cdot p/r$ of the momentum and the magnitude $L := |x \wedge p|$ of the angular momentum, that is,

$$f(t,x,p) = \mathcal{F}(t,r,p_r,L). \tag{4}$$

In this case, the density ρ and the potential Φ_{gas} depend only on t and r, and the Vlasov equation (1) reduces to

$$\frac{\partial\mathcal{F}}{\partial t} + p_r\frac{\partial\mathcal{F}}{\partial r} - \frac{\partial V_L}{\partial r}\frac{\partial\mathcal{F}}{\partial p_r} = 0, \qquad V_L(t,r) := \Phi(t,r) + \frac{L^2}{2r^2}, \tag{5}$$

with V_L the effective potential for the radial motion. Since no derivative with respect to L appears in Eq. (5), the gas particles with a given value of L behave like a one-dimensional gas in the potential V_L. However, particles with different values of L interact with each other through the potential Φ_{gas}, which is determined by the spherically reduced Poisson equation

$$\frac{1}{r^2}\frac{\partial}{\partial r}\left(r^2\frac{\partial\Phi_{gas}}{\partial r}\right) = 4\pi G\rho, \qquad \rho(t,r) = \frac{2\pi}{r^2}\int\limits_0^\infty\int\limits_{-\infty}^\infty \mathcal{F}(t,r,p_r,L)\,dp_r\,LdL, \tag{6}$$

see for instance Ref. [17]. The solution of Eq. (6) which is regular at the center and vanishes at infinity is

$$\Phi_{gas}(t,r) = -4\pi G\int\limits_0^\infty \frac{\rho(t,s)}{\max\{r,s\}}s^2ds, \tag{7}$$

and its gradient is $\partial\Phi_{gas}/\partial r = Gm(t,r)/r^2$, where $m(t,r) := 4\pi\int_0^r \rho(t,s)s^2ds$ is the mass of the gas contained in the sphere of radius r at time t.

For the following, it is useful to note that when a spherically symmetric function is integrated over phase space, the volume element can be replaced with

$$d^3x\,d^3p = 8\pi^2 dr\,dp_r\,LdL. \tag{8}$$

In particular, the total mass and the total energy of the gas are given by

$$M_{gas} = 8\pi^2\int\limits_0^\infty\int\limits_{-\infty}^\infty\int\limits_0^\infty \mathcal{F}(t,r,p_r,L)\,dr\,dp_r\,LdL, \tag{9}$$

$$\mathcal{E}_{gas} = 8\pi^2\int\limits_0^\infty\int\limits_{-\infty}^\infty\int\limits_0^\infty \mathcal{F}(t,r,p_r,L)\left[\frac{p_r^2}{2} + \frac{L^2}{2r^2} + \Phi_{ext}(r) + \frac{1}{2}\Phi_{gas}(t,r)\right]dr\,dp_r\,LdL, \tag{10}$$

both of which are conserved in time. The conservation of $\mathcal{E}_{gas}$ provides a test for the accuracy of our numerical evolutions, see section V.

B. The reduced model with fixed angular momentum

A further simplification, which has been considered in Refs. [14, 17, 31], is obtained by assuming that all the gas particles have the same magnitude $L_0 > 0$ of the angular momentum.[3] In this case, the DF is of the form

$$\mathcal{F}(t, r, p_r, L) = \mathcal{F}_0(t, r, p_r)\delta(L - L_0), \tag{11}$$

and Eqs. (5,6) yield

$$\frac{\partial \mathcal{F}_0}{\partial t} + p_r \frac{\partial \mathcal{F}_0}{\partial r} - \frac{\partial V_{L_0}}{\partial r}\frac{\partial \mathcal{F}_0}{\partial p_r} = 0, \qquad \rho(t, r) = \frac{2\pi L_0}{r^2} \int_{-\infty}^{\infty} \mathcal{F}_0(t, r, p_r)\, dp_r, \tag{12}$$

where the potential Φ_{gas} is still given by Eq. (7). Likewise, Eqs. (9,10) hold with $\mathcal{F}$ replaced by $\mathcal{F}_0$, L by L_0, and the integral over L by the factor L_0. Equations (12,7) constitute a Vlasov-Poisson system on a two-dimensional phase space with coordinates (r, p_r). It describes a gas of concentric spherical shells which move in the effective potential V_{L_0}. The centrifugal term in this potential prevents the shells from reaching the center, and the gas configurations considered in these notes are confined to a region of the form $0 < r_{in} \leq r \leq r_{out}$. In spite of its simplicity, the reduced model retains the two ingredients which are relevant for these notes, namely the dependency of the frequency of the radial motion on the orbit, which is responsible for the mixing, and the self-gravity of the gas.

C. Dimensionless variables

In our numerical computations we use dimensionless variables, which are obtained by measuring masses in units of M, lengths in units of b and times in units of the dynamical time

$$t_{dyn} := \sqrt{\frac{b^3}{GM}}. \tag{13}$$

Accordingly, p_r is measured in units of $\sqrt{GM/b}$, L in units of $\sqrt{GMb}$, the potentials and the energy per unit mass in units of GM/b, and $\mathcal{F}$ in units of $M/(GMb)^{3/2}$. In terms of these variables, all the equations of this section retain their form with $G = M = b = 1$; in particular, the isochrone potential is $\Phi_{ext}(r) = -1/(1 + \sqrt{1 + r^2})$. The resulting system contains no free parameters, and the strength of the self-gravity is controlled by the ratio M_{gas}/M between the mass of the gas and the mass which generates the external potential. In the limit $M_{gas}/M \to 0$ the potential Φ_{gas} can be neglected, and one recovers the model analyzed in Ref. [14].

In the Kepler case $b = 0$ the external potential does not provide a length scale. However, in this case the system (5,6) is invariant with respect to the rescaling

$$(t, r, p_r, L, \mathcal{F}, \Phi) \mapsto \left(\alpha^{3/2} t, \alpha r, \alpha^{-1/2} p_r, \alpha^{1/2} L, \alpha^{-3/2}\mathcal{F}, \alpha^{-1}\Phi\right), \qquad \alpha > 0, \tag{14}$$

which leaves M_{gas} unchanged. Therefore, b can be replaced with an arbitrary length in the definitions above, such that $\Phi_{ext}(r) = -1/r$, and in the reduced model this freedom can be used to fix the value of L_0.

To give an idea of the physical time scale, we note that $M = 10^5 M_\odot$ and $b = 1$ pc, which are of the order of magnitude of the mass and of the core radius of a globular cluster [1], yield $t_{dyn} \approx 4.7 \times 10^4$ yr. As will be seen below, $2\pi t_{dyn}$ is a lower bound for the periods of the radial motion in the isochrone potential.

---

[3] Note that in Ref. [31] the symbol L refers to the square of the magnitude of the angular momentum.

D. Action-angle variables and steady states

Suppose that the total potential Φ is independent of time. In our applications, Φ is the external potential (3), the total potential Φ_{eq} of a self-consistent steady state (defined below), or the potential of an evolving gas at a fixed instant of time, see section II E. In all these cases $\Phi'(r) = Gm_{tot}(r)/r^2$, where $m_{tot}(r)$ denotes the total mass contained in the sphere of radius r, including the one which generates the external potential. The energy per unit mass

$$E := \frac{p_r^2}{2} + V_L(r) \tag{15}$$

is conserved along the trajectories of the gas particles, and following Refs. [1, 14] we introduce action-angle variables for their radial motion. Since the function $r \mapsto rm_{tot}(r)$ increases strictly monotonically from 0 to ∞, for each $L > 0$ the effective potential V_L has a unique critical point $r = r_{min}(L)$. It is determined by the equation $r^3\Phi'(r) = L^2$, it is a non-degenerate minimum of V_L, and it corresponds to a circular orbit with energy $E_{min}(L) := V_L(r_{min}(L))$. Since $V_L(r) \to \infty$ for $r \to 0$ and $V_L(r) \to 0$ for $r \to \infty$, the orbits with energies in the range $E_{min}(L) < E < 0$ are bound, and they oscillate between the turning points $r_1(E, L) < r_2(E, L)$, which are the two solutions of the equation $V_L(r) = E$.

As in Ref. [14] we define the area function

$$A(E, L) := \oint p_r dr = 2 \int_{r_1(E,L)}^{r_2(E,L)} \sqrt{2(E - V_L(r))}\, dr, \tag{16}$$

which represents the area enclosed by the orbit in the (r, p_r)-plane, and whose partial derivative with respect to E yields the period of the radial motion,

$$T(E, L) := \frac{\partial A}{\partial E}(E, L) = 2 \int_{r_1(E,L)}^{r_2(E,L)} \frac{dr}{\sqrt{2(E - V_L(r))}}. \tag{17}$$

The action and angle variables associated with the radial motion are

$$J := \frac{A(E, L)}{2\pi}, \qquad Q := \frac{2\pi}{T(E, L)} \int_{\gamma_r} \frac{dr}{p_r}, \tag{18}$$

where γ_r denotes the segment of the orbit in the (r, p_r)-plane which connects, in the clockwise direction, the inner turning point $(r_1(E, L), 0)$ with the given point (r, p_r). Hence, Q/Ω with $\Omega := 2\pi/T(E, L)$ is the time which has elapsed since the last passage of the particle through its pericenter; in particular, $Q = 0$ at the pericenter, $Q = \pi$ at the apocenter, and $0 < Q < \pi$ when $p_r > 0$. For each fixed value of L, the map $(r, p_r) \mapsto (Q, J)$ is symplectic, that is $dr \wedge dp_r = dQ \wedge dJ$, and it provides smooth coordinates on the set of bound orbits which are not circular. Since $T > 0$, the first relation in Eq. (18) can be inverted for each L, which yields the energy as a function $E(J, L)$ of the action variable and the angular momentum, and the frequency of the radial motion is

$$\Omega(J, L) = \frac{\partial E}{\partial J}(J, L) = \frac{2\pi}{T(E, L)}. \tag{19}$$

[(Erik) We denote the orbital frequency by Ω (it is called ω in Ref. [14]) and reserve ω for the frequency of the collective oscillation.] In terms of the variables (Q, J, L), the volume element (8) is $8\pi^2 dQ\, dJ\, LdL$, and the Vlasov equation (5) for a time-independent potential assumes the simple form

$$\frac{\partial \mathcal{F}}{\partial t} + \Omega(J, L)\frac{\partial \mathcal{F}}{\partial Q} = 0. \tag{20}$$

In particular, the time-independent solutions of Eq. (20) are the DFs which do not depend on the angle variable. Since $E(J, L)$ is a strictly increasing function of J, they are of the form

$$\mathcal{F}(r, p_r, L) = \psi(E, L), \tag{21}$$

with a nonnegative function ψ. Such a DF is a solution of the Vlasov-Poisson system if the potential Φ_{gas}, which enters the definition of E through V_L, is the one generated by the density of the DF through Eqs. (6,7). This is a nonlinear

problem for Φ_{gas}. A solution $(\mathcal{F}_{eq}, \Phi_{eq})$ of this problem, with Φ_{eq} the total potential and ρ_{eq} the corresponding density, will be referred to as a self-consistent steady state.

For the external potential (3) the area function can be computed explicitly, see Refs. [1, 14]. In terms of the dimensionless variables introduced in the previous subsection, one finds

$$A(E,L) = 2\pi \left[\frac{1}{\sqrt{-2E}} - c(L) \right], \qquad E_{min}(L) = -\frac{1}{2c(L)^2}, \tag{22}$$

with $c(L) := \left(L + \sqrt{L^2+4}\right)/2$ for the isochrone potential and $c(L) := L$ for the Kepler potential. This yields

$$E(J,L) = -\frac{1}{2\left[J + c(L)\right]^2}, \qquad \Omega(J,L) = \frac{1}{\left[J + c(L)\right]^3} = (-2E)^{3/2}. \tag{23}$$

Therefore, the frequency depends only on the energy, it decreases monotonically when J or L increase, and for each fixed L its maximum is attained in the limit $J \to 0$ of circular orbits. For the isochrone potential, $c(L) > 1$ implies that $\Omega < 1$, such that the periods of the radial motion are larger than 2π. An explicit expression for the angle variable Q in terms of (r, p_r, L) can be found in section 3.5.2 of Ref. [1]; its inversion requires the solution of an equation of Kepler's type. In contrast to this, when the self-gravity of the gas is taken into account the potential is only known numerically, and in this case we compute the action-angle variables by numerical quadrature of Eqs. (16,17,18), see Appendix A.

E. Observables

As in Ref. [14], we describe the macroscopic state of the gas through observables of the form

$$N_\varphi(t) := \int f(t,x,p)\varphi(x,p)\, d^3x\, d^3p, \tag{24}$$

where φ is a given test function on phase space. In these notes we consider two classes of such observables.

The first one refers to the action-angle variables (Q, J) belonging to a time-independent potential. When the self-gravity of the gas is neglected, this is the external potential. When it is taken into account and the gas is far from equilibrium, there is no fixed potential with respect to which the gas relaxes, and we use the action-angle variables of the instantaneous potential: at each time t, (Q, J) are the action-angle variables which belong to the potential $r \mapsto \Phi(t, r)$, regarded as a time-independent potential. For a spherically symmetric configuration and a test function of the form $\varphi = a(Q)B(J, L)$, with real-valued functions a and B, Eqs. (8,24) and the expansion of a in a Fourier series yield

$$N_\varphi(t) = \sum_{n\in\mathbb{Z}} \hat{a}_n^*\, h_n(t), \qquad \hat{a}_n := \frac{1}{2\pi} \int_0^{2\pi} a(Q) e^{-inQ} dQ, \tag{25}$$

with the functions

$$h_n(t) := 8\pi^2 \int_0^\infty \int_0^\infty \int_0^{2\pi} \mathcal{F}(t,Q,J,L) B(J,L) e^{-inQ}\, dQ\, dJ\, LdL, \qquad n \in \mathbb{Z}. \tag{26}$$

In the reduced model, B depends only on J and Eq. (26) holds with $\mathcal{F}$ replaced by $\mathcal{F}_0$ and the integral over L by the factor L_0. Since $\mathcal{F}$ is real, $h_{-n} = h_n^*$, such that it is sufficient to consider $n \geq 0$. For a steady state h_n vanishes for all $n \neq 0$, whereas h_0 is equal to the mass of the gas weighted with the function B. Therefore, the functions $h_n(t)$ with $n \neq 0$ measure the departure of the gas configuration from a steady state, and their decay in time is the signature of the relaxation process we are interested in. We stress that this property relies on the use of the action-angle variables which belong to the total potential. If those of the external potential are used instead, a gas whose self-gravity is not negligible gives rise to nonvanishing values of h_n with $n \neq 0$ even after it has relaxed, as we show in section V. When the action-angle variables of the instantaneous potential are used, the test function depends on the DF through $\Phi(t, \cdot)$, such that h_n is no longer a linear functional of the DF. On the other hand, h_n with $n \neq 0$ then vanishes for any steady state, including a steady state which the gas may approach at late times and which is not known in advance.

The second class of observables consists of the deviation of the gravitational potential from its average $\langle\Phi\rangle(r)$ over a late time interval, which is specified in section VI,

$$\delta\Phi(t,r) := \Phi(t,r) - \langle\Phi\rangle(r) = -4\pi G \int_0^\infty \frac{\rho(t,s) - \langle\rho\rangle(s)}{\max\{r,s\}} s^2 ds, \tag{27}$$

where we have used Eq. (7) and $\langle\rho\rangle$ denotes the average of the density over the same interval. For each fixed r, $\Phi_{gas}(t,r)$ is the observable (24) with the test function $\varphi(x,p) = -G/\max\{r,|x|\}$. In contrast to the functions $h_n(t)$, the quantity $\delta\Phi$ does not refer to any particular coordinates on phase space. The results in Refs. [31, 32] refer to the decay of the force field $-\partial\delta\Phi/\partial r$ which is generated by it, or of the time derivative of this field. To measure its magnitude we use the root mean square

$$\|\delta\Phi(t)\| := \left(\frac{1}{r_b - r_a} \int_{r_a}^{r_b} \delta\Phi(t,r)^2 dr \right)^{1/2} \tag{28}$$

over an interval $[r_a, r_b]$ which covers the region occupied by the gas.

III. PHASE SPACE MIXING IN THE ABSENCE OF SELF-GRAVITY

In this section we review the mixing phenomenon in the absence of self-gravity, for an arbitrary initial DF, and discuss the rate at which the observables introduced in the previous section decay. The derivations are formal; for rigorous statements we refer the reader to Refs. [14, 16].

When the self-gravity of the gas is neglected, $\Phi = \Phi_{ext}$ is independent of time, and the solution of Eq. (20) is

$$\mathcal{F}(t,Q,J,L) = F(Q - \Omega(J,L)t, J, L), \tag{29}$$

where the function F, which is 2π-periodic in its first argument, describes the initial DF. No assumption on F is needed here: it need not be close to a steady state. Expanding F in a Fourier series with respect to the angle variable,

$$F(Q,J,L) = \sum_{n\in\mathbb{Z}} \hat{F}_n(J,L) e^{inQ}, \qquad \hat{F}_n(J,L) := \frac{1}{2\pi} \int_0^{2\pi} F(Q,J,L) e^{-inQ} dQ, \tag{30}$$

and introducing Eq. (29) into Eq. (26) one obtains

$$h_n(t) = 16\pi^3 \int_0^\infty \int_0^\infty \hat{F}_n(J,L) B(J,L) e^{-in\Omega(J,L)t}\, dJ\, L dL. \tag{31}$$

Hence, h_0 is constant in time, whereas for $n \neq 0$ the function $h_n(t)$ is a superposition of oscillations whose frequencies $n\Omega(J,L)$ vary from one orbit to the other. Provided that the gradient of Ω with respect to (J,L) is different from zero for almost all (J,L) (in the reduced model, provided that $\partial\Omega/\partial J \neq 0$ for almost all J), it follows from the Riemann-Lebesgue lemma and its generalizations (cf. Refs. [14, 16]) that $h_n(t) \to 0$ for $t \to \infty$ and all $n \neq 0$. As a consequence, the observables (25) converge to the ones which belong to the angle-averaged DF $\hat{F}_0(J,L)$, which is a steady state according to the remark below Eq. (20). This is the phase space mixing property, and $\hat{F}_0$ is the final state of the gas. For the isochrone and the Kepler potentials, Eq. (23) gives $\partial\Omega/\partial J = -3[J + c(L)]^{-4} < 0$, such that the non-degeneracy condition is satisfied.

The rate at which $h_n(t)$ decays is not universal. It depends on the regularity of the function $\hat{F}_n B$ and on its behavior at the boundary of its support. If this function is smooth and vanishes together with all its derivatives at the boundary, repeated integration by parts in Eq. (31) shows that $h_n(t)$ decays faster than any inverse power of t. A different situation arises in the reduced model when the DF has its support in an interval $0 \leq J \leq J_{max}$ and vanishes like a power at the outer edge, such that

$$\hat{F}_n(J) B(J) = (J_{max} - J)^k g(J) \tag{32}$$

with an exponent $k > 0$ and a smooth function g satisfying $g(J_{max}) \neq 0$. Suppose that $\Omega'(J) < 0$ for all $0 < J \leq J_{max}$, as is the case in all the examples of these notes, and that g vanishes sufficiently fast for $J \to 0$. Then, the behavior

of the integral in Eq. (31) for large times is determined by the endpoint $J = J_{max}$ (see, for instance, chapter 6 in Ref. [51]), and for $n \geq 1$ one finds

$$h_n(t) \simeq 16\pi^3 L_0\, \Gamma(k+1)\, g(J_{max}) \frac{e^{-in\Omega_{min}t}}{[in|\Omega'(J_{max})|t]^{k+1}}, \qquad t \to \infty, \tag{33}$$

where $\Omega_{min} := \Omega(J_{max})$ is the frequency of the outermost orbit and Γ denotes the Gamma function. Therefore, an edge at which the DF vanishes with the exponent k gives rise to an algebraic tail which is proportional to $t^{-(k+1)}$ and which oscillates with the frequency $n\Omega_{min}$: the less regular the DF is at the edge, the slower is the decay. The general rule behind this result is that a singularity of the form $(\omega - \omega_0)^a$ of the Laplace transform of an observable (with a logarithmic factor when a is an integer) gives rise to a tail which is proportional to $t^{-(a+1)}$ and oscillates with the frequency ω_0; the singularities which occur for systems with one and with two action variables have been classified in Refs. [52, 53]. Decay estimates for the mixing in the potential of a steady state whose DF vanishes like a power at the edge of its support have been proven in Ref. [23].

When the angular momenta of the gas particles are distributed, the integration over L in Eq. (31) provides an additional source of decay, since Ω depends on L as well. For the isochrone and the Kepler potentials, Eq. (23) yields $\partial\Omega/\partial L = c'(L)\,\partial\Omega/\partial J$ with $1/2 < c'(L) \leq 1$, such that a dispersion in L is almost as efficient for the mixing as a dispersion of the same magnitude in J. In these notes we restrict ourselves to the reduced model.

When the self-gravity of the gas is taken into account, the potential depends on time, and there is no longer a fixed set of action-angle variables in which the motion of the gas particles is trivial. If the initial DF is close to a self-consistent steady state, the deviation from that state can be described by the equations linearized about it, as in the works on Landau damping and oscillating modes cited in section I. The initial data of these notes are far from any steady state, and their evolution has to be followed with the nonlinear Vlasov-Poisson system, which we solve with the PIC method described in the next section. In this case we compute the observables (26) with the action-angle variables of the instantaneous potential, see section II E.

IV. NUMERICAL METHODS

In this section we describe the PIC method which we use to solve the nonlinear Vlasov-Poisson system, and the representation of the initial data by the computational particles.

A. Particle-in-cell method

In the PIC method [44, 45] the DF is represented by a finite number N of computational particles, each of which stands for a large number of gas particles. In spherical symmetry a computational particle is a spherical shell, which is characterized by its radius $r_j(t)$, its radial momentum $p_j(t)$, its angular momentum L_j and its mass m_j, $j = 1, 2, \ldots, N$, the last two quantities being constant in time. This corresponds to the approximation

$$\mathcal{F}(t, r, p_r, L) \approx \sum_{j=1}^{N} \frac{m_j}{8\pi^2 L_j}\, \delta(r - r_j(t))\, \delta(p_r - p_j(t))\, \delta(L - L_j) \tag{34}$$

of the DF, where the factor $8\pi^2 L_j$ is due to the volume element (8). Introducing Eq. (34) into the Vlasov equation (5) yields the equations of motion

$$\frac{dr_j}{dt} = p_j, \qquad \frac{dp_j}{dt} = \frac{L_j^2}{r_j^3} - \frac{\partial\Phi_{ext}}{\partial r}(r_j) - \frac{\partial\Phi_{gas}}{\partial r}(t, r_j) \tag{35}$$

for the computational particles; in the reduced model $L_j = L_0$ for all j. The first two terms on the right-hand side of the second equation are evaluated analytically, whereas the last one requires the solution of the Poisson equation (6). To this purpose we introduce the uniform grid $R_i := (i - 1/2)\Delta r$, $i = 1, 2, \ldots, N_r$, which is staggered with respect to the center, and proceed in four steps.

(i) The mass of each computational particle is assigned to its two neighboring grid points with the linear weights of the cloud-in-cell scheme [44, 45], such that the grid point R_i receives the mass

$$\mu_i := \sum_{j=1}^{N} m_j W\left(\frac{R_i - r_j}{\Delta r}\right), \qquad W(y) := \max\{1 - |y|, 0\}. \tag{36}$$

(ii) The density is approximated by the continuous, piecewise linear function $\rho(t,r)$ which assumes the values

$$\rho_i := \frac{\mu_i}{4\pi\Delta r\,(R_i^2 + \Delta r^2/6)} \tag{37}$$

at the grid points and which is constant for $0 \le r \le R_1$. The denominator in Eq. (37) is the integral of the function $4\pi r^2 W((r-R_i)/\Delta r)$ over r, which guarantees that the mass of the reconstructed density coincides with the total mass of the computational particles.

(iii) The mass function $m(t,r)$ and the potential are obtained by integrating the equations $\partial m/\partial r = 4\pi r^2\rho$ and $\partial\Phi_{gas}/\partial r = Gm/r^2$ outwards from the center. Since ρ is linear on each interval $[R_{i-1}, R_i]$, both integrals are computed in closed form, such that no further discretization error is introduced. The additive constant in Φ_{gas} is fixed by requiring that $\Phi_{gas} + r\,\partial\Phi_{gas}/\partial r = 0$ at the outermost grid point, which matches the potential to the exterior solution $-GM_{gas}/r$.

(iv) The potential and its gradient are interpolated from the grid points to the positions of the computational particles with the same weights as in Eq. (36).

For particles which lie within the distance Δr from the center, the weights in steps (i) and (iv) are complemented by those of their mirror images at $-r_j$, in accordance with the regularity of the density at the center, and beyond the outermost grid point the exterior solution is used. Neither of these situations plays a role for the configurations considered in these notes, in which the gas occupies a shell that lies well inside the grid; the computational particles which leave the grid for the widest initial data of section VI carry less than 10^{-11} of the mass of the gas. For the Kepler potential the code uses the potential of a homogeneous sphere of mass M and unit radius, which coincides with the Kepler potential for $r \ge 1$. In all the simulations of these notes with this potential the gas remains in the region $r > 1.7$, such that the two potentials are equivalent.

Note that the potential which acts on a computational particle includes the contribution of the particle itself. In PIC schemes based on Cartesian coordinates the use of the same weights for the assignment and for the interpolation guarantees that a particle exerts no force on itself [44, 45]. In spherical symmetry this is not the case, and it should not be: a thin shell of mass m and radius r is subject to the mean value of the gravitational fields on its two sides, that is, to the force $-Gm/(2r^2)$ per unit mass. This is the self-gravity of the shell; the scheme reproduces it up to a correction of the order $\Delta r/r$, and we do not subtract it. In fact, we have verified that subtracting it reduces the order of convergence with respect to $1/N$ from two to one in the collapse of a cold homogeneous sphere, for which the exact solution is known.

Equations (35) are integrated in time with the fourth-order symplectic method of Yoshida [54], which is a composition of three steps of the second-order leapfrog method with step sizes $w_1\Delta t$, $w_0\Delta t$ and $w_1\Delta t$, where $w_1 := 1/(2-2^{1/3})$ and $w_0 := 1 - 2w_1$. A leapfrog step with step size h consists of the update $p_j \mapsto p_j + hf_j/2$ of the momenta, with f_j the right-hand side of the second equation in (35), followed by the update $r_j \mapsto r_j + hp_j$ of the radii, the computation of the potential for the new radii, and a second update of the momenta with the new values of f_j. Hence, the Poisson equation is solved three times per time step. We use a symplectic method because, for the motion in a given potential, it does not introduce a secular drift of the energy and of the action variables of the particles, which would otherwise limit the accuracy of the observables at late times. For the same reason we keep the time step Δt fixed.

At selected times we compute the total energy (10),

$$\mathcal{E}_{gas} \approx \sum_{j=1}^{N} m_j \left[\frac{p_j^2}{2} + \frac{L_j^2}{2r_j^2} + \Phi_{ext}(r_j) + \frac{1}{2}\Phi_{gas}(t,r_j)\right], \tag{38}$$

the potential Φ_{gas} on the grid, from which $\delta\Phi$ is obtained according to Eq. (27), and the functions

$$h_n(t) \approx \sum_{j=1}^{N} m_j B(J_j, L_j) e^{-inQ_j}, \tag{39}$$

which follow from Eqs. (26,34). Here (Q_j, J_j) denote the action-angle variables of the particle j at the time t with respect to the potential specified in section II E. They are computed from (r_j, p_j) with the method described in Appendix A; for the instantaneous potential, Φ_{gas} is the potential computed by the PIC code on its grid at the time t. In the tests of section V we choose the weight function

$$B(J) = J^2 \exp\left[-\frac{(J-J_1)^2}{\sigma_1^2}\right], \tag{40}$$

with parameters J_1 and σ_1 which are specified below. The factor J^2 makes the function $B(J)e^{-inQ}$ continuous at the circular orbit for $|n| \leq 4$. For the evolution of the gas cloud in section VI we choose $B = 1$, such that h_n/h_0 is the average of e^{-inQ} over the mass of the gas.

The simulations of these notes use an implementation of this method for the reduced model. Unless otherwise stated, they use $\Delta r = 0.1$, a grid which extends to $r = 20$, a time step $\Delta t = 0.05$ or 0.1, and $N = 10^4$ computational particles. [(Erik) To be decided: whether the codes and the scripts which reproduce the results are made public, in which case the repository should be cited here.]

B. Initial data

The initial data considered in these notes are given as functions of the action-angle variables (Q, J) of the external potential $\Phi_0 := \Phi_{ext}$.

If the radii and momenta of the computational particles were drawn at random from the initial DF, the observables would be affected by fluctuations of the order of $N^{-1/2}$. For the numbers of particles used in this work, these fluctuations would exceed the values of $|h_n|/h_0$ which are reached during the mixing of the gas. For this reason we place the computational particles on a regular lattice in the variables (Q, J),

$$J_a := \left(a - \frac{1}{2}\right)\frac{J_{max}}{N_J}, \quad a = 1, \ldots, N_J, \qquad Q_b := \left(b - \frac{1}{2}\right)\frac{2\pi}{N_Q}, \quad b = 1, \ldots, N_Q, \tag{41}$$

where $[0, J_{max}]$ is the range of the action variable which is covered by the DF, such that $N = N_J N_Q$. The radii and momenta of the particles are obtained by inverting the map $(r, p_r) \mapsto (Q, J)$ which belongs to Φ_0, which only requires the solution of an equation of Kepler's type (see below), and the particle at the lattice point (Q_b, J_a) is given a mass m_{ab} which is proportional to the value of the initial DF at this point, with a constant of proportionality which is determined by the total mass M_{gas}. Since the map $(r, p_r) \mapsto (Q, J)$ is symplectic, all the cells of the lattice have the same volume in phase space. Hence, the sums over the computational particles in Eqs. (38,39) are midpoint rules for the corresponding integrals over phase space, and they are free of the fluctuations of a random sample. In plasma physics, a loading of the particles which avoids these fluctuations is known as a quiet start [44]. The method is only reliable over a finite time span: in the absence of self-gravity the rows of the lattice produce a spurious recurrence of the functions h_n, which is quantified in section V, and in its presence the regularity of the lattice is eventually lost due to the discreteness of the particles.

As initial datum we consider the gas cloud

$$\mathcal{F}_0(0, Q, J) = \alpha \exp\left[-\frac{\sin^2(Q/2)}{\sigma_Q^2}\right] J^2 \exp\left(-\frac{J^2}{\sigma_J^2}\right), \tag{42}$$

where (Q, J) are the action-angle variables of the external potential and the constant $\alpha > 0$ is determined by the mass of the gas. For $\sigma_Q \ll 1$ the gas particles are concentrated near the pericenters $Q = 0$ of their orbits, and their action variables are distributed about $J = \sigma_J$, see Fig. 1. The factor J^2 makes the DF smooth at the circular orbit $J = 0$, where the angle variable is not defined. In the absence of self-gravity, the mixing of this cloud is the test problem of section V A, and in section VI we analyze how its evolution changes when the mass of the gas is increased. We truncate the DF at $J_{max} = 6\sigma_J$, where it has decayed to about 2×10^{-14} of its maximum. Unless otherwise stated, $\sigma_Q = \sigma_J = 0.1$ and $N_J \times N_Q = 400 \times 25$. For the isochrone and the Kepler potentials the inversion of the map $(r, p_r) \mapsto (Q, J)$ only requires the solution of an equation of Kepler's type (see section II D), which we solve with Newton's method. In the Kepler case, for instance,

$$r = a(1 - e\cos\eta), \qquad p_r = \frac{\sqrt{a}\, e \sin\eta}{r}, \qquad Q = \eta - e\sin\eta, \tag{43}$$

with $a := (J + L_0)^2$ and $e := (1 - L_0^2/a)^{1/2}$. When the self-gravity of the gas is taken into account, the observables are computed with the action-angle variables of the instantaneous potential, see section II E.

V. CODE VALIDATION

In this section we validate the numerical methods described in the previous section. We start with the mixing of the gas cloud (42) in the absence of self-gravity, for which the exact solution is known. Next, we include the self-gravity of the cloud and show that the functions $h_n(t)$ need to be computed with the action-angle variables which belong to the total potential. Finally, we test the computation of the gravitational field in the collapse of a cold homogeneous sphere, and we report the conservation of the total energy in the simulations of section VI.

A. Mixing in the absence of self-gravity

When the last term in the second equation in (35) is omitted, the computational particles move independently from each other in the external potential, and the functions $h_n(t)$ are given by Eq. (31). This provides a test for the time integration, for the representation of the DF by the lattice (41) and for the computation of the observables. We consider the reduced model with $L_0 = 2$ in the isochrone potential and the gas cloud (42) with $\sigma_Q = \sigma_J = 0.1$ (see Fig. 1), and we choose the weight function $B(J) = J^2 \exp(-J^2/\sigma_J^2)$, that is, Eq. (40) with $J_1 = 0$ and $\sigma_1 = \sigma_J$.

The dependency of the DF (42) on the angle variable has been chosen such that it is smooth and 2π-periodic and such that its Fourier coefficients (30) are known explicitly. In fact, the identity $\sin^2(Q/2) = (1 - \cos Q)/2$ yields

$$\exp\left[-\frac{\sin^2(Q/2)}{\sigma_Q^2}\right] = e^{-z}e^{z\cos Q} = e^{-z}\sum_{n\in\mathbb{Z}} I_n(z)e^{inQ}, \qquad z := \frac{1}{2\sigma_Q^2}, \tag{44}$$

where I_n denotes the modified Bessel function of the first kind of order n, which satisfies $I_{-n} = I_n$. The second equality in Eq. (44) is the generating function of these functions, see Eq. (10.35.2) in Ref. [55]; equivalently, $I_n(z)$ is the n-th Fourier coefficient of the function $e^{z\cos Q}$ (Eq. (10.32.3) in Ref. [55]). Consequently, the Fourier coefficients of the DF (42) with respect to the angle variable are

$$\hat{F}_n(J) = \alpha\, e^{-z} I_n(z)\, J^2 \exp\left(-\frac{J^2}{\sigma_J^2}\right). \tag{45}$$

For $\sigma_Q = 0.1$ one has $z = 50$, and in this case $I_n(z)/I_0(z)$ differs from $e^{-n^2/(2z)}$ by less than one percent for $|n| \le 16$. Hence, the narrower the pulse is in the angle variable, the more harmonics it contains: here the amplitudes of those with $|n| \le 2\sqrt{z} \approx 14$ are larger than $e^{-2} \approx 0.14$ times the one of the harmonic $n = 0$. It follows from Eqs. (31,45) that $|h_n(0)|/h_0 = I_n(z)/I_0(z)$, which is equal to 0.990, 0.960, 0.913 and 0.851 for $n = 1, 2, 3, 4$, and that $h_n(t)$ is given by an integral over J, which we compute with the trapezoidal rule on 2×10^5 subintervals of $0 \le J \le 6\sigma_J$.




FIG. 1: Mixing of the initial datum (42) in the isochrone potential when self-gravity is neglected. Shown are the computational particles in the (r, p_r)-plane (upper panels) and in the plane of the action-angle variables (lower panels) at four different times, for a lattice with $N_J \times N_Q = 800 \times 64$ points. The color indicates the mass of each particle relative to the largest one, and only particles whose mass exceeds 10^{-3} of the latter are shown.

Table I shows the error

$$e_n := \frac{1}{h_0} \max_{0\le t\le 2000} \left|h_n(t) - h_n^{exact}(t)\right| \tag{46}$$

for different lattices and time steps. For $N_J \times N_Q = 400 \times 64$ and $\Delta t = 0.1$ it does not exceed 6×10^{-9} for $n \le 4$. In this case the error is dominated by the time integration. In fact, when the Yoshida integrator is replaced with the

N_J	N_Q	Δt	e_1	e_2	e_3	e_4
50	64	0.1	4.1×10^{-6}	1.6×10^{-4}	0.15	0.72
100	64	0.1	3.3×10^{-8}	1.2×10^{-7}	5.1×10^{-7}	3.9×10^{-6}
200	64	0.1	5.3×10^{-9}	5.4×10^{-9}	5.9×10^{-9}	5.5×10^{-9}
400	64	0.1	5.3×10^{-9}	5.5×10^{-9}	6.0×10^{-9}	5.5×10^{-9}
400	16	0.1	9.2×10^{-3}	3.7×10^{-2}	8.2×10^{-2}	0.14
400	32	0.1	1.7×10^{-5}	7.3×10^{-5}	1.8×10^{-4}	3.7×10^{-4}
400	64	0.2	8.6×10^{-8}	8.8×10^{-8}	9.6×10^{-8}	8.8×10^{-8}
400	64	0.05	3.3×10^{-10}	3.4×10^{-10}	3.7×10^{-10}	3.4×10^{-10}
400	64	0.025	2.1×10^{-11}	2.0×10^{-11}	2.2×10^{-11}	2.7×10^{-11}
400	64	exact	6.6×10^{-12}	1.3×10^{-11}	2.0×10^{-11}	2.7×10^{-11}

TABLE I: Error e_n defined in Eq. (46) for the mixing of the initial datum (42) in the absence of self-gravity, for different lattices and time steps. In the last row the Yoshida integrator is replaced with the exact advance of the angle variables.

exact advance $Q_j \mapsto Q_j + \Omega(J_j)\Delta t$ of the angle variables of the particles, the error drops to less than 3×10^{-11}, and the difference with respect to this solution decreases by the factors 16.0, 16.0 and 15.4 each time Δt is halved between 0.2 and 0.025 (for $n = 1$), as expected for a fourth-order method.

The remaining rows of Table I show the effect of the lattice. Since the sum (39) is a midpoint rule for an integral whose integrand is smooth and, with respect to Q, periodic, the error decreases rapidly when N_J or N_Q increase, until it reaches the one of the time integration.

With respect to the angle variable the error of the lattice can be computed explicitly. On a lattice with N_Q points per row the harmonics $n + mN_Q$, $m \in \mathbb{Z}$, cannot be distinguished from the harmonic n, such that the sum over the points Q_b yields $\sum_m (-1)^m \hat{F}_{n+mN_Q}$ instead of $\hat{F}_n$, where the signs are due to the fact that the Q_b are the midpoints of the subintervals. Since the transport does not mix different harmonics, this factor is independent of time. Taking into account that the masses of the computational particles are normalized such that their sum is equal to M_{gas}, Eq. (45) gives

$$e_n = \left| \frac{\sum_m (-1)^m I_{n+mN_Q}(z)}{\sum_m (-1)^m I_{mN_Q}(z)} - \frac{I_n(z)}{I_0(z)} \right|, \tag{47}$$

where the sums extend over all $m \in \mathbb{Z}$. This expression reproduces the values in the rows with $N_Q = 16$ and $N_Q = 32$ of Table I to nine significant digits, and for $N_Q = 64$ it is smaller than 5×10^{-15}. Hence, the number of points per row has to be a few times larger than the number $2\sqrt{z}$ of harmonics contained in the DF.

With respect to the action variable the situation is different, since for a given lattice the sum is only reliable over a finite time span. The reason is that the phases $n\Omega(J_a)t$ of two neighboring rows differ by $n|\Omega'(J)|\Delta J\, t$, with ΔJ the spacing between the rows. When this difference reaches 2π the rows are in phase again, such that h_n exhibits a spurious recurrence [44] at the time

$$t_{rec} = \frac{2\pi}{n|\Omega'(J)|\Delta J}. \tag{48}$$

In the present example, $\Delta J = 0.6/N_J$ and $|\Omega'(J)| \le 3/c(L_0)^4 \approx 0.088$, which gives $t_{rec} \approx 120\, N_J/n$. For $N_J = 50$ this is 1980 and 1480 for $n = 3$ and $n = 4$, respectively, which explains the large values of e_3 and e_4 in the first row of the table. For the lattices which are used in the next subsection and in section VI the recurrence time for $n = 1$ exceeds the duration of the simulations: computed with the frequencies of the external potential, it is 4.7×10^4 for the one of the next subsection and at least 2.2×10^4 for those of section VI. For the higher harmonics it is shorter by the factor $1/n$.

Figure 2 shows the result of a simulation with $N_J \times N_Q = 800\times64$ and $\Delta t = 0.1$ which extends to $t = 10^4$. During this time $|h_1|$ and $|h_4|$ decay by seven and ten orders of magnitude, respectively, and the numerical solution reproduces the exact one over the whole range: for $n \le 4$ the difference does not exceed $6\times10^{-9}h_0$ at any time, and it is smaller than $2\times10^{-10}h_0$ for $t \ge 5000$. At late times $|h_n(t)|$ is proportional to t^{-5} (a fit over $5000 \le t \le 10^4$ gives the exponents -5.03 and -5.01 for $n = 1$ and $n = 2$). This is the analogue of the algebraic tail (33) for the endpoint $J = 0$ of the support, at which the function $\hat{F}_n B$ vanishes like J^4. In the same simulation h_0, which is constant for the exact solution, varies by 1.1×10^{-9}, and the total energy by 7×10^{-11}, in relative terms.




FIG. 2: Left panel: the functions $|h_n(t)|/h_0$, $n = 1, 2, 3, 4$, for the mixing of the initial datum (42) in the absence of self-gravity, obtained from the PIC code with $N_J \times N_Q = 800 \times 64$ and $\Delta t = 0.1$ (solid lines) and from the exact solution (31) (circles). Right panel: error of the time integration, measured through the difference with the solution in which the angle variables are advanced exactly, as a function of the time step. The dashed line is proportional to Δt^4.

B. Action-angle variables of the total potential

As emphasized in section II E, the functions $h_n(t)$ with $n \neq 0$ measure the departure of the gas configuration from a steady state only if they are computed with the action-angle variables which belong to the total potential. To illustrate this point we have repeated the simulation of the previous subsection including the self-gravity of the gas, for $M_{gas}/M = 10^{-4}$, 10^{-3} and 10^{-2}, with $N_J \times N_Q = 400 \times 25$, $\Delta r = 0.1$, $\Delta t = 0.1$ and up to $t = 2 \times 10^4$, using the weight function (40) with $J_1 = \sigma_1 = 0.1$. The cloud is far from a steady state, and the gas undergoes mixing while its potential Φ_{gas} readjusts: for $M_{gas}/M = 10^{-3}$ it changes by up to 40 percent of its largest value between $t = 0$ and $t = 2000$, and afterwards it is stationary up to fluctuations whose root mean square is smaller than 10^{-3} of this value.




FIG. 3: Dependency of the function h_1 on the action-angle variables which are used to compute it. (a) $|h_1|/h_0$ for the mixing of the initial datum (42) including self-gravity, computed with the action-angle variables of the external potential, for different masses of the gas. (b) Real part of h_1/h_0 for $M_{gas}/M = 10^{-3}$, computed with the action-angle variables of the external potential and with those of the potential $\Phi_{ext} + \langle\Phi_{gas}\rangle$, where $\langle\Phi_{gas}\rangle$ is the average of Φ_{gas} over $t \geq 2000$. (c) Relative variation of the action variable of a single computational particle for the same mass, for both sets of variables; here $\langle\Phi_{gas}\rangle$ is the average over $1500 \leq t \leq 2000$, and $\langle J\rangle$ denotes the mean value over this interval.

Figure 3(a) shows the function $|h_1(t)|/h_0$ which is obtained with the action-angle variables of the external potential. When self-gravity is neglected it decays according to the law t^{-5} found in the previous subsection, until it reaches the level of the numerical errors, which is of the order of 10^{-9}, at $t \approx 10^4$. In contrast to this, when self-gravity is taken into account it settles down to a plateau: the magnitude of the time average of h_1/h_0 over the interval $2000 \leq t \leq 2 \times 10^4$ is 2.2×10^{-4}, 2.2×10^{-3} and 1.5×10^{-2} for the three values of M_{gas}/M, whereas it is 9×10^{-9} in the absence of self-gravity. For $M_{gas}/M = 10^{-3}$ the same value, 2.16×10^{-3}, is obtained with $N_J \times N_Q = 800 \times 25$ and 400×50. Such a plateau could be mistaken for a collective mode which is not damped.

However, the plateau disappears when the same particle data are analyzed with the action-angle variables of the potential $\Phi_{ext}+\langle\Phi_{gas}\rangle$, where $\langle\Phi_{gas}\rangle$ denotes the time average of the potential of the gas after it has become stationary. This is shown in Fig. 3(b) for $M_{gas}/M = 10^{-3}$: with these variables h_1/h_0 fluctuates about zero instead of about the value 2.16×10^{-3}, and the magnitude of its time average over $2000 \leq t \leq 2 \times 10^4$ is reduced to 5.4×10^{-7}, which is 4000 times smaller. The action-angle variables of the instantaneous potential, which we use for the initial data of section VI, give the same result: with them the magnitude of the time average is 1.5×10^{-6}. Therefore, the plateau is due to the choice of the coordinates and not to the dynamics. The reason is that the action variable of the external potential is not conserved along the trajectories of the gas particles in the total potential. As shown in Fig. 3(c), for a given particle it oscillates with the period of the radial motion; the relative difference between its largest and its smallest value is 5.6×10^{-3} for the particle shown, which has $J = 0.10$, and 4.4×10^{-3} for a particle with $J = 0.30$. In contrast to this, the action variable which belongs to the total potential is constant to within a relative variation of 2×10^{-5}. Consequently, a DF which depends only on the action variable of the total potential is not independent of the angle variable of the external one, the difference between both sets of variables being of the first order in M_{gas}/M.

What remains after subtracting the time average is a fluctuation which is the same for both sets of variables, see Fig. 3(b). Its root mean square is $2.0 \times 10^{-4} h_0$ in the simulation with 400×25 particles, and $1.1 \times 10^{-4} h_0$ and $9.6 \times 10^{-5} h_0$ in those with 800×25 and 400×50 particles, respectively. For $M_{gas}/M = 10^{-2}$ it grows in time and is visible in Fig. 3(a) for $t > 10^4$; in this case its root mean square is $9.1 \times 10^{-3} h_0$ with 400×25 particles and $2.2 \times 10^{-3} h_0$ with 800×25 particles. Hence, these fluctuations are caused by the finite number of particles.

C. Self-gravity and conservation of the energy

We have tested the computation of the gravitational field in a situation which is dominated by self-gravity, namely the collapse of a cold homogeneous sphere of mass M_{gas} and radius R in the absence of an external potential. Since in this case all the particles have $L = 0$, this test has been performed with an implementation of the same algorithm for a general distribution of the angular momenta. Before the shells cross each other, the shell with initial radius r_0 follows the cycloid $r = r_0 \cos^2\theta$, $t = (\theta + \sin\theta\cos\theta)/\sqrt{2GM_{gas}/R^3}$. For $\Delta r = 0.01R$, the median of the relative error of the radii of the shells with $0.1 < r_0/R < 0.9$ at the time $t = 0.8\sqrt{R^3/(GM_{gas})}$ is 1.8×10^{-5} for $N = 200$ and 4.3×10^{-6} for $N = 400$ shells, which corresponds to second-order convergence in $1/N$.

In the simulations of the gas cloud in section VI, which use $\Delta t = 0.05$ and extend to $t = 10^4$ (2×10^4 for $\sigma_J = 0.05$), the largest relative change of the total energy (38) is 1.7×10^{-4} for $M_{gas}/M = 0.3$, it does not exceed 7.2×10^{-5} for the remaining simulations, and its median is 1.0×10^{-5}. [(Erik) A test of the dependency on the time step for the cloud (for instance $M_{gas}/M = 0.01$ and 0.1 with $\Delta t = 0.1$ and 0.025) is to be added; it requires new simulations.]

VI. INITIAL DATA FAR FROM EQUILIBRIUM

In this section we consider the gas cloud (42), which is concentrated about one orbital phase and is far from any steady state, and we analyze how its evolution changes when its mass is increased. We start with the simulations and with a parameter μ which compares the amplitude of the potential generated by the cloud with the spread of its orbital frequencies. Next, we show that the cloud mixes and its potential relaxes when μ is small, whereas it collapses into a self-bound clump which does not mix when μ exceeds a critical value, and we analyze the dependence of this value on the widths of the cloud and on the external potential. Finally, we describe the state of the gas at late times.

A. Simulations and the parameter μ

The reference family of simulations consists of the cloud (42) with $\sigma_Q = \sigma_J = 0.1$ and $L_0 = 2$ in the isochrone potential, with the masses $M_{gas}/M = 10^{-3}$, 0.01, 0.015, 0.02, 0.025, 0.03, 0.04, 0.05, 0.07, 0.1 and 0.3, together with a simulation without self-gravity. In order to analyze the dependence on the shape of the cloud, we vary its width in angle, $\sigma_Q = 0.2$ and 0.4 with $M_{gas}/M = 0.01$, 0.02, 0.04 and 0.08, and its width in action, $\sigma_J = 0.05$ with

$M_{gas}/M = 0.005$, 0.01 and 0.02 and $\sigma_J = 0.2$ with $M_{gas}/M = 0.02$, 0.04 and 0.08. Finally, we consider the cloud with $\sigma_Q = \sigma_J = 0.1$ in the Kepler potential, with $M_{gas}/M = 0.01$, 0.02, 0.04 and 0.08. This is the external potential of the nonlinear phase mixing result of Ref. [35], although the reduced model and the support of the DF, which extends to the circular orbit, differ from the assumptions made there. The simulations use the lattice (41) with $N_J \times N_Q = 400 \times 25$, except for $\sigma_Q = 0.2$ ($N_Q = 50$) and $\sigma_J = 0.2$ ($N_J = 800$), and the time step $\Delta t = 0.05$, and they extend to $t = 10^4$ (2×10^4 for $\sigma_J = 0.05$). In order to test the convergence, we have repeated the simulations of the reference family with $M_{gas}/M = 0.01$, 0.02 and 0.1 with $N_Q = 100$, and the one with $M_{gas}/M = 0.01$ with $N_J = 800$. In all the simulations the gas particles remain bound, and the observables are computed with the action-angle variables of the instantaneous potential and the weight function $B = 1$, see sections II E and IV A.

In the absence of self-gravity the cloud mixes on a time scale which is set by the spread of its orbital frequencies. We characterize this spread by the interval $[\underline{J}, \overline{J}]$ in which the factor $J^2 \exp(-J^2/\sigma_J^2)$ of the DF (42) exceeds one percent of its maximum, by its width $\Delta J := \overline{J} - \underline{J}$, and by the width $\Delta\Omega := \Omega(\underline{J}) - \Omega(\overline{J})$ of the corresponding frequency band of the external potential. After the time

$$\tau_1 := \frac{2\pi}{\Delta\Omega} \tag{49}$$

the orbits at the two ends of this interval have drifted apart by one revolution, such that the first harmonic of the cloud has been spread over the orbit. For $\sigma_Q = \sigma_J = 0.1$ in the isochrone potential one has $\Delta J = 0.270$, $[\Omega(\overline{J}), \Omega(\underline{J})] = [0.0513, 0.0705]$ and $\tau_1 = 327$, for $\sigma_J = 0.05$ and 0.2 one has $\tau_1 = 589$ and 198, respectively, and in the Kepler potential $\tau_1 = 161$. Hence, the simulations extend over at least 30 units of τ_1.

The self-gravity of the cloud couples the motions of its particles and acts against the mixing. In order to compare both effects, consider the motion of the gas particles relative to the center of the cloud. In the absence of self-gravity, the angle variable of a particle relative to the one of the center drifts with a velocity which varies by $\Delta\Omega$ across the support of the cloud. Approximating the derivative of the frequency by the constant $|\Omega'| = \Delta\Omega/\Delta J$, the energy which is associated with this differential drift, $|\Omega'|(J - J_c)^2/2$ with J_c the action variable of the center, is of the order of $\Delta\Omega\,\Delta J$. On the other hand, the strength of the self-gravity is measured by the amplitude A of the variation of the potential which is generated by the cloud. This suggests the parameter

$$\mu := \left(\frac{A}{\Delta\Omega\,\Delta J}\right)^{1/2} = \frac{\omega_b}{\Delta\Omega}, \qquad \omega_b := (|\Omega'|A)^{1/2}, \tag{50}$$

whose square compares these two energies, and one expects the self-gravity of the cloud to prevent the mixing when μ is of order one. The frequency ω_b is the frequency of the small oscillations of a pendulum with the Hamiltonian $|\Omega'|(J - J_c)^2/2 - A\cos\vartheta$, which describes the trapping of particles by a wave in a plasma [56]; in terms of μ, $2\pi/\omega_b = \tau_1/\mu$. This argument is a heuristic one: it does not describe how the potential of the cloud depends on the positions of its particles, and it does not predict the critical value of μ. We measure A as half of the largest variation of $\Phi_{gas}(t, r)$ at fixed r during the first radial period $0 \le t \le 2\pi/\Omega(\overline{J})$, maximized over $3 \le r \le 15$. For small masses A is proportional to the mass of the gas: in the reference family $A = 4.9 \times 10^{-5}$ and 4.9×10^{-4} for $M_{gas}/M = 10^{-3}$ and 10^{-2}, such that μ grows like $(M_{gas}/M)^{1/2}$. For these two masses $\mu = 0.10$ and 0.31, and for $M_{gas}/M = 0.1$ and 0.3 one has $\mu = 0.89$ and 1.16.

B. Mixing and self-bound clump

Figure 4 shows the computational particles in the (r, p_r)-plane for two clouds of the reference family. For $M_{gas}/M = 0.01$ ($\mu = 0.31$) the evolution resembles the mixing without self-gravity of Fig. 1: the cloud is sheared into a spiral which winds up around the circular orbit, and at $t = 30\tau_1$ the gas fills the region of phase space which is covered by its orbits, without any visible concentration. For $M_{gas}/M = 0.1$ ($\mu = 0.89$), in contrast, a part of the cloud contracts within the first radial periods into a compact clump, which does not mix: at $t = 30\tau_1$ it is still present, and the rest of the gas is mixed around it. The center of the clump oscillates radially about the circular orbit. Defining the clump by the particles whose action variable is smaller than 0.05 at $t = 10^4$, we find that for $t \ge 15\tau_1$ its center moves between $r = 4.8$ and $r = 6.5$, with the frequency 0.075, while the root mean square distance of its particles from the center is 0.3. This frequency is close to the frequency $\Omega(0) = 0.0711$ of the circular orbit in the external potential.

The function $|h_1(t)|/h_0$ and the oscillation of the potential are shown in Fig. 5 for the reference family. Without self-gravity, $|h_1|$ decays like t^{-3} (a fit of the envelope over $10 \le t/\tau_1 \le 30$ gives the exponent -3.03), which is the analogue of the tail (33) for the endpoint $J = 0$ of the support, at which $\hat{F}_1$ vanishes like J^2. With self-gravity the decay slows down, and the level which is reached at late times increases with μ. Averaged over $10 \le t/\tau_1 \le 20$, $|h_1|/h_0$ is 1.1×10^{-4}, 2.4×10^{-3} and 1.2×10^{-2} for $\mu = 0.10$, 0.31 and 0.43 (the last two values for $N_Q = 100$),




FIG. 4: The computational particles in the (r, p_r)-plane at four different times for the cloud (42) with $\sigma_Q = \sigma_J = 0.1$ in the isochrone potential, for $M_{gas}/M = 0.01$ (upper panels) and $M_{gas}/M = 0.1$ (lower panels). The color indicates the mass of each particle relative to the largest one. For $M_{gas}/M = 0.01$ the cloud mixes, whereas for $M_{gas}/M = 0.1$ it forms a clump which persists until the end of the simulation.




FIG. 5: Evolution of the cloud (42) with $\sigma_Q = \sigma_J = 0.1$ in the isochrone potential for different masses, labeled by the value of μ. (a) Envelope of $|h_1|/h_0$ (largest value in a moving window of width 100), computed with the action-angle variables of the instantaneous potential, for $N_J \times N_Q = 400 \times 25$ (solid lines) and 400×100 (dotted lines). (b) Deviation of the potential generated by the gas from its average over the second half of the simulation, measured with the norm (28) over $3 \le r \le 15$ in units of the largest value of $|\langle\Phi_{gas}\rangle|$ (envelope as in (a)). The potential is computed from the radii of the particles; without self-gravity (dashed line) it is the potential which the gas would generate if it acted on itself.

compared to 7.0×10^{-5} without self-gravity, and it lies between 0.09 and 0.85 for $\mu \geq 0.48$. The potential which is generated by the gas behaves in the same way, see Fig. 5(b). Without self-gravity, the potential which the gas would generate relaxes to its final profile through the mixing alone: its deviation (27) from the average over the second half of the simulation falls below 10^{-3} of the largest value of $|\langle\Phi_{gas}\rangle|$ at $t = 2.9\tau_1$. With self-gravity and $\mu \leq 0.43$ the potential still relaxes, but more slowly: the same level is reached at $t = 3.3\tau_1$, $4.8\tau_1$, $5.5\tau_1$ and $6.2\tau_1$ for $\mu = 0.10$, 0.31, 0.37 and 0.43, respectively. Over $10 \leq t/\tau_1 \leq 20$ the deviation lies between 7×10^{-5} and 3×10^{-4}, compared to 4×10^{-5} without self-gravity, and it is reduced when the number of particles is increased (dotted lines). For $\mu \geq 0.48$, in contrast, the deviation never falls below 10^{-3}: over $10 \leq t/\tau_1 \leq 20$ it lies between 1.6×10^{-3} and 4.6×10^{-2}, since the potential oscillates with the clump. Therefore, below the transition the gravitational field generated by the gas relaxes to a stationary field while the DF continues to mix, which is the analogue of Landau damping, whereas above the transition the field does not relax. (Without self-gravity the deviation increases again for $t \gtrsim 15\tau_1$, which we attribute to the spurious recurrence (48) of the higher harmonics of the potential on the lattice; for $n = 10$ the recurrence time is about $4800 \approx 15\tau_1$.)

C. The transition




FIG. 6: Mean value of $|h_1|/h_0$ over the interval $10 \leq t/\tau_1 \leq 20$ as a function of the parameter (50), for all the simulations of the cloud (42). The families differ in the widths σ_Q and σ_J of the cloud (equal to 0.1 unless otherwise stated) and in the external potential (isochrone unless otherwise stated). The open circles refer to the simulations of the reference family with $N_Q = 100$ or $N_J = 800$. The shaded band indicates the interval $0.43 \leq \mu \leq 0.48$.

Figure 6 shows the mean value of $|h_1|/h_0$ over the interval $10 \leq t/\tau_1 \leq 20$ as a function of μ for all the simulations. In each family, $|h_1|/h_0$ grows by one to two orders of magnitude within a narrow interval of μ. With the exception of the family with $\sigma_Q = 0.4$, the location of this interval is the same for all the families: all the simulations with $\mu \leq 0.43$ have $|h_1|/h_0 \leq 2.1 \times 10^{-2}$, and all those with $\mu \geq 0.48$ have $|h_1|/h_0 \geq 0.09$. Table II lists, for each family, the largest value of μ for which the cloud mixes and the smallest one for which it forms a clump. The intervals which are delimited by these values have the common part $0.43 \leq \mu \leq 0.48$ for the five families with $\sigma_Q \leq 0.2$, whereas for $\sigma_Q = 0.4$ the transition lies between $\mu = 0.52$ and $\mu = 0.69$. Therefore, for clouds which are compact in angle the critical value μ_c of the parameter (50) lies between 0.43 and 0.48, independently of the width of the cloud in action and of the external potential, whereas a cloud which is wide in angle requires a larger value of μ in order to form a clump. The critical value is of order one, as expected from the argument of section VI A, but this argument does not predict it. Hence, μ captures the competition between self-gravity and shear for compact clouds, but not the complete dependence on the shape of the cloud. [(Erik) A theory of μ_c, including its dependence on σ_Q, is still missing.]

Away from the transition the results do not depend on the resolution. For $\mu = 0.31$ we obtain $|h_1|/h_0 = 2.62 \times 10^{-3}$ with $N_J \times N_Q = 400 \times 25$ and 2.44×10^{-3} with both 400×100 and 800×25, and for $\mu = 0.89$ we obtain 0.80 with 400×25 and 0.85 with 400×100. Close to the transition, in contrast, the result depends on the number of

potential	widths	mixing μ	mixing $\lvert h_1\rvert/h_0$	clump μ	clump $\lvert h_1\rvert/h_0$
isochrone	$\sigma_Q = \sigma_J = 0.1$	0.43	1.2×10^{-2}	0.48	0.091
	$\sigma_J = 0.05$	0.35	8.3×10^{-3}	0.49	0.105
	$\sigma_J = 0.2$	0.38	5.8×10^{-3}	0.53	0.33
	$\sigma_Q = 0.2$	0.42	7.4×10^{-3}	0.58	0.14
	$\sigma_Q = 0.4$	0.52	1.4×10^{-2}	0.69	0.10
Kepler	$\sigma_Q = \sigma_J = 0.1$	0.37	8.8×10^{-3}	0.51	0.23

TABLE II: Bracketing of the transition: for each family, the largest value of μ for which the cloud mixes and the smallest one for which it forms a clump, together with the mean value of $|h_1|/h_0$ over $10 \leq t/\tau_1 \leq 20$. In the family with $\sigma_J = 0.05$ ($\sigma_Q = 0.2$, 0.4) the remaining width is $\sigma_Q = 0.1$ ($\sigma_J = 0.1$). For $\mu = 0.43$ the value refers to $N_Q = 100$; for $N_Q = 25$ it is 2.1×10^{-2}.

particles: for $\mu = 0.43$ it is 2.1×10^{-2} with 400×25 and 1.2×10^{-2} with 400×100. This is to be expected, since near the transition the outcome is sensitive to small perturbations, such as those which are due to the discreteness of the particles. However, both simulations lie on the same side of the transition.

In the Kepler potential the clouds with $M_{gas}/M = 0.01$ and 0.02 ($\mu = 0.26$ and 0.37) mix, which is the behavior described by Ref. [35] for small data, whereas those with $M_{gas}/M = 0.04$ and 0.08 ($\mu = 0.51$ and 0.69) form a clump. Since μ grows with the mass, for a cloud of a given shape the mixing fails above a critical mass, which lies between $M_{gas}/M = 0.02$ and 0.04 in this case. Hence, in the reduced model, the phase mixing of such a cloud requires its mass to be small.

D. The state at late times

Below the transition, the mixing continues after the decay of Fig. 5(a) has slowed down. In the simulations of the reference family with $\mu = 0.31$ and $N_Q = 100$ or $N_J = 800$, $|h_1|/h_0$ decreases from 2.4×10^{-3} for $10 \leq t/\tau_1 \leq 20$ to 7.5×10^{-4} and 8.6×10^{-4} for $t/\tau_1 \geq 20$, respectively. With the default lattice, in contrast, it remains at 2.8×10^{-3}, which is due to the fluctuations generated by the discreteness of the particles (see section V B). Hence, below the transition the self-gravity of the cloud slows down the mixing, but it does not stop it. A quantitative analysis of this late decay requires larger numbers of particles than those used here. [(Erik) To be determined with $N \geq 10^5$ particles: the decay law of $|h_1|$ below the transition.]

Above the transition the clump persists until the end of the simulations. For $0.76 \leq \mu \leq 1.16$ the amplitude of the radial oscillation of its center decreases by at most 25 percent between the intervals $3 \leq t/\tau_1 \leq 10$ and $20 \leq t/\tau_1 \leq 30$, and not at all for $\mu = 1.16$, such that within the time span of our simulations the oscillation does not die out. The value of $|h_1|$ is not a monotonic function of μ above the transition: for $\mu = 0.52$, 0.60, 0.66, 0.76, 0.89 and 1.16 it is 0.22, 0.32, 0.19, 0.58, 0.80 and 0.69, respectively. This reflects the amplitude of the oscillation of the clump, which determines how strongly the gas is concentrated in angle. For $\mu = 0.66$, for instance, the center of the clump moves by ± 0.16 about its mean radius, which is less than the size 0.32 of the clump, whereas for $\mu = 0.89$ it moves by ± 0.9. The mass of the clump cannot be determined in a unique way by means of a cut in the action variable: for $\mu = 0.89$ the particles with $J < 0.03$, 0.05 and 0.08 at $t = 10^4$ carry 58, 75 and 83 percent of the mass of the gas, and below the transition such cuts also select part of the mixed gas (for $\mu = 0.31$, 21 percent of the mass has $J < 0.05$). For this purpose a dynamical criterion would be required, which identifies the particles that are bound to the clump. [(Erik) To be done: a dynamical criterion for the mass of the clump.]

The state above the transition, which consists of a compact clump whose oscillation persists and of mixed gas around it, resembles the core-halo states which are formed by collisionless relaxation in systems with long-range interactions [36]. In the scenario described in that reference, the particles which are in parametric resonance with the oscillation of the core escape to the halo, and this damps the oscillation until a quasi-stationary state is reached. In our simulations the oscillation of the clump is damped only slowly, if at all, over 30 units of τ_1.

VII. CONCLUSIONS

In these notes we have studied numerically the relaxation through phase space mixing of a spherically symmetric collisionless gas which is trapped in an external central potential and whose initial DF is concentrated about one orbital phase, taking into account the self-gravity of the gas, in the reduced model in which all the gas particles have

the same magnitude of the angular momentum. Our analysis is based on a PIC code whose particles are placed on a regular lattice in action-angle variables. In this section we summarize the results and list some open problems.

In the absence of self-gravity, the PIC code reproduces the exact solution of the mixing problem to within 6×10^{-9} in relative terms over 10^4 time units, during which the first harmonic of the DF decays by seven orders of magnitude. When the self-gravity of the gas is taken into account, the observables which measure the mixing have to be computed with the action-angle variables of the total potential, or of the instantaneous potential when the gas is far from equilibrium. The action-angle variables of the external potential leave a time-independent residue which is proportional to the mass of the gas (2×10^{-3} for $M_{gas}/M = 10^{-3}$) and which could be mistaken for an incomplete mixing.

The self-gravity of the cloud first slows down the mixing and then stops it. When the parameter (50), which compares the amplitude of the potential generated by the cloud with the spread of its orbital frequencies, is smaller than a critical value μ_c, the gas mixes, more slowly than in the absence of self-gravity, and the potential which it generates relaxes to a stationary profile while the DF continues to mix, which is the analogue of Landau damping. The time in which the deviation of the potential falls below 10^{-3} of its largest value grows from 2.9 units of the mixing time τ_1 without self-gravity to $6.2\,\tau_1$ for $\mu = 0.43$. Above μ_c the gas collapses into a compact, self-bound clump which oscillates radially about the circular orbit, with a frequency close to the one of the circular orbit of the external potential, and which does not mix during the 30 units of τ_1 covered by our simulations; accordingly, the potential does not relax. For clouds which are compact in angle, μ_c lies between 0.43 and 0.48 in the isochrone potential as well as in the Kepler potential, independently of the width of the cloud in action, whereas for a cloud which is wide in angle the transition lies between $\mu = 0.52$ and $\mu = 0.69$. In the Kepler potential this means that, for a cloud of a given shape, phase mixing in the sense of Ref. [35] only takes place below a critical mass, which lies between 0.02 and 0.04 times the central mass for the clouds considered here.

Several questions remain open. A theory of the critical value μ_c is missing, which should also explain why a cloud which is wide in angle requires a larger value; the heuristic argument of section VI A only shows that μ_c should be of order one. The mass of the clump has to be defined by a dynamical criterion, which identifies the particles that are bound to it. The decay laws of $|h_1|$ and of the deviation of the potential below the transition, and the dependence of the results on the number of particles close to the transition, require simulations with more particles than those used here. It is not known whether the oscillation of the clump is eventually damped, as in the core-halo states of systems with long-range interactions [36], and on which time scale; answering this question requires longer simulations. Finally, all our results refer to the reduced model. When the angular momenta of the gas particles are distributed, the frequencies of the orbits depend also on L, which provides an additional source of mixing (see section III), and the initial data can be chosen to satisfy the assumptions of Ref. [35]. Whether this shifts the critical value of μ is a natural next step, for which the methods of these notes can be extended in a direct way. [(Erik) To be completed after the discussion with Olivier: the outlook regarding the distribution of angular momenta, and the reference to the companion article on perturbations of steady states.]

Acknowledgments

[(Erik) To be written.]

Appendix A: Numerical computation of the action-angle variables

In this appendix we describe the numerical evaluation of the action-angle variables defined in Eqs. (16,17,18) for a potential of the form $\Phi = \Phi_{ext} + \Phi_{gas}$, in which Φ_{gas} is only known on a radial grid. This is required for the construction of the self-consistent steady states, for the generation of the initial data and for the computation of the observables $h_n(t)$.

The values of Φ_{gas} on the grid are interpolated with a natural cubic spline, and beyond the last grid point we set $\Phi_{gas} = -GM_{gas}/r$. Given a point (r, p_r, L) we compute its energy E from Eq. (15), locate the minimum $r_{min}(L)$ of the effective potential V_L, and determine the turning points $r_1 < r_{min}(L) < r_2$ by solving the equation $V_L(r) = E$ with the bisection method on either side of the minimum. The integrand in Eq. (16) behaves like a square root at the turning points and the one in Eq. (17) like an inverse square root, which spoils the convergence of standard quadrature rules. We avoid this problem by means of the substitution

$$r = r_m + r_a \sin\theta, \qquad r_m := \frac{r_1 + r_2}{2}, \quad r_a := \frac{r_2 - r_1}{2}, \qquad -\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}. \tag{A1}$$

Since for a non-circular orbit the zeros of the function $E - V_L$ at the turning points are simple, one has $E - V_L(r) =$

$r_a^2 \cos^2\theta\, u(\theta)$ with a smooth and positive function u, and it follows that

$$A(E,L) = 2r_a^2 \int\limits_{-\pi/2}^{\pi/2} \cos^2\theta \sqrt{2u(\theta)}\, d\theta, \qquad T(E,L) = 2 \int\limits_{-\pi/2}^{\pi/2} \frac{d\theta}{\sqrt{2u(\theta)}}. \tag{A2}$$

Both integrands are smooth, and we evaluate the integrals with a Gauss-Legendre rule with 64 nodes. The angle variable is obtained in the same way from

$$Q = \frac{2\pi}{T(E,L)} \int\limits_{-\pi/2}^{\theta(r)} \frac{d\theta'}{\sqrt{2u(\theta')}}, \qquad \theta(r) := \arcsin\left(\frac{r-r_m}{r_a}\right), \tag{A3}$$

when $p_r \geq 0$, and from 2π minus the right-hand side of Eq. (A3) when $p_r < 0$. For the isochrone potential without self-gravity, for which the exact expressions (22,23) are available, this procedure yields the action variable with an error of the order of 10^{-14} and the period with an error of the order of 10^{-11}. Without the substitution (A1) the corresponding errors are 2×10^{-6} and 9×10^{-3}, and they only decrease like the third and the first inverse power of the number of nodes, respectively.

For the inverse map $(Q, J) \mapsto (r, p_r)$ at fixed L we tabulate the function $J = A(E,L)/2\pi$ for 4000 values of the energy between $E_{min}(L)$ and a value slightly above the cutoff energy, and obtain $E(J,L)$ by interpolation. Next, we determine the turning points and find, by bisection in θ, the point of the orbit at which the time elapsed since the passage through the pericenter is equal to $QT(E,L)/2\pi$ when $Q \leq \pi$ and to $(2\pi - Q)T(E,L)/2\pi$ otherwise. This gives the radius r, and the momentum is $p_r = \pm\sqrt{2(E - V_L(r))}$, with the positive sign when $Q \leq \pi$. Applying the direct map to the result reproduces the points of the lattice (41) with errors which are smaller than 10^{-8} in J and 5×10^{-7} in Q. The frequency is obtained from $\Omega = 2\pi/T(E,L)$.

[1] J. Binney and S. Tremaine. *Galactic Dynamics*. Princeton University Press, Princeton, second edition, 2008.

[2] G. Rein. Collisionless kinetic equations from astrophysics – the Vlasov-Poisson system. In *Handbook of Differential Equations: Evolutionary Equations*, volume 3, pages 383–476. Elsevier, 2007.

[3] H. Andréasson. The Einstein-Vlasov System/Kinetic Theory. *Living Rev. Relativ.*, 14:4, 2011.

[4] J. R. Oppenheimer and H. Snyder. On Continued Gravitational Contraction. *Phys. Rev.*, 56:455–459, 1939.

[5] D. Christodoulou. Violation of cosmic censorship in the gravitational collapse of a dust cloud. *Commun. Math. Phys.*, 93:171–195, 1984.

[6] N. Ortiz and O. Sarbach. Conformal diagrams for the gravitational collapse of a spherical dust cloud. *Class. Quantum Grav.*, 28:235001, 2011.

[7] K. Pfaffelmoser. Global classical solutions of the Vlasov-Poisson system in three dimensions for general initial data. *J. Differ. Equ.*, 95:281–303, 1992.

[8] P. L. Lions and B. Perthame. Propagation of moments and regularity for the 3-dimensional Vlasov-Poisson system. *Invent. Math.*, 105:415–430, 1991.

[9] J. Schaeffer. Global existence of smooth solutions to the Vlasov Poisson system in three dimensions. *Commun. Partial Differ. Equ.*, 16:1313–1335, 1991.

[10] G. Rein and A. D. Rendall. Global existence of solutions of the spherically symmetric Vlasov-Einstein system with small initial data. *Commun. Math. Phys.*, 150:561–583, 1992.

[11] H. Andréasson, M. Kunze, and G. Rein. The formation of black holes in spherically symmetric gravitational collapse. *Math. Ann.*, 350:683–705, 2011.

[12] D. Lynden-Bell. The Stability and Vibrations of a Gas of Stars. *Mon. Not. R. Astron. Soc.*, 124:279–296, 1962.

[13] D. Lynden-Bell. Statistical Mechanics of Violent Relaxation in Stellar Systems. *Mon. Not. R. Astron. Soc.*, 136:101–121, 1967.

[14] P. Rioseco and O. Sarbach. Phase space mixing in an external gravitational central potential. *Class. Quantum Grav.*, 37:195027, 2020.

[15] S. Tremaine, M. Hénon, and D. Lynden-Bell. H-functions and mixing in violent relaxation. *Mon. Not. R. Astron. Soc.*, 219:285–297, 1986.

[16] C. Mitchell. Weak convergence to equilibrium of statistical ensembles in integrable Hamiltonian systems. *J. Math. Phys.*, 60:052702, 2019.

[17] P. Domínguez-Fernández, E. Jiménez-Vázquez, M. Alcubierre, E. Montoya, and D. Núñez. Description of the evolution of inhomogeneities on a dark matter halo with the Vlasov equation. *Gen. Relativ. Gravit.*, 49:123, 2017.

[18] A. L. García-Perciante, M. Guerrero, D. Núñez, and O. Sarbach. Relaxation of a Vlasov gas to an inhomogeneous state due to phase space mixing in an axisymmetric potential: A Newtonian analogy of the Kerr orbital motion. *arXiv:2610.06913*, 2026.

[19] P. Rioseco and O. Sarbach. Phase space mixing in the equatorial plane of a Kerr black hole. *Phys. Rev. D*, 98:124024, 2018.

[20] P. Rioseco and O. Sarbach. Phase Space Mixing of a Vlasov Gas in the Exterior of a Kerr Black Hole. *Commun. Math. Phys.*, 405:105, 2024.

[21] M. Moreno, P. Rioseco, and H. Van Den Bosch. Mixing in anharmonic potential well. *J. Math. Phys.*, 63:071502, 2022.

[22] S. Chaturvedi and J. Luk. Phase mixing for solutions to 1D transport equation in a confining potential. *Kinet. Relat. Models*, 15:403–416, 2022.

[23] M. Hadžić, G. Rein, M. Schrecker, and C. Straub. Quantitative phase mixing for Hamiltonians with trapping. *arXiv:2405.17153*, 2024.

[24] L. D. Landau. On the vibrations of the electronic plasma. *J. Phys. (USSR)*, 10:25–34, 1946.

[25] C. Mouhot and C. Villani. On Landau damping. *Acta Math.*, 207:29–201, 2011.

[26] M. Hadžić, G. Rein, and C. Straub. On the Existence of Linearly Oscillating Galaxies. *Arch. Ration. Mech. Anal.*, 243:611–696, 2022.

[27] M. Kunze. *A Birman-Schwinger Principle in Galactic Dynamics*, volume 77 of *Progress in Mathematical Physics*. Birkhäuser, Cham, 2021.

[28] M. Kunze. A Birman–Schwinger principle in galactic dynamics: ESI, Vienna, 07–11 February 2022. *Class. Quantum Grav.*, 39:244001, 2022.

[29] S. D. Mathur. Existence of oscillation modes in collisionless gravitating systems. *Mon. Not. R. Astron. Soc.*, 243:529–536, 1990.

[30] M. Moreno, P. Rioseco, and H. Van Den Bosch. Spectrum of the linearized Vlasov–Poisson equation around steady states from galactic dynamics. *J. Math. Phys.*, 67:091501, 2026.

[31] M. Hadžić, G. Rein, M. Schrecker, and C. Straub. Damping Versus Oscillations for a Gravitational Vlasov–Poisson System. *Arch. Ration. Mech. Anal.*, 249:45, 2025.

[32] M. Hadžić and M. Schrecker. On quantitative linear gravitational relaxation. *arXiv:2505.14856*, 2025.

[33] T. Ramming and G. Rein. Oscillating solutions of the Vlasov–Poisson system—A numerical investigation. *Physica D*, 365:72–79, 2018.

[34] C. Straub. Numerical experiments on stationary, oscillating, and damped spherical galaxy models. *Physica D*, 470:134351, 2024.

[35] S. Chaturvedi and J. Luk. Linear and Nonlinear Phase Mixing for the Gravitational Vlasov–Poisson System Under an External Kepler Potential. *Arch. Ration. Mech. Anal.*, 250:68, 2026.

[36] Y. Levin, R. Pakter, F. B. Rizzato, T. N. Teles, and F. P. C. Benetti. Nonequilibrium statistical mechanics of systems with long-range interactions. *Phys. Rep.*, 535:1–60, 2014.

[37] M. Hénon. Numerical experiments on the stability of spherical stellar systems. *Astron. Astrophys.*, 24:229–238, 1973.

[38] P. Klinko and B. N. Miller. Angular-momentum-induced phase transition in spherical gravitational systems: N-body simulations. *Phys. Rev. E*, 65:056127, 2002.

[39] S. L. Shapiro and S. A. Teukolsky. Relativistic stellar dynamics on the computer. I - Motivation and numerical method. *Astrophys. J.*, 298:34–57, 1985.

[40] G. Rein, A. D. Rendall, and J. Schaeffer. Critical collapse of collisionless matter: A numerical investigation. *Phys. Rev. D*, 58:044007, 1998.

[41] I. Olabarrieta. Critical collapse of collisionless matter in spherical symmetry. Master's thesis, University of British Columbia, 2000. arXiv:gr-qc/0012059.

[42] I. Olabarrieta and M. W. Choptuik. Critical phenomena at the threshold of black hole formation for collisionless matter in spherical symmetry. *Phys. Rev. D*, 65:024007, 2002.

[43] H. Andréasson and G. Rein. A numerical investigation of the stability of steady states and critical phenomena for the spherically symmetric Einstein–Vlasov system. *Class. Quantum Grav.*, 23:3659–3677, 2006.

[44] C. K. Birdsall and A. B. Langdon. *Plasma Physics via Computer Simulation*. CRC Press, Boca Raton, 1991.

[45] R. W. Hockney and J. W. Eastwood. *Computer Simulation Using Particles*. Adam Hilger, Bristol, 1988.

[46] G. Lapenta. Kinetic plasma simulation: particle in cell method. In *XII Carolus Magnus Summer School on Plasma and Fusion Energy Physics*. 2015.

[47] M. Shalaby, A. E. Broderick, P. Chang, C. Pfrommer, A. Lamberts, and E. Puchwein. SHARP: A Spatially Higher-order, Relativistic Particle-in-cell Code. *Astrophys. J.*, 841:52, 2017.

[48] G. Rein and T. Rodewis. Convergence of a particle-in-cell scheme for the spherically symmetric Vlasov-Einstein system. *Indiana Univ. Math. J.*, 52:821–861, 2003.

[49] M. Hénon. L'amas isochrone: I. *Ann. Astrophys.*, 22:126, 1959.

[50] J. Binney. Hénon's isochrone model. *arXiv:1411.4937*, 2014.

[51] C. M. Bender and S. A. Orszag. *Advanced Mathematical Methods for Scientists and Engineers I: Asymptotic Methods and Perturbation Theory*. Springer, New York, 1999.

[52] J. Barré, A. Olivetti, and Y. Y. Yamaguchi. Algebraic damping in the one-dimensional Vlasov equation. *J. Phys. A: Math. Theor.*, 44:405502, 2011.

[53] J. Barré and Y. Y. Yamaguchi. On algebraic damping close to inhomogeneous Vlasov equilibria in multi-dimensional spaces. *J. Phys. A: Math. Theor.*, 46:225501, 2013.

[54] H. Yoshida. Construction of higher order symplectic integrators. *Phys. Lett. A*, 150:262–268, 1990.

[55] NIST Digital Library of Mathematical Functions. `https://dlmf.nist.gov/`, Release 1.2.8 of 2026-09-15. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl,

and M. A. McClain, eds.
[56] T. O'Neil. Collisionless damping of nonlinear plasma oscillations. *Phys. Fluids*, 8:2255–2262, 1965.