

ON THE INVERSE PROBLEM OF THE k -TH DAVENPORT CONSTANTS FOR GROUPS OF RANK 2

QINGHAI ZHONG

ABSTRACT. For a finite abelian group G and a positive integer k , let $D_k(G)$ denote the smallest integer ℓ such that each sequence over G of length at least ℓ has k disjoint nontrivial zero-sum subsequences. It is known that $D_k(G) = n_1 + kn_2 - 1$ if $G \cong C_{n_1} \oplus C_{n_2}$ is a rank 2 group, where $1 < n_1 | n_2$. We investigate the associated inverse problem for rank 2 groups, that is, characterizing the structure of zero-sum sequences of length $D_k(G)$ that can not be partitioned into $k + 1$ nontrivial zero-sum subsequences.

1. INTRODUCTION

Let $(G, +, 0)$ be a finite abelian group. By a sequence S over G , we mean a finite sequence of terms from G which is unordered, repetition of terms allowed. We say that S is a zero-sum sequence if the sum of its terms equals zero and denote by $|S|$ the length of the sequence.

Let k be a positive integer. We denote by $D_k(G)$ the smallest integer ℓ such that every sequence over G of length at least ℓ has k disjoint nontrivial zero-sum subsequences. We call $D_k(G)$ the k -th Davenport constant of G , while the Davenport constant $D(G) = D_1(G)$ is one of the most important zero-sum invariants in Combinatorial Number Theory and, together with Erdős-Ginzburg-Ziv constant, η -constant, etc., has been studied a lot (see [39, 40, 1, 29, 49, 50, 30, 16, 21, 41, 5, 43, 6, 14, 7, 22, 38]). This variant $D_k(G)$ of the Davenport constant was introduced and investigated by F. Halter-Koch [37], in the context of investigations on the asymptotic behavior of certain counting functions of algebraic integers defined via factorization properties (see the monograph [27, Section 6.1], and the survey article [19, Section 5]). In 2014, K. Cziszter and M. Domokos ([9, 8]) introduced the generalized Noether Number $\beta_k(G)$ for general groups, which equals $D_k(G)$ when G is abelian (see [11, 12, 10] for more about this direction). Knowledge of those constants is highly relevant when applying the inductive method to determine or estimate the Davenport constant of certain finite abelian groups (see [13, 4, 3, 42]).

In 2010, M. Freeze and W. Schmid ([17]) showed that for each finite abelian group G we have $D_k(G) = D_0(G) + k \exp(G)$ for some $D_0(G) \in \mathbb{N}_0$ and all sufficiently large k . In fact, it is known that for groups of rank at most two, and for some other types of groups, an equality of the form

2010 *Mathematics Subject Classification.* 11B30, 11P70.

Key words and phrases. zero-sum sequences, Davenport constant, k -th Davenport constant.

This work was supported by the Austrian Science Fund FWF [grant DOI:10.55776/P36852].

$D_k(G) = D_0(G) + k \exp(G)$ for some $D_0(G) \in \mathbb{N}_0$ holds for all k . In particular, for a rank two abelian group $G = C_m \oplus C_n$, where $m \mid n$, we have $D_k(G) = m + kn - 1$ ([27, Theorem 6.1.5]). Yet, it fails for elementary 2 and 3-groups of rank at least 3 (see [13, 4]). In general, computing (even bounding) $D_k(G)$ is quite more complicated than for $D(G)$, in particular for (elementary) p -groups, while $D(G)$ is known for p -groups.

In zero-sum theory, the associated inverse problems of zero-sum invariants study the structure of extremal sequences that do not have enough zero-sum subsequences with the prescribed properties. The inverse problems of the Davenport constant, the η -constant, and the Erdős-Ginzburg-Ziv constant are central topics (see [51, 52, 45, 46, 23, 24, 15, 34, 35, 47, 48, 31, 36]). The associated inverse problem of $D_k(G)$ is to characterize the maximal length zero-sum sequences that can not be partitioned into $k+1$ nontrivial zero-sum subsequences. In particular, the inverse problem of $D(G)$ is to characterize the structure of minimal zero-sum subsequences of length $D(G)$, which was accomplished for groups of rank 2 in a series of papers [44] [18] [20] [47] [2], where a minimal zero-sum sequence is a zero-sum sequence that can not be partitioned into two nontrivial zero-sum subsequences. Those inverse results can be used to construct minimal generating subsets in Invariant Theory (see [11, Proposition 4.7]).

Let $\mathcal{B}(G)$ be the set of all zero-sum sequences over G . We define

$$\mathcal{M}_k(G) = \{S \in \mathcal{B}(G) : S \text{ can not be partitioned into } k+1 \text{ nontrivial zero-sum subsequences}\}.$$

Then it is easy to see that $D_k(G) = \max\{|S| : S \in \mathcal{M}_k(G)\}$. In this paper, we investigate the inverse problem of general Davenport constant $D_k(G)$ for all rank 2 groups, that is, to study the structure of sequences of $\mathcal{M}_k(G)$ of length $D_k(G)$. In 2003, Gao and Geroldinger ([18, Theorem 7.1]) studied the inverse problem of $D_k(G)$ for $G = C_n \oplus C_n$ under some assumptions of G , which had been confirmed later. We reformulate this result in the following and a proof will be given in Section 3.

Theorem 1.1. *Let $G = C_n \oplus C_n$ with $n \geq 2$, let $k \geq 1$, and let $U \in \mathcal{B}(G)$ with $|U| = D_k(G)$. Then $U \in \mathcal{M}_k(G)$ if and only if there exists a basis (e_1, e_2) of G such that it has one of the following two forms.*

(I)

$$U = e_1^{k_1 n - 1} \prod_{i \in [1, k_2 n]} (x_i e_1 + e_2), \quad \text{where}$$

- (a) $k_1, k_2 \in \mathbb{N}$ with $k_1 + k_2 = k + 1$,
- (b) $x_1, \dots, x_{k_2 n} \in [0, n - 1]$ and $x_1 + \dots + x_{k_2 n} \equiv 1 \pmod{n}$.

(II)

$$U = e_1^{an} e_2^{bn-1} (x e_1 + e_2)^{cn-1} (x e_1 + 2e_2), \quad \text{where}$$

- (a) $x \in [2, n - 2]$ with $\gcd(x, n) = 1$,
- (b) $a, b, c \geq 1$ with $a + b + c = k + 1$.

Note that in this case, we have $k \geq 2$.

For general groups, we have the following main theorem.

Theorem 1.2. *Let $G = C_{n_1} \oplus C_{n_2}$ with $1 < n_1 \mid n_2$ and $n_1 < n_2$, let $k \geq 1$, and let $U \in \mathcal{B}(G)$ with $|U| = D_k(G)$. Then $U \in \mathcal{M}_k(G)$ if and only if it has one of the following four forms.*

(I)

$$U = e_1^{\text{ord}(e_1)-1} \prod_{i \in [1, k \text{ord}(e_2)]} (x_i e_1 + e_2), \quad \text{where}$$

- (a) (e_1, e_2) is a basis for G with $\text{ord}(e_1) = n_1$ and $\text{ord}(e_2) = n_2$,
- (b) $x_1, \dots, x_{k \text{ord}(e_2)} \in [0, \text{ord}(e_1) - 1]$ and $x_1 + \dots + x_{k \text{ord}(e_2)} \equiv 1 \pmod{\text{ord}(e_1)}$.

(II)

$$U = e_1^{k \text{ord}(e_1)-1} \prod_{i \in [1, \text{ord}(e_2)]} (x_i e_1 + e_2), \quad \text{where}$$

- (a) (e_1, e_2) is a basis for G with $\text{ord}(e_1) = n_2$ and $\text{ord}(e_2) = n_1$,
- (b) $x_1, \dots, x_{\text{ord}(e_2)} \in [0, \text{ord}(e_1) - 1]$ and $x_1 + \dots + x_{\text{ord}(e_2)} \equiv 1 \pmod{\text{ord}(e_1)}$.

(III)

$$U = g_1^{n_1-1} \prod_{i \in [1, kn_2]} (-x_i g_1 + g_2), \quad \text{where}$$

- (a) (g_1, g_2) is a generating set of G with $\text{ord}(g_1) > n_1$ and $\text{ord}(g_2) = n_2$,
- (b) $x_1, \dots, x_{kn_2} \in [0, n_1 - 1]$ with $x_1 + \dots + x_{kn_2} = n_1 - 1$.

(IV)

$$U = e_1^{sn_1-1} \prod_{i \in [1, kn_2-(s-1)n_1]} ((1-x_i)e_1 + e_2), \quad \text{where}$$

- (a) (e_1, e_2) is a basis of G with $\text{ord}(e_1) = n_2$ and $\text{ord}(e_2) = n_1$,
- (b) $s \in [2, kn_2/n_1 - 1]$,
- (c) $x_1, \dots, x_{kn_2-(s-1)n_1} \in [0, n_1 - 1]$ with $x_1 + \dots + x_{kn_2-(s-1)n_1} = n_1 - 1$.

2. NOTATION AND PRELIMINARIES

Our notations and terminology are consistent with [25] and [32]. Let \mathbb{N} denote the set of positive integers and $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. For real numbers $a, b \in \mathbb{R}$, we set the discrete interval $[a, b] = \{x \in \mathbb{Z} : a \leq x \leq b\}$. Throughout this paper, all abelian groups will be written additively, and for $n \in \mathbb{N}$, we denote by C_n a cyclic group with n elements.

Let G be a finite abelian group. It is well-known that $|G| = 1$ or $G \cong C_{n_1} \oplus \dots \oplus C_{n_r}$ with $1 < n_1 \mid \dots \mid n_r \in \mathbb{N}$, where $r = r(G) \in \mathbb{N}$ is the *rank* of G , and $n_r = \exp(G)$ is the *exponent* of G . We denote by $|G|$ the *order* of G , and $\text{ord}(g)$ the *order* of $g \in G$.

Let $\mathcal{F}(G)$ be the free abelian (multiplicatively written) monoid with basis G . Then sequences over G could be viewed as elements of $\mathcal{F}(G)$. A sequence $S \in \mathcal{F}(G)$ could be written as

$$S = g_1 \cdot \dots \cdot g_l = \prod_{g \in G} g^{\mathbf{v}_g(S)},$$

where $\mathbf{v}_g(S) \in \mathbb{N}_0$ is the multiplicity of g in S . We call

- $\text{supp}(S) = \{g \in G : \mathbf{v}_g(S) > 0\} \subset G$ the *support* of S , and
- $\sigma(S) = \sum_{i=1}^l g_i = \sum_{g \in G} \mathbf{v}_g(S)g \in G$ the *sum* of S .

Let $t \in \mathbb{N}$. We denote

$$\Sigma_{\leq t}(S) = \left\{ \sum_{i \in I} g_i : I \subseteq [1, l] \text{ with } 1 \leq |I| \leq t \right\}.$$

A sequence $T \in \mathcal{F}(G)$ is called a subsequence of S if $\mathbf{v}_g(T) \leq \mathbf{v}_g(S)$ for all $g \in G$, and denoted by $T \mid S$. If $T \mid S$, then we denote

$$T^{-1}S = \prod_{g \in G} g^{\mathbf{v}_g(S) - \mathbf{v}_g(T)} \in \mathcal{F}(G).$$

Let $T_1, T_2 \in \mathcal{F}(G)$. We set

$$T_1 T_2 = \prod_{g \in G} g^{\mathbf{v}_g(T_1) + \mathbf{v}_g(T_2)} \in \mathcal{F}(G).$$

If $T_1, \dots, T_t \in \mathcal{F}(G)$ such that $T_1 \cdot \dots \cdot T_t \mid S$, where $t \geq 2$, then we say T_1, \dots, T_t are disjoint subsequences of S .

Every map of abelian groups $\phi : G \rightarrow H$ extends to a map from the sequences over G to the sequences over H by setting $\phi(S) = \phi(g_1) \cdot \dots \cdot \phi(g_l)$. If ϕ is a homomorphism, then $\phi(S)$ is a zero-sum sequence if and only if $\sigma(S) \in \ker(\phi)$.

We denote by $\mathbf{E}(G)$ the Gao's constant which is the smallest integer ℓ such that every sequence over G of length ℓ has a zero-sum subsequence of length $|G|$ and by $\eta(G)$ the smallest integer ℓ such that every sequence over G of length ℓ has a zero-sum subsequence T of length $1 \leq |T| \leq \exp(G)$. Let $\mathbf{d}(G)$ be the maximal length of a sequence over G that has no zero-sum subsequence. Then it is easy to see that $\mathbf{d}(G) = \mathbf{D}(G) - 1$. The following result is well-known and we may use it without further mention.

Lemma 2.1. *Let G be a finite abelian group. Then $\mathbf{E}(G) = |G| + \mathbf{d}(G) \leq 2|G| - 1$.*

Proof. **TOPROVE 0** □

We also need the following lemmas.

Lemma 2.2. *Let G be a finite abelian group. If $\mathbf{D}(G) = |G|$, then G is cyclic and for every minimal zero-sum sequence S over G of length $|G|$, there exists $g \in G$ with $\text{ord}(g) = |G|$ such that $S = g^{|G|}$.*

Proof. **TOPROVE 1** □

Lemma 2.3. *Let G be a finite abelian group and let $H \subset G$ be a proper subgroup. Then $\mathbf{D}_k(H) < \mathbf{D}_k(G)$ for all $k \in \mathbb{N}$.*

Proof. **TOPROVE 2** □

Theorem 2.4. *Let $G = C_{n_1} \oplus C_{n_2}$ with $n_1 \mid n_2$, where $n_1, n_2 \in \mathbb{N}$, and let $k \in \mathbb{N}$. Then $\eta(G) = 2n_1 + n_2 - 2$ and $D_k(G) = n_1 + kn_2 - 1$. In particular, $D(G) = n_1 + n_2 - 1$.*

Proof. **TOPROVE 3** □

Theorem 2.5. *Let $G = C_n \oplus C_{mn}$ with $n \geq 2$ and $m \geq 1$. A sequence S over G of length $D(G) = n + mn - 1$ is a minimal zero-sum sequence if and only if it has one of the following two forms:*

(I)

$$S = e_1^{\text{ord}(e_1)-1} \prod_{i=1}^{\text{ord}(e_2)} (x_i e_1 + e_2),$$

where

(a) $\{e_1, e_2\}$ is a basis of G ,

(b) $x_1, \dots, x_{\text{ord}(e_2)} \in [0, \text{ord}(e_1) - 1]$ and $x_1 + \dots + x_{\text{ord}(e_2)} \equiv 1 \pmod{\text{ord}(e_1)}$.

In this case, we say that S is of type I(a) or I(b) according to whether $\text{ord}(e_2) = n$ or $\text{ord}(e_2) = mn > n$.

(II)

$$S = f_1^{sn-1} f_2^{(m-s)n+\epsilon} \prod_{i=1}^{n-\epsilon} (-x_i f_1 + f_2),$$

where

(a) $\{f_1, f_2\}$ is a generating set for G with $\text{ord}(f_2) = mn$ and $\text{ord}(f_1) > n$,

(b) $\epsilon \in [1, n-1]$ and $s \in [1, m-1]$,

(c) $x_1, \dots, x_{n-\epsilon} \in [1, n-1]$ with $x_1 + \dots + x_{n-\epsilon} = n-1$,

(d) either $s = 1$ or $nf_1 = nf_2$, with both holding when $m = 2$, and

(e) either $\epsilon \geq 2$ or $nf_1 \neq nf_2$.

In this case, we say that S is of type II.

Proof. **TOPROVE 4** □

Lemma 2.6. *Let G be a finite abelian group, let H be a cyclic subgroup of G , and let $\varphi: G \rightarrow G/H$ be the canonical epimorphism. If $S \in \mathcal{M}_k(G)$, then $\varphi(S) \in \mathcal{M}_{k|H|}(G/H)$.*

Proof. **TOPROVE 5** □

3. PROOF OF MAIN THEOREMS

Proposition 3.1. *Let G be a finite abelian group of rank at most 2, let $k \in \mathbb{N}$, and let S be a zero-sum sequence over G of length $D_k(G)$. Then $S \in \mathcal{M}_k(G)$ if and only if $0 \notin \Sigma_{\leq \exp(G)-1}(S)$.*

Proof. **TOPROVE 6** □

We first investigate the associated inverse problem for cyclic groups.

Theorem 3.2. *Let G be cyclic, let $k \in \mathbb{N}$, and let S be a zero-sum sequence over G of length $D_k(G)$. Then $S \in \mathcal{M}_k(G)$ if and only if there exists $g \in G$ with $\text{ord}(g) = |G|$ such that $S = g^{k|G|}$.*

Proof. **TOPROVE 7** □

Next, we prove Theorem 1.1 which could be handled easily by Proposition 3.1 and [18, Theorem 7.1].

Proof. **TOPROVE 8** □

Lemma 3.3. *Let $G = C_n \oplus C_n$ with $n \geq 2$ and let $k \geq 2$. If $S \in \mathcal{F}(G)$ is a zero-sum sequence with $|S| = (k+1)n - 1$ and $0 \notin \Sigma_{\leq n-1}(S)$, then there is a basis (e_1, e_2) for G such that either*

1. $\text{supp}(S) \subseteq \{e_1\} \cup (\langle e_1 \rangle + e_2)$ and $v_{e_1}(S) \equiv -1 \pmod{n}$, or
2. $S = e_1^{an} e_2^{bn-1} (xe_1 + e_2)^{cn-1} (xe_1 + 2e_2)$ for some $x \in [2, n-2]$ with $\gcd(x, n) = 1$, and some $a, b, c \geq 1$ with $k+1 = a + b + c$.

Proof. **TOPROVE 9** □

The following lemma shows two special cases of Theorem 1.2.

Lemma 3.4. *Let $G = C_{n_1} \oplus C_{n_2}$ with $1 < n_1 \mid n_2$ and $n_1 < n_2$, let $k \geq 2$, and let $U \in \mathcal{M}_k(G)$ with $|U| = D_k(G)$.*

1. *If there is some $e_1 \in \text{supp}(U)$ such that $\text{ord}(e_1) = n_1$ and $v_{e_1}(U) \geq n_1 - 1$, then there exists $e_2 \in G$ with $\text{ord}(e_2) = n_2$ such that (e_1, e_2) is a basis of G and*

$$U = e_1^{n_1-1} \prod_{i \in [1, kn_2]} (x_i e_1 + e_2),$$

where $x_1, \dots, x_{kn_2} \in [0, n_1 - 1]$ and $x_1 + \dots + x_{kn_2} \equiv 1 \pmod{n_1}$.

2. *If there is some $e_2 \in \text{supp}(U)$ such that $\text{ord}(e_2) = n_2$ and $v_{e_2}(U) \geq kn_2 - 1$, then there exists $e_1 \in G$ with $\text{ord}(e_1) = n_1$ such that (e_1, e_2) is a basis of G and*

$$U = e_2^{kn_2-1} \prod_{i \in [1, n_1]} (e_1 + x_i e_2),$$

where $x_1, \dots, x_{n_1} \in [0, n_2 - 1]$ and $x_1 + \dots + x_{n_1} \equiv 1 \pmod{n_2}$.

Proof. **TOPROVE 10** □

Now we are ready to prove our main Theorem 1.2.

Proof. **TOPROVE 11** □

REFERENCES

- [1] Sukumar Das Adhikari, David J. Gryniewicz, and Zhi-Wei Sun, *On weighted zero-sum sequences*, Advances in Applied Mathematics **48**(2012), 506-527.
- [2] G. Bhowmik, I. Halupczok, and J. Schlage-Puchta, *The structure of maximal zero-sum free sequences*, Acta Arith. **143** (2010), 21-50.
- [3] G. Bhowmik, I. Halupczok, and J. Schlage-Puchta, *Inductive methods and zero-sum free sequences*, Integers, **9**(2009), 515-536.
- [4] G. Bhowmik and J. Schlage-Puchta, *Davenport's constant for groups of the form $\mathbb{Z}_3 \oplus \mathbb{Z}_3 \oplus \mathbb{Z}_{3d}$* , In *Additive combinatorics*, volume 43 of *CRM Proc. Lecture Notes*, pages 307–326. Amer. Math. Soc., Providence, RI, 2007.
- [5] J. Bitz, S. Griffith, and X. He, *Exponential lower bounds on the generalized Erdős–Ginzburg–Ziv constant*, Discrete Mathematics **343**(2020), 112083.
- [6] F.E. Brochero Martínez and S. Ribas, *Extremal product-one free sequences in Dihedral and Dicyclic Groups*, Discrete Math. **341** (2018), 570 – 578.
- [7] Y. Caro, *Remarks on a zero-sum theorem*, J. Comb. Theory Ser. A **76** (1996), 315 – 322.
- [8] K. Csiszter and M. Domokos, *On the generalized Davenport constant and the Noether number*, Central European J. Math. **11** (2013), 1605 – 1615.
- [9] K. Csiszter and M. Domokos, *Groups with large Noether bound*, Ann. Inst. Fourier (Grenoble) **64** (2014), 909 – 944.
- [10] K. Csiszter and M. Domokos, *The Noether number for the groups with a cyclic subgroup of index two*, J. Algebra **399** (2014), 546 – 560.
- [11] K. Csiszter, M. Domokos, and A. Geroldinger, *The interplay of invariant theory with multiplicative ideal theory and with arithmetic combinatorics*, Multiplicative Ideal Theory and Factorization Theory, Springer, 2016, pp. 43 – 95.
- [12] K. Csiszter, M. Domokos, and I. Szöllősi, *The Noether numbers and the Davenport constants of the groups of order less than 32*, J. Algebra **510** (2018), 513 – 541.
- [13] Ch. Delorme, O. Ordaz, and D. Quiroz, *Some remarks on Davenport constant*, Discrete Math. **237**(2001), 119-128.
- [14] Y. Edel, C. Elsholtz, A. Geroldinger, S. Kubertin, and L. Rackham, *Zero-sum problems in finite abelian groups and affine caps*, Quart. J. Math. **58** (2007), 159 – 186.
- [15] J. Ebert and D.J. Gryniewicz, *Structure of a sequence with prescribed zero-sum subsequences: rank two p -groups*, European J. Combin. **118** (2024), Paper No. 103888, 13pp.
- [16] Y. Fan, W. Gao, and Q. Zhong, *On the Erdős–Ginzburg–Ziv constant of finite abelian groups of high rank*, J. Number Theory **131** (2011), 1864 – 1874.
- [17] M. Freeze and W. Schmid, *Remarks on a generalization of the Davenport constant*, Discrete mathematics **310**(2010), 3373-89.
- [18] W. Gao and A. Geroldinger, *On zero-sum sequences in $\mathbb{Z}/n\mathbb{Z} \oplus \mathbb{Z}/n\mathbb{Z}$* , Integers **3** (2003), Paper A08, 45p.
- [19] W. Gao and A. Geroldinger, *Zero-sum problems in finite abelian groups: a survey*, Expo. Math. **24**(2006), 337-369.
- [20] Weidong Gao, A. Geroldinger, D. J. Gryniewicz, *Inverse zero-sum problems III*, Acta Arith. **141**(2010), 103–152.
- [21] W. Gao, D. Han, J. Peng, and F. Sun, *On zero-sum subsequences of length $k \exp(G)$* , J. Combin. Theory Ser. A **125**(2014), 240-53.
- [22] W. Gao and Z. Lu, *The Erdős–Ginzburg–Ziv theorem for dihedral groups*, J. Pure Appl. Algebra **212** (2008), 311 – 319.

- [23] W. Gao, J. Peng, and G. Wang, *Behaving sequences*, J. Comb. Theory, Ser. A **118** (2011), 613 – 622.
- [24] F. Ge, *Solution to the index conjecture in zero-sum theory*, J. Combin. Theory Ser. A **180** (2021), 105410.
- [25] A. Geroldinger, *Additive group theory and non-unique factorizations*, Combinatorial Number Theory and Additive Group Theory (A. Geroldinger and I. Ruzsa, eds.), Advanced Courses in Mathematics CRM Barcelona, Birkhäuser, 2009, pp. 1–86.
- [26] A. Geroldinger, D.J. Grynkiewicz, and P. Yuan, *On products of k atoms II*, Mosc. J. Comb. Number Theory **5** (2015), 73 – 129.
- [27] A. Geroldinger and F. Halter-Koch, *Non-Unique Factorizations*, Algebraic, Combinatorial and Analytic Theory, Pure and Applied Mathematics, vol. 278, 700p, Chapman & Hall/CRC, 2006.
- [28] B. Girard. *On the existence of distinct lengths zero-sum subsequences*, Rocky Mountain J. Math. **42**(2012), 583-596.
- [29] B. Girard, *An asymptotically tight bound for the Davenport constant*, Journal de l'École polytechnique—Mathématiques **5** (2018), 605-611.
- [30] B. Girard and W.A. Schmid, *Direct zero-sum problems for certain groups of rank three*, J. Number Theory **197** (2019), 297 – 316.
- [31] B. Girard and W.A. Schmid, *Inverse zero-sum problems for certain groups of rank three*, Acta Math. Hungar. **160**(2020), 229-247.
- [32] D. J. Grynkiewicz, *Structural Additive Theory*, Developments in Mathematics, 30, Springer, 2013, 426 pp.
- [33] D. J. Grynkiewicz, *Inverse Zero-Sum Problems III: Addendum*, <https://arxiv.org/abs/2107.10619>.
- [34] D.J. Grynkiewicz and C. Liu, *A multiplicative property for zero-sums I*, Discrete Math. **345** (2022), Paper No. 112974, 21pp.
- [35] D.J. Grynkiewicz and C. Liu, *A multiplicative property for zero-sums II*, Electron. J. Combin. **29** (2022), Paper No. 3.12, 16pp.
- [36] D. J. Grynkiewicz and U. Vishne, *The index of small length sequences*, Internat. J. Algebra Comput. **30**(2020), 977–1014.
- [37] F. Halter-Koch, *A generalization of Davenport's constant and its arithmetical applications*, Colloq. Math. **63**(1992), 203-210.
- [38] Dongchun Han and Hanbin Zhang, *Erdős-Ginzburg-Ziv Theorem and Noether number for $C_m \rtimes_{\varphi} C_{mn}$* , J. Number Theory **198** (2019), 159 – 175.
- [39] S. Hong and K. Zhao, *On the zero-sum subsequences of modular restricted lengths*, Discrete Mathematics **347**(2024), 113967.
- [40] E. Naslund, *Exponential bounds for the Erdős-Ginzburg-Ziv constant*, J. Combin. Theory Ser. A **174** (2020), 105185.
- [41] C. Liu, *On the lower bounds of Davenport constant*, J. Combin. Theory Ser. A **171**(2020), 105162.
- [42] A. Plagne and W. Schmid. *An application of coding theory to estimating Davenport constants*, Designs, Codes and Cryptography **61**(2011), 105-118.
- [43] Y. Qu, Y. Li, and D. Teeuwsen, *On a conjecture of the small Davenport constant for finite groups*, J. Combin. Theory Ser. A **189**(2022), 105617.
- [44] C. Reiher, *A proof of the theorem according to which every prime number possesses property B*, PhD Thesis, Rostock, 2010.
- [45] S. Savchev and F. Chen, *Long zero-free sequences in finite cyclic groups*, Discrete Math. **307** (2007), 2671 – 2679.
- [46] S. Savchev and F. Chen, *Long minimal zero-sum sequences in the group $C_2 \oplus C_{2k}$* , Integers **12** (2012), Paper A51, 18pp.
- [47] W.A. Schmid, *Inverse zero-sum problems II*, Acta Arith. **143** (2010), 333 – 343.

- [48] W.A. Schmid, *Restricted inverse zero-sum problems in groups of rank two*, Q. J. Math. **63** (2012), 477 – 487.
- [49] A. Sidorenko, *On generalized Erdős–Ginzburg–Ziv constants for $Z2d$* , J. Combin. Theory Ser. A **174**(2020), 105254.
- [50] J. E. Olson, *A combinatorial problem on finite Abelian groups II*, J. Number Theory **1**(1969), 195-199.
- [51] P. Yuan, *On the index of minimal zero-sum sequences over finite cyclic groups*, J. Comb. Theory, Ser. A **114** (2007), 1545 – 1551.
- [52] P. Yuan and X. Zeng, *Indexes of long zero-sum free sequences over cyclic groups*, European J. Combin. **32** (2011), 1213 – 1221.

UNIVERSITY OF GRAZ, NAWI GRAZ, DEPARTMENT OF MATHEMATICS AND SCIENTIFIC COMPUTING, HEINRICHSTRASSE 36, 8010 GRAZ, AUSTRIA

Email address: qinghai.zhong@uni-graz.at

URL: <https://imsc.uni-graz.at/zhong/>