

Reducing Stochastic Games to Semidefinite Programming

Manuel Bodirsky^{*} Georg Loho[†] Mateusz Skomra[‡]

Abstract

We present a polynomial-time reduction from max-average constraints to the feasibility problem for semidefinite programs. This shows that Condon’s simple stochastic games, stochastic mean payoff games, and in particular mean payoff games and parity games can all be reduced to semidefinite programming.

1 Introduction

There are two important clusters of computational problems that are not known to be in the complexity class P: the first is related to numeric computation, and contains the sums-of-square-roots problem [GGJ76, EHS24], the Euclidean shortest path problem, PosSLP [ABKPM09], and the feasibility problem for semidefinite programs [NN94, Ram97]. The second cluster is related to tropical geometry, and contains for instance the model checking problem for the propositional μ -calculus [EJ91, EJS93], parity games [Mar75, Mos91], mean payoff games [EM79], and/or scheduling [MSS04], stochastic mean payoff games [AM09], and simple stochastic games [Con92]. So far, no polynomial-time reduction from a problem of one of the clusters to a problem from the other cluster was known. We show that all of the mentioned problems from the second cluster can be reduced in polynomial time to the feasibility problem for semidefinite programs from the first cluster.

Semidefinite programming is a generalisation of linear programming with many important algorithmic applications in discrete and continuous optimisation, both from a theoretical and a practical perspective [GLS94, BTN01, Las09]. However, already the feasibility problem for semidefinite programs is not known to be in P. By Ramana’s duality [Ram97], the problem is either in the intersection of NP and coNP, or outside of the union of NP and coNP. The semidefinite feasibility problem falls into the existential theory of the reals, which is known to be in PSPACE [Can88]. However, it has been observed by Khachiyan that the smallest feasible solution to an SDP might be of doubly exponential size (see, e.g., [PK97]), which is an obstacle for the polynomial-time algorithmic methods known for linear programming. In fact, it is possible to reduce the *PosSLP problem* to semidefinite programming [TV08]. PosSLP has been introduced in [ABKPM09], motivated by the ‘generic task of numerical computation’. For example, the sums-of-square-roots problem, which is a numeric computational problem not even known to be in NP and the barrier for polynomial-time tractability for numerous problems in computational geometry, such as the Euclidean shortest path problem, has a polynomial-time reduction to PosSLP [ABKPM09].

Mean payoff games are turn-based two-player games on graphs [EM79, ZP96]; it is a famous open problem in theoretical computer science whether there is a polynomial-time algorithm to

^{*}Institut für Algebra, TU Dresden, Germany.

[†]FU Berlin, Germany & U Twente, The Netherlands.

[‡]LAAS-CNRS, Université de Toulouse, CNRS, Toulouse, France.




Figure 1: Computational problems that are not known to be in P and not known to be NP-hard, and their relationship. Arcs indicate polynomial-time (many-one) reductions, the dotted arc indicates a polynomial-time Turing reduction.

decide for a given graph which of the players has a winning strategy. Finding the winning region in mean payoff games is polynomial-time equivalent to various other computational problems in theoretical computer science, for instance scheduling with and-or-constraints [MSS04], the max-atoms problem [BNRC08], as well as solvability of max-plus systems and tropical linear feasibility [ABG07, AGG12]. The latter can be seen as the tropical analog of testing feasibility of linear inequalities (which for classical geometry is known to be in P, e.g. via the ellipsoid method [Kha79, GLS94]).

Furthermore, there is a simple reduction from *parity games* to mean payoff games by using the priorities of the parity game with a suitably large basis [Pur95]. Parity games are polynomial-time equivalent to the model-checking problem of the propositional μ -calculus [EJS93], which has been called ‘*the most important logics in model checking*’ [BW18]. For parity games, a quasipolynomial algorithm has been found recently [CJK⁺22]. Subsequent work indicates that the various quasipolynomial methods that have been obtained lately for parity games [LPSW22, JMT22, LB20] do not extend to mean payoff games [CFG022].

Mean payoff games can be generalised to *stochastic mean payoff games* (sometimes called *2 $\frac{1}{2}$ -player games*), where the graph might contain stochastic nodes. If the strategy of one of the two players in such a game is fixed, the game turns into a Markov decision process, for which polynomial-time algorithms based on linear programming are known [Put05]. Stochastic mean payoff games are equivalent under polynomial-time Turing reductions to Condon’s *simple stochastic games* [AM09], which are known to be in the intersection of NP and coNP [Con92] (we may even assume that the simple stochastic games are *stopping*, see the discussion in Section 2.2.) There are many other variants of games that all have the same complexity, see [AM09]. Stochastic mean payoff games can also be reduced to max-plus-average constraints, which are still in $\text{NP} \cap \text{coNP}$ [BM18].

We introduce a fragment of max-plus-average constraints, which we call *max-average constraints*. There is a polynomial-time reduction from stopping simple stochastic games to max-average constraints (Section 3). Max-average constraints have the advantage that they can be further reduced to the feasibility problem for semidefinite programs. This implies that all the mentioned computational problems from the second cluster can be reduced to semidefinite programming. See Figure 1 for an overview of the mentioned computational problems and their relationship.

The reduction uses a similar idea as the reduction of Schewe [Sch09] to reduce parity games and mean payoff games to linear programming, which is, however, not a proper polynomial-time reduction. The idea is to replace constraints of the form $x_0 = \max(x_1, \dots, x_k)$ by $x_0 = \log_b(b^{x_1} + \dots + b^{x_k})$ for a sufficiently large b . Following [AGS18], we extend this idea to max-average constraints and obtain a proper polynomial-time reduction to non-Archimedean SDPs. From there, we use bounds from quantifier elimination results to translate non-Archimedean SDPs to real SDPs. However, we still have the obstacle that these SDPs involve coefficients of doubly exponential size, which means that they have exponential representation size and do not lead to a polynomial-time reduction. We overcome this difficulty by using the duality of semidefinite programming to find small expressions that define coefficients of doubly exponential size, which combined with the above ideas leads to a polynomial-time reduction from max-average constraints to (real) SDPs.

Our results imply that if there is a polynomial-time algorithm for SDPs, then all the mentioned problems in the second cluster can be solved in polynomial time as well. Conversely, our reduction can be used to translate interesting families of instances for (stochastic) mean payoff games or parity games into SDP instances, which might yield interesting families of instances for (algorithmic approaches to) semidefinite programming. For linear programming, this idea turned out to be very fruitful: certain instances of parity games have been used to prove exponential lower bounds for several famous pivoting rules for the simplex method in linear programming [FHZ11, Fri11, FHZ14, AF17, DFH23]. However, these results were not based on a general polynomial-time reduction from parity games to linear programming (which is not known!). We also hope that algorithmic ideas for solving games might generalise to (certain classes of) SDPs.

Our results also show that two large research communities (convex real algebraic geometry, and more generally convex optimisation on the one hand, and verification and automata theory on the other hand) were working on closely related computational problems and might profit from joining forces in the future.

1.1 Related work

There are multiple works that reduce games to continuous optimization problems. As mentioned above, one-player mean payoff games (Markov decision processes) can be reduced to linear programming [Put05], which implies that they can be solved in polynomial time. For the case of two-player deterministic mean payoff games, Schewe [Sch09] proposed a reduction to linear programs with coefficients of exponential representation size. Such programs can also be interpreted as linear programs over non-Archimedean ordered fields [ABGJ15, ABGJ14, ABGJ21]. The idea of Schewe was generalized to stochastic games in [BEGM17], where the authors prove that stochastic mean payoff games can be encoded by convex programs with constraints of exponential encoding size. Such programs can be solved by the ellipsoid method in pseudo-polynomial time provided that the number of stochastic nodes is fixed. The non-Archimedean approach was generalized from deterministic games to stochastic games in the work [AGS18], which proves that stochastic mean payoff games can be reduced to semidefinite programs over non-Archimedean fields. The paper [AGS18] serves as a basis for our results.

These reductions express mean payoff games as convex optimization problems, but these problems are “non-standard” in the sense that they either use constraints of exponential encoding size or are expressed over non-Archimedean fields. There is yet another line of work deriving polynomial-time reductions from mean payoff games to classical optimization problems which are not convex. In particular, [Con93] expressed simple stochastic games as non-convex quadratic programs. This reduction is further studied in [KRSW22]. Another reduction to (generalized) complementarity problems with P -matrices was proposed in [GR05, SV06] and extended in sub-

sequent works [JS08, FJS10, HIJ13]. Such linear complementarity problems form a subclass of quadratic programs and belong to the complexity class UniqueEOPL [FGMS20]. However, they are in general non-convex and can therefore not be expressed by semidefinite programs.

2 Preliminaries

The set of natural numbers $\{0, 1, 2, \dots\}$ is denoted by \mathbb{N} , and the set of real numbers by \mathbb{R} . We assume familiarity with first-order logic; see, e.g., [Hod97]. Let τ be a *signature*, i.e., a set of function and relation symbols. A first-order τ -formula is a formula build from the Boolean connectives, atomic τ -formulas and quantification in the usual way. If A is a τ -structure and ϕ is a τ -formula with free variables x_1, \dots, x_k , then ϕ *defines* over A the relation

$$\{(a_1, \dots, a_k) \in A^k \mid A \models \phi(a_1, \dots, a_k)\}.$$

A set is called *semialgebraic* if it has a first-order definition over the field $(\mathbb{R}; +, \cdot)$, allowing real parameters. A first-order formula $\phi(x_1, \dots, x_k)$ with free variables x_1, \dots, x_k is called *primitive positive* if it is of the form

$$\exists y_1, \dots, y_\ell (\psi_1 \wedge \dots \wedge \psi_m)$$

where ψ_1, \dots, ψ_m are atomic formulas over the variables $x_1, \dots, x_k, y_1, \dots, y_\ell$. A relation is called *primitive positive definable in a τ -structure A* if there exists a primitive positive τ -formula which defines the relation over A .

2.1 Semidefinite Programming

For a symmetric matrix $S \in \mathbb{R}^{m \times m}$ we write $S \succeq 0$ if S is *positive semidefinite*, i.e., if $x^\top S x \geq 0$ for all $x \in \mathbb{R}^m$, and $S \succ 0$ if it is *positive definite*, i.e., if $x^\top S x > 0$ for all $x \in \mathbb{R}^m \setminus \{0\}$. A *semidefinite program (SDP)* is an optimisation problem of the following form:

$$\begin{aligned} \inf \quad & \sum_{i=1}^n c_i x_i \\ \text{such that} \quad & \sum_{i=1}^n x_i A_i - B \succeq 0, \end{aligned} \tag{P}$$

where $A_1, \dots, A_n, B \in \mathbb{R}^{m \times m}$ are symmetric and $c_1, \dots, c_n \in \mathbb{R}$. If the data $A_1, \dots, A_n, B, c_1, \dots, c_n$ is rational, then the problem of deciding whether the supremum exists, and if so, to compute the supremum, is a computational problem known as the *semidefinite programming problem*. The *representation size* of the SDP then consists of the sum of the bit lengths of all the rational numbers in the input.

The expression $\sum_{i=1}^n x_i A_i - B \succeq 0$ in (P) is called a *linear matrix inequality (LMI)*, and defines the *feasible region* of the SDP, which is called a *spectrahedron*. The representation size of LMIs is defined analogously as for SDPs. We say that an SDP is *strictly feasible* if $\sum_{i=1}^n x_i A_i - B \succ 0$ has a solution $x \in \mathbb{R}^n$.

Definition 2.1. Given symmetric matrices $A_1, \dots, A_n, B \in \mathbb{Q}^{m \times m}$ the *feasibility problem for semidefinite programs* asks to decide whether there is a solution $x \in \mathbb{R}^n$ to an LMI

$$x_1 A_1 + \dots + x_n A_n - B \succeq 0 \ .$$

Note that spectrahedra are convex and semialgebraic. It has been conjectured that every convex semialgebraic set is the projection of a spectrahedron (also called a *spectrahedral shadow*), but this turned out to be false [Sch18] (see [BKT24] for an alternative proof of an even stronger inexpressibility result). It is easy to see that the set of spectrahedral shadows is closed under primitive positive definability, and that for every primitive positive formula ϕ we may find an LMI for the spectrahedral shadow defined by ϕ whose representation size is bounded by the sum of the representation sizes of conjuncts of ϕ . We also note that the definition of a semidefinite program makes sense over any real closed field, not only over the real numbers. In particular, it is meaningful to consider semidefinite programs over non-Archimedean real closed fields, such as the field of Puiseux series which we discuss in Section 4.

The (*Frobenius*) *inner product* of two matrices $S, T \in \mathbb{R}^{m \times m}$ is defined as $\langle S, T \rangle := \text{trace}(ST) = \sum_{i,j} S_{ij}T_{ij}$. The *dual program* to the SDP from (P) (to which we refer as the *primal*) is given by

$$\begin{aligned} & \sup_{Y \succeq 0} \langle B, Y \rangle \\ & \text{such that } \langle A_i, Y \rangle = c_i \text{ for every } i \in \{1, \dots, n\}. \end{aligned} \tag{D}$$

It is well-known that the value of the primal SDP is bounded from below by the value of the dual; however, the lower bound might not be tight, i.e., there might be a non-zero *duality gap*. Moreover, both primal and dual might not have optimal solutions, i.e., the supremum may not be a maximum and the infimum might not be a minimum. See [Ram97] for a more complicated, but exact dual. However, for us the following sufficient condition for a vanishing duality gap will be enough.

Theorem 2.2 (see Theorem 2.4.1 in [BTN01]). *If the primal SDP is strictly feasible and bounded from below, then the dual program attains its supremum and there is no duality gap.*

2.2 Stochastic Games

A *stochastic mean payoff game* is a two-player game played on a finite edge-weighted directed graph $\vec{G} = (V, E, r)$, i.e., V is a finite set of nodes, $E \subseteq V^2$ is a set of edges, and $r: E \rightarrow \mathbb{Q}$ is a *weight function*. The nodes of the graph are partitioned into three subsets, $V = V_{\text{Min}} \sqcup V_{\text{Max}} \sqcup V_{\text{Rand}}$. The nodes in V_{Min} are controlled by player Min, the nodes in V_{Max} are controlled by player Max, and the nodes in V_{Rand} are controlled by nature. The players play the game by moving a pawn on the graph. When the pawn is on a node $v \in V_{\text{Min}}$, player Min chooses an outgoing edge (v, u) and moves the pawn to u . Analogously, player Max chooses the next move when the pawn is on a node $v \in V_{\text{Max}}$. When the pawn is on a node $v \in V_{\text{Rand}}$, the next move is chosen by nature randomly, according to some probability distribution on the outgoing edges of v . This probability distribution is known to both players and does not change during the game. To ensure that players can always make a move, we suppose that all nodes of the graph have at least one outgoing edge. The weights of the edges of the graph represent payoffs that player Max receives from player Min after each move of the pawn. In order to state the objectives of both players, we first specify the strategies that they are allowed to use. A *pure and positional* strategy (also called a *policy*) for player Min is a function $\sigma: V_{\text{Min}} \rightarrow V$ that to each node controlled by Min associates a neighbor, $(v, \sigma(v)) \in E$. Player Min plays according to σ if they always move the pawn to $\sigma(v)$ when the pawn lands on v . We analogously define a policy $\tau: V_{\text{Max}} \rightarrow V$ of player Max. Note that if we fix a pair of policies and an initial position of the pawn, then the resulting movement of the pawn is a Markov chain on the graph (V, E) . Given an initial position $u \in V$ of the pawn and a pair of policies (σ, τ) , the

mean payoff reward of player Max is the quantity

$$g(\sigma, \tau)_u := \lim_{N \rightarrow \infty} \frac{1}{N} \mathbb{E}_{\sigma, \tau}^u (r_{u_0 u_1} + \cdots + r_{u_{N-1} u_N}),$$

where $u = u_0, u_1, \dots, u_N$ is the path of the pawn in the first N turns of the game and the expectation is taken with respect to the Markov chain obtained by fixing (σ, τ) . The ergodic theorem of Markov chains shows that $g(\sigma, \tau)_u$ is well-defined. The objective of player Max is to maximize the mean payoff reward while the objective of player Min is to minimize this reward. It is known that both players in a stochastic mean payoff game have optimal policies [LL69]. In other words, there exist policies (σ^*, τ^*) and a vector $\chi \in \mathbb{R}^V$ such that for every u and every (σ, τ) we have the inequality

$$g(\sigma^*, \tau)_u \leq \chi_u \leq g(\sigma, \tau^*)_u.$$

The reward $\chi = g(\sigma^*, \tau^*)$ is called the *value* of the game. In order to make these games computationally tractable, we suppose that the weights and the probability distributions at nodes in V_{Rand} are rational. The nodes of the graph are also called the *states* of the game.

A *simple stochastic game* is a special case of a stochastic mean payoff game. In this game, every node in V_{Rand} has out-degree exactly two and the probability distribution at every such node is $(1/2, 1/2)$. Furthermore, the graph is equipped with two special terminal states called Win and Lose. These states are absorbing, i.e., Win has only one outgoing edge that goes back to Win and Lose has only one outgoing edge that goes back to Lose. The weight of the edge going from Win to Win is equal to 1, while the weight of all of the other edges of the graph is equal to 0. In this case, the mean payoff reward of player Max is simply the probability that the pawn reaches the winning state, $g(\sigma, \tau)_u = \mathbb{P}_{\sigma, \tau}^u(\text{Reach}(\text{Win}))$. A simple stochastic game is called *stopping* if for every initial position $u \in V$ and every pair of policies (σ, τ) , the pawn reaches a terminal state with probability one, $\mathbb{P}_{\sigma, \tau}^u(\text{Reach}(\text{Win}, \text{Lose})) = 1$. In other words, the game is stopping if the Markov chain obtained by fixing any pair of policies has only two recurrent classes, namely $\{\text{Win}\}$ and $\{\text{Lose}\}$.

The computational problem associated with stochastic mean payoff games is the following: given a stochastic mean payoff game, find its value and a pair of optimal policies. In [AM09], Andersson and Miltersen proved that this problem is polynomial-time (Turing) reducible to the problem of finding the value of a stopping simple stochastic game. By a binary search argument, the problem of computing the value of a stopping simple stochastic game can be further reduced to the decision problem, in which we want to decide whether the value of a given state of such a game is at least $1/2$. The details of this reduction can be found, e.g., in [AGS18] or [Sko18, Section 7.1].

Theorem 2.3 ([AGS18, Lemma 48]). *The problem of computing the value and a pair of optimal policies of a stochastic mean payoff game is polynomial-time (Turing) reducible to the problem of deciding whether a given state of a stopping simple stochastic game has value at least $1/2$.*

3 From stochastic games to max-average constraints

In this section, we focus on the decision problem stated in Theorem 2.3 above: given a stopping simple stochastic game we want to decide whether the value of a given state is at least $1/2$. This problem can be reduced to max-average constraint satisfaction problems. The ideas for this reduction are not new, but we are not aware of an explicit reference for this fact, and we therefore present a short proof in this section. In the next section, we will show that max-average CSPs then further reduce to the feasibility problem of non-Archimedean semidefinite programs. The

fact that stochastic games can be encoded as non-Archimedean SDPs was proven in [AGS18] and our reduction is very similar. While their approach relies on reducing stochastic games to ergodic games, we omit this step and obtain a slightly more straightforward reduction.

Definition 3.1. An instance of a *max-average constraint satisfaction problem* is a conjunction of constraints, each of which has one of the following forms:

1. $x_0 \leq \max(x_1, \dots, x_k)$ where $k \in \mathbb{N}$,
2. $x_0 \leq \frac{x_1 + x_2}{2}$,
3. $x_0 = c$ where c is a rational constant.

The computational task is to determine whether such a conjunction has a solution over $\mathbb{Q} \cup \{-\infty\}$ (equivalently, over $\mathbb{R} \cup \{-\infty\}$).

Note that max-average CSP instances may also use constraints of the form $x_0 \leq \min(x_1, \dots, x_k)$ where $k \in \mathbb{N}$, $x_0 \leq c$, or $x_0 \geq c$ for some rational constant c , because the respective relations are primitively positively definable over the other relations.

We now show that the problem of solving stochastic mean payoff games can be reduced to max-average CSPs. A reduction to slightly more general CSPs, including constraints of the form $x_0 \leq c + x_1$, follows from [AGS18, Theorem 18] and is based on a Collatz–Wielandt property of order-preserving and additively homogeneous maps. However, for our purposes, it is useful that the reduction does not use these additional constraints. Instead, we give a proposition that characterizes the value of a stopping simple stochastic game. It was already proven in [Con92] that this value is a unique fixed point of an operator involving min, max, and averaging operations. By replacing the fixed point equations with inequalities, we obtain a set of certificates which prove that the value is at least $1/2$. This observation is proven, e.g., in [BBDFS23] and gives inequalities very similar to the inequalities on Shapley operators used in [AGS18].

Proposition 3.2. Consider a stopping simple stochastic game played on a directed graph $\vec{\mathcal{G}} = ([n], E)$. Then, the value of a state $k \in [n]$ is at least $1/2$ if and only if the following system of inequalities (an instance of a max-average constraint satisfaction problem) has a solution $x \in \mathbb{Q}_{\geq 0}^n$:

$$\begin{aligned}
x_i &\leq \max_{(i,j) \in E} \{x_j\} \quad \text{for every } i \in V_{\text{Max}}, \\
x_i &\leq \min_{(i,j) \in E} \{x_j\} \quad \text{for every } i \in V_{\text{Min}}, \\
x_i &\leq \frac{1}{2} \sum_{(i,j) \in E} x_j \quad \text{for every } i \in V_{\text{Rand}}, \\
x_{\text{Lose}} &= 0, \\
x_{\text{Win}} &= 1, \\
x_k &\geq 1/2.
\end{aligned} \tag{1}$$

Proof. **TOPROVE 0** □

4 From max-average constraints to non-Archimedean SDPs

By “lifting” the inequalities (1) to Puiseux series, we get the reduction from max-average constraint satisfaction problems to non-Archimedean SDPs. A (*formal real*) *Puiseux series* is a series of the form

$$\mathbf{x} = \mathbf{x}(t) = c_0 t^{a/n} + c_1 t^{(a-1)/n} + c_2 t^{(a-2)/n} + \dots,$$

where t is a formal parameter, the number n is a positive integer, a is an integer, and the coefficients c_i are real. We further assume that $c_0 \neq 0$ for every Puiseux series except the zero series. The number a/n is called the *valuation* of the series \mathbf{x} and is denoted by $\text{val}(\mathbf{x})$, with the convention that $\text{val}(0) = -\infty$. Puiseux series can be added and multiplied in the natural way. Moreover, they can be ordered by defining $\mathbf{x} \geq 0$ if $c_0 \geq 0$ and $\mathbf{x} \geq \mathbf{y}$ if $\mathbf{x} - \mathbf{y} \geq 0$. It is known that Puiseux series form a real closed field, see, e.g., [BPR06, Section 2.6] for a detailed discussion.¹ We denote the field of Puiseux series by \mathbb{K} . For any $\mathbf{x}, \mathbf{y} \in \mathbb{K}$, the following properties of the valuation function are immediate from the definitions:

$$\begin{aligned} \text{val}(\mathbf{x} + \mathbf{y}) &\leq \max\{\text{val}(\mathbf{x}), \text{val}(\mathbf{y})\}, \\ \text{val}(\mathbf{x}\mathbf{y}) &= \text{val}(\mathbf{x}) + \text{val}(\mathbf{y}), \\ \mathbf{x} \geq \mathbf{y} \geq 0 &\implies \text{val}(\mathbf{x}) \geq \text{val}(\mathbf{y}). \end{aligned} \tag{2}$$

Theorem 4.1. *An instance ϕ with variables x_1, \dots, x_n of a max-average constraint satisfaction problem has a solution in $(\mathbb{Q} \cup \{-\infty\})^n$ if and only if the following system of inequalities $S_\phi(t)$, which is a (quadratic) semidefinite program over Puiseux series, has a solution in \mathbb{K}^n : for every variable x_i of ϕ we have a variable \mathbf{x}_i , and the inequalities are*

$$\begin{aligned} \mathbf{x}_{i_0} &\leq \mathbf{x}_{i_1} + \dots + \mathbf{x}_{i_k} && \text{for every constraint } x_{i_0} \leq \max(x_{i_1}, \dots, x_{i_k}) \text{ in } \phi, \\ \mathbf{x}_{i_0}^2 &\leq \mathbf{x}_{i_1} \mathbf{x}_{i_2} && \text{for every constraint } x_{i_0} \leq \frac{x_{i_1} + x_{i_2}}{2} \text{ in } \phi, \\ \mathbf{x}_{i_0} &= t^c && \text{for every constraint } x_{i_0} = c \text{ in } \phi, \\ \mathbf{x}_{i_0} &\geq 0 && \text{for every } i_0 \in \{1, \dots, n\}. \end{aligned} \tag{3}$$

Proof. **TOPROVE 1** □

5 From non-Archimedean to real SDPs

To conclude, we aim to reduce the non-Archimedean semidefinite program from (3) to a real semidefinite program. We will do this in two steps. First, we replace the formal parameter t by a real number that is sufficiently large (doubly exponential in the size of the CSP). In fact, any non-Archimedean semialgebraic feasibility problem can be reduced to a semialgebraic feasibility problem over the reals by replacing t with a sufficiently large real number. This follows from the quantifier elimination in real closed fields coupled with the definition of the order in Puiseux series. By using the effective bounds on quantifier elimination discussed, e.g., in [BPR06], we can give a bound on this “sufficiently large” number. Second, we will show that this SDP with very large coefficients can be reduced to another SDP that has coefficients of polynomial bit-size.

Before presenting the proofs, we note that a reduction from non-Archimedean SDPs coming from games to real SDPs with large coefficients was already studied in [AGS18, Theorem 36]. However, the bounds given in [AGS18] rely on an additional assumption on the underlying game. In particular, they cannot be applied directly to the SDP given in (3). For max-average systems coming from games, one could overcome this restriction by perturbing the underlying game slightly. Nevertheless, the bounds coming from quantifier elimination which we present below do not require any additional assumptions and can be of independent interest.

¹In this work, we use a convention in which one interprets t as a “very large” parameter. Many authors use the opposite but equivalent convention in which t is a very small parameter. This explains the differences in notation between our work and [BPR06].

Theorem 5.1. Let $P_1, \dots, P_m \in \mathbb{Z}[x_0, x_1, \dots, x_n]$ be a collection of polynomials with integer coefficients and let $\diamond \in \{\leq, \geq, <, >, =, \neq\}^m$ be a sign pattern. Consider the semialgebraic set over Puiseux series defined by

$$\mathbf{S} := \{\mathbf{x} \in \mathbb{K}^n : \forall i \in [m], P_i(t, \mathbf{x}) \diamond_i 0\}.$$

Furthermore, for any real $L > 0$ consider the semialgebraic set over the real numbers defined by

$$S(L) := \{x \in \mathbb{R}^n : \forall i \in [m], P_i(L, x) \diamond_i 0\}.$$

Then, \mathbf{S} is nonempty if and only if $S(L)$ is nonempty for all sufficiently large L . More precisely, there exists an absolute constant $C > 0$ with the following property. If the maximal degree of any polynomial P_i is bounded by $d \geq 2$ and s denotes the maximal bit-size of any coefficient of P_i , then for $L > 2^{\text{sd}^{Cn}}$ either all $S(L)$ are empty or all $S(L)$ are nonempty. In the former case \mathbf{S} is empty and in the latter case \mathbf{S} is nonempty.

Proof. **TOPROVE 2** □

In the next proposition and later, we use the following notation. Fix an instance ϕ of a max-average CSP and let $(c_i)_i$ be the rational constants used in the constraints of ϕ . For every i we denote $c_i = p_i/q_i$ where $p_i \in \mathbb{Z}$ and $q_i \in \mathbb{N}^*$ and we put $W := 2 + \max_i \{|p_i| + q_i\}$.

Proposition 5.2. There exists an absolute integer constant $M > 0$ such that an instance ϕ of a max-average constraint satisfaction problem has a solution if and only if the real SDP $S_\phi(L)$ obtained from $S_\phi(t)$ (see Theorem 4.1) by replacing t with $L > 2^{2^{Mn \log_2 W}}$, is feasible.

Proof. **TOPROVE 3** □

In Proposition 5.2, we obtain a (real) semidefinite program, whose size, however, is exponential in the size of ϕ and therefore does not provide a polynomial-time reduction to the semidefinite program feasibility problem. However, by the following result we can also reduce to a semidefinite program of polynomial representation size.

Theorem 5.3. For every $n \in \mathbb{Z}$, the singleton set $\{2^n\}$ is the projection of a spectrahedron whose representation size is polynomial in the bit-size of n .

Proof. **TOPROVE 4** □

Corollary 5.4. There is a polynomial-time reduction from the max-average constraint satisfaction problem to the feasibility problem for semidefinite programs.

Proof. **TOPROVE 5** □

Corollary 5.5. There is a polynomial-time reduction from stopping simple stochastic games to the feasibility problem for semidefinite programs.

Proof. **TOPROVE 6** □

Acknowledgments

This research benefited from the support of the FMJH Program PGMO. Manuel Bodirsky received funding from the ERC (Grant Agreement no. 101071674, POCOCOP). Views and opinions expressed are however those of the authors only and do not necessarily reflect those of the European Union or the European Research Council Executive Agency.

References

- [ABG07] Marianne Akian, Ravindra Bapat, and Stéphane Gaubert. Max-plus algebra. In *Chapter 25 in “Handbook of linear algebra”*, volume 39 of *Discrete Mathematics and Its Applications*. Chapman and Hall, 2007.
- [ABGJ14] X. Allamigeon, P. Benchimol, S. Gaubert, and M. Joswig. Combinatorial simplex algorithms can solve mean payoff games. *SIAM J. Optim.*, 24(4):2096–2117, 2014.
- [ABGJ15] X. Allamigeon, P. Benchimol, S. Gaubert, and M. Joswig. Tropicalizing the simplex algorithm. *SIAM J. Discrete Math.*, 29(2):751–795, 2015.
- [ABGJ21] X. Allamigeon, P. Benchimol, S. Gaubert, and M. Joswig. What tropical geometry tells us about the complexity of linear programming. *SIAM Rev.*, 63(1):123–164, 2021.
- [ABKPM09] Eric Allender, Peter Bürgisser, Johan Kjeldgaard-Pedersen, and Peter Bro Miltersen. On the complexity of numerical analysis. *SIAM J. Comput.*, 38(5):1987–2006, 2008/09.
- [AF17] David Avis and Oliver Friedmann. An exponential lower bound for Cunningham’s rule. *Math. Program.*, 161(1-2):271–305, 2017.
- [AGG12] M. Akian, S. Gaubert, and A. Guterman. Tropical polyhedra are equivalent to mean payoff games. *Int. J. Algebra Comput.*, 22(1):125001 (43 pages), 2012.
- [AGS18] X. Allamigeon, S. Gaubert, and M. Skomra. Solving generic nonarchimedean semidefinite programs using stochastic game algorithms. *J. Symbolic Comput.*, 85:25–54, 2018.
- [AM09] D. Andersson and P. B. Miltersen. The complexity of solving stochastic games on graphs. In *Proceedings of the 20th International Symposium on Algorithms and Computation (ISAAC)*, volume 5878 of *Lecture Notes in Comput. Sci.*, pages 112–121. Springer, 2009.
- [BBDFS23] N. Bertrand, P. Bouyer-Decitre, N. Fijalkow, and M. Skomra. Stochastic games. In N. Fijalkow, editor, *Games on Graphs*. 2023.
- [BEGM17] Endre Boros, Khaled Elbassioni, Vladimir Gurvich, and Kazuhisa Makino. A convex programming-based algorithm for mean payoff stochastic games with perfect information. *Optim. Lett.*, 11(8):1499–1512, 2017.
- [BKT24] Manuel Bodirsky, Mario Kummer, and Andreas Thom. Spectrahedral shadows and completely positive maps on real closed fields. *J. Eur. Math. Soc.*, 2024.
- [BM18] Manuel Bodirsky and Marcello Mamino. Tropically convex constraint satisfaction. *Theory of Computing Systems*, 62(3):481–509, 2018. An extended abstract of the paper appeared under the title “Max-Closed Semilinear Constraints” in the proceedings of CSR’16; preprint available under ArXiv:1506.04184.
- [BNRC08] Marc Bezem, Robert Nieuwenhuis, and Enric Rodríguez-Carbonell. The max-atom problem and its relevance. In *Proceedings of the International Conference on Logic for Programming, Artificial Intelligence, and Reasoning (LPAR)*, pages 47–61, 2008.

- [BPR06] S. Basu, R. Pollack, and M.-F. Roy. *Algorithms in Real Algebraic Geometry*, volume 10 of *Algorithms Comput. Math.* Springer, Berlin, 2006.
- [BTN01] A. Ben-Tal and A. Nemirovski. *Lectures on Modern Convex Optimization: Analysis, Algorithms, and Engineering Applications*, volume 2 of *MOS-SIAM Ser. Optim.* SIAM, Philadelphia, PA, 2001.
- [BW18] Julian C. Bradfield and Igor Walukiewicz. The mu-calculus and model checking. In Edmund M. Clarke, Thomas A. Henzinger, Helmut Veith, and Roderick Bloem, editors, *Handbook of Model Checking*, pages 871–919. Springer, 2018.
- [Can88] John Canny. Some algebraic and geometric computations in PSPACE. In *Proceedings of the Twentieth Annual ACM Symposium on Theory of Computing (STOC)*, pages 460–467, New York, NY, USA, 1988. ACM.
- [CFG022] Thomas Colcombet, Nathanaël Fijalkow, Pawel Gawrychowski, and Pierre Ohlmann. The theory of universal graphs for infinite duration games. *Log. Methods Comput. Sci.*, 18(3), 2022.
- [CJK⁺22] Cristian S. Calude, Sanjay Jain, Bakhadyr Khoussainov, Wei Li, and Frank Stephan. Deciding parity games in quasi-polynomial time. *SIAM J. Comput.*, 51(2):17–152, 2022.
- [Con92] A. Condon. The complexity of stochastic games. *Inform. and Comput.*, 96(2):203–224, 1992.
- [Con93] Anne Condon. On algorithms for simple stochastic games. In Jin-Yi Cai, editor, *Advances in Computational Complexity Theory*, volume 13 of *DIMACS Series in Discrete Mathematics and Theoretical Computer Science*, pages 51–72. AMS, 1993.
- [DFH23] Yann Disser, Oliver Friedmann, and Alexander V. Hopp. An exponential lower bound for Zadeh’s pivot rule. *Math. Program.*, 199(1):865–936, 2023.
- [EHS24] Friedrich Eisenbrand, Matthieu Haeberle, and Neta Singer. An Improved Bound on Sums of Square Roots via the Subspace Theorem. In Wolfgang Mulzer and Jeff M. Phillips, editors, *40th International Symposium on Computational Geometry (SoCG 2024)*, volume 293 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 54:1–54:8, Dagstuhl, Germany, 2024. Schloss Dagstuhl – Leibniz-Zentrum für Informatik.
- [EJ91] E. Allen Emerson and Charanjit S. Jutla. Tree automata, mu-calculus and determinacy. In *Proceedings of the 32nd annual Symposium on Foundations of Computer Science (FOCS’91)*, pages 368–377, Washington, DC, USA, 1991. IEEE Computer Society.
- [EJS93] E. A. Emerson, C. S. Jutla, and A. P. Sistla. On model-checking fragments of μ -calculus. In *Proceedings of the 5th International Conference on Computer-Aided Verification (CAV)*, volume 697 of *Lecture Notes in Comput. Sci.*, pages 385–396. Springer, 1993.
- [EM79] Andrzej Ehrenfeucht and Jan Mycielski. Positional strategies for mean payoff games. *International Journal of Game Theory*, 8(2):109–113, 1979.

- [FGMS20] John Fearnley, Spencer Gordon, Ruta Mehta, and Rahul Savani. Unique end of potential line. *Journal of Computer and System Sciences*, 114:1–35, 2020.
- [FHZ11] Oliver Friedmann, Thomas Dueholm Hansen, and Uri Zwick. Subexponential lower bounds for randomized pivoting rules for the simplex algorithm. In *Proceedings of the Forty-third Annual ACM Symposium on Theory of Computing (STOC)*, pages 283–292, New York, NY, USA, 2011. ACM.
- [FHZ14] Oliver Friedmann, Thomas Dueholm Hansen, and Uri Zwick. Errata for: A subexponential lower bound for the random facet algorithm for parity games. *CoRR*, abs/1410.7871, 2014.
- [FJS10] John Fearnley, Marcin Jurdziński, and Rahul Savani. Linear complementarity algorithms for infinite games. In *SOFSEM 2010: Theory and practice of computer science. 36th conference on current trends in theory and practice of computer science, Špindlerův Mlýn, Czech Republic, January 23–29, 2010. Proceedings*, pages 382–393. Berlin: Springer, 2010.
- [Fri11] O. Friedmann. A subexponential lower bound for Zadeh’s pivoting rule for solving linear programs and games. In *Proceedings of the 15th International Conference on Integer Programming and Combinatorial Optimization (IPCO)*, volume 6655 of *Lecture Notes in Comput. Sci.*, pages 192–206. Springer, 2011.
- [GGJ76] M. R. Garey, R. L. Graham, and D. S. Johnson. Some NP-complete geometric problems. In *Eighth Annual ACM Symposium on Theory of Computing (Hershey, Pa., 1976)*, pages 10–22. Assoc. Comput. Mach., New York, 1976.
- [GLS94] M. Grötschel, L. Lovász, and L. Schrijver. *Geometric Algorithms and Combinatorial Optimization*. Springer, Heidelberg, 1994. Second edition.
- [GR05] Bernd Gärtner and Leo Rüst. Simple stochastic games and P-matrix generalized linear complementarity problems. In *Fundamentals of computation theory. 15th international symposium, FCT 2005, Lübeck, Germany, August 17–20, 2005. Proceedings.*, pages 209–220. Berlin: Springer, 2005.
- [HIJ13] Thomas Dueholm Hansen and Rasmus Ibsen-Jensen. The complexity of interior point methods for solving discounted turn-based stochastic games. In *The nature of computation. Logic, algorithms, applications. 9th conference on computability in Europe, CiE 2013, Milan, Italy, July 1–5, 2013. Proceedings*, pages 252–262. Berlin: Springer, 2013.
- [Hod97] Wilfrid Hodges. *A shorter model theory*. Cambridge University Press, Cambridge, 1997.
- [JMT22] Marcin Jurdzinski, Rémi Morvan, and K. S. Thejaswini. Universal algorithms for parity games and nested fixpoints. In Jean-François Raskin, Krishnendu Chatterjee, Laurent Doyen, and Rupak Majumdar, editors, *Principles of Systems Design - Essays Dedicated to Thomas A. Henzinger on the Occasion of His 60th Birthday*, volume 13660 of *Lecture Notes in Computer Science*, pages 252–271. Springer, 2022.
- [JS08] Marcin Jurdziński and Rahul Savani. A simple P-matrix linear complementarity problem for discounted games. In *Logic and theory of algorithms. 4th conference on*

- computability in Europe, CiE 2008, Athens, Greece, June 15–20, 2008. Proceedings*, pages 283–293. Berlin: Springer, 2008.
- [Kha79] L. Khachiyan. A polynomial algorithm in linear programming. *Doklady Akademii Nauk SSSR*, 244:1093–1097, 1979.
 - [KRSW22] Jan Křetínský, Emanuel Ramneantu, Alexander Slivinskiy, and Maximilian Weininger. Comparison of algorithms for simple stochastic games. *Information and Computation*, 289:104885, 2022.
 - [Las09] J. B. Lasserre. *Moments, Positive Polynomials and Their Applications*. Ser. Optim. Appl. Imperial College Press, 2009.
 - [LB20] Karoliina Lehtinen and Udi Boker. Register games. *Log. Methods Comput. Sci.*, 16(2), 2020.
 - [LL69] T. M. Liggett and S. A. Lippman. Stochastic games with perfect information and time average payoff. *SIAM Rev.*, 11(4):604–607, 1969.
 - [LPSW22] Karoliina Lehtinen, Pawel Parys, Sven Schewe, and Dominik Wojtczak. A recursive approach to solving parity games in quasipolynomial time. *Log. Methods Comput. Sci.*, 18(1), 2022.
 - [Mar75] Donald A. Martin. Borel determinacy. *Annals of Mathematics*, 102(2):363–371, 1975.
 - [Mos91] A.W. Mostowski. Games with forbidden positions. Tech. Rep. 78, University of Gdansk, 1991.
 - [MSS04] Rolf H. Möhring, Martin Skutella, and Frederik Stork. Scheduling with and/or precedence constraints. *SIAM Journal on Computing*, 33(2):393–415, 2004.
 - [NN94] Y. Nesterov and A. Nemirovskii. *Interior-point polynomial algorithms in convex programming*, volume 13 of *Studies in Applied and Numerical Mathematics*. SIAM, Philadelphia, PA, 1994.
 - [PK97] Lorant Porkolab and Leonid Khachiyan. On the complexity of semidefinite programs. *Journal of Global Optimization*, 10(4):351–365, 1997.
 - [Pur95] Anuj Puri. *Theory of hybrid systems and discrete event systems*. University of California at Berkeley, 1995.
 - [Put05] M. L. Puterman. *Markov Decision Processes: Discrete Stochastic Dynamic Programming*. Wiley Ser. Probab. Stat. Wiley, Hoboken, NJ, 2005.
 - [Ram97] Motakuri V. Ramana. An exact duality theory for semidefinite programming and its complexity implications. *Mathematical Programming*, 77:129–162, 1997.
 - [Sch09] Sven Schewe. From parity and payoff games to linear programming. In *Proceedings of the 34th International Symposium on Mathematical Foundations of Computer Science (MFCS)*, pages 675–686, 2009.
 - [Sch18] C. Scheiderer. Spectrahedral shadows. *SIAM J. Appl. Algebra Geom.*, 2(1):26–44, 2018.

- [Sko18] M. Skomra. *Tropical spectrahedra: Application to semidefinite programming and mean payoff games*. PhD thesis, Université Paris-Saclay, 2018.
- [SV06] Ola Svensson and Sergei Vorobyov. Linear complementarity and P-matrices for stochastic games. In *Perspectives of Systems Informatics. 6th International Andrei Ershov Memorial Conference, PSI 2006*, volume 4378 of *Lecture Notes in Comput. Sci.*, pages 409–423. Springer, 2006.
- [TV08] Sergey P. Tarasov and Mikhail N. Vyalyi. Semidefinite programming and arithmetic circuit evaluation. *Discrete Applied Mathematics*, 156(11):2070 – 2078, 2008. In Memory of Leonid Khachiyan (1952-2005).
- [ZP96] Uri Zwick and Mike Paterson. The complexity of mean payoff games on graphs. *Theor. Comput. Sci.*, 158(1&2):343–359, 1996.