

Sudan National Micronutrient Survey Indicators Definition

Mark Myatt and Ernest Guevarra

18 May 2020

Contents

| | | |
|----------|---|----------|
| 1 | Background | 3 |
| 2 | Biomarkers variables | 3 |
| 2.1 | Haemoglobin | 3 |
| 2.2 | Serum ferritin and c-reactive protein | 6 |
| 2.3 | Calcium | 7 |
| 2.4 | Iodine | 8 |
| 3 | Micronutrient indicators | 8 |
| 3.1 | Anaemia prevalence indicators | 9 |
| 3.1.1 | AN1: Mild anaemia | 10 |
| 3.1.2 | AN2: Moderate anaemia | 13 |
| 3.1.3 | AN2C: Moderate anaemia in pregnant carers | 15 |
| 3.1.4 | AN3: Severe anaemia | 16 |
| 3.2 | Serum iron stores indicators | 19 |
| 3.2.1 | IR1: Iron deficiency indicators | 19 |
| 3.2.2 | IR2: Iron overload indicators | 21 |
| 3.3 | Acute inflammation indicators | 21 |
| 3.3.1 | AI1: Acute inflammation in children 6-59 months | 22 |
| 3.3.2 | AI2: Acute inflammation in non-pregnant carers | 22 |
| 3.3.3 | AI3: Acute inflammation in pregnant carers | 22 |

| | | |
|----------|---|-----------|
| 3.4 | Calcium stores indicators | 22 |
| 3.4.1 | CA1: Hypocalcemia indicators | 23 |
| 3.4.2 | CA2: Hypercalcemia indicators | 23 |
| 3.5 | Iodine concentration indicators | 24 |
| 3.5.1 | ID1: Iodine insufficiency indicator | 25 |
| 3.5.2 | ID2: Mild iodine deficiency in non-pregnant non-lactating | 25 |
| 3.5.3 | ID3: Moderate iodine deficiency in non-pregnant non-lactating | 25 |
| 3.5.4 | ID4: Severe iodine deficiency in non-pregnant non-lactating | 25 |
| 3.5.5 | ID5: Excessive iodine | 26 |
| 4 | Summary | 26 |
| 4.1 | Results presentation | 27 |

1 Background

To aid the analysis of the Sudan National Micronutrient Survey 2017-2018 data, appropriate indicators needed to be defined. The only documentation of indicators to be assessed from the survey was the last version of the S3M-II indicators list dated 16 November 2018. However, this document does not clearly define the indicators with no cut-off values provided. As such, indicator definitions were made based on a rapid literature review including micronutrient survey reports done elsewhere and reflected upon based actual available data from the survey itself to update the indicator definitions. This document presents these definitions.

2 Biomarkers variables

2.1 Haemoglobin

In the main S3M-II survey, we defined multiple indicators based on Hb data. These indicators represented the different severities of anaemia by different respondent groupings. Classification into these severity categories was based on Hb level cut-offs defined by WHO [[World Health Organization and Centers for Disease Control and Prevention, 2007](#), [World Health Organization, 2011b](#)] as follows:

Table 1: Hb levels to diagnose anaemia at sea level in grams per litre (g/L)

| Population | Mild | Moderate | Severe |
|---|-----------|----------|--------|
| Children 6-59 months of age | 100 - 109 | 70 - 99 | < 70 |
| Children 5-11 years of age | 110 - 114 | 80 - 109 | < 80 |
| Children 12-14 years of age | 110 - 119 | 80 - 109 | < 80 |
| Non-pregnant women (15 years and above) | 110 - 119 | 80 - 109 | < 80 |
| Pregnant women | 100 - 109 | 70 - 99 | < 70 |
| Men (15 years and above) | 110 - 129 | 80 - 109 | < 80 |

For the Sudan S3M-II main survey, no data was collected for children 5-17 years of age and for adult men 15 years of age and above so the indicator for this age group was not calculated and reported. When categorising respondents based on the above cut-offs in the main S3M-II survey, no adjustments to Hb were done based on altitude and for smoking history as recommended by WHO [[World Health Organization and Centers for Disease Control and Prevention, 2007](#), [World Health Organization, 2011b](#)].

We propose to analyse the Sudan National Micronutrient Survey data using the same indicator definitions used in the Sudan S3M-II main survey. We also propose to adjust Hb based on altitude of the PSU from where the data was collected. Altitude data will be gathered from publicly available elevation model data (such as the Shuttle Radar Topography Mission or

SRTM data that is available freely through various outlets for Sudan) if no altitude data can be provided by UNICEF. Map below shows elevation for Sudan based on publicly available SRTM data [[for Spatial Information](#)].




Figure 1: Shuttle Radar Topography Mission (SRTM) 90m Digital Elevation Model (DEM) for Sudan overlaid with the Sudan National Micronutrient survey primary sampling unit locations

With this data, we are able to extract elevation data for each of the PSUs in the Sudan Micronutrient Survey dataset.

Table 2: Sudan National Micronutrient Survey dataset with altitude extracted from SRTM 90m DEM

| psu | state | locality | sex | hb | altitude |
|-----|-------|----------|-----|--------------------|----------|
| 45 | 16 | 164 | 2 | 11.6 | 978 |
| 45 | 16 | 164 | 2 | 13.9 | 978 |
| 45 | 16 | 164 | 2 | 12.4 | 978 |
| 45 | 16 | 164 | 2 | 13.1 | 978 |
| 45 | 16 | 164 | 2 | 13.2 | 978 |
| 45 | 16 | 164 | 2 | 10.3 | 978 |
| 45 | 16 | 164 | 2 | 12.4 | 978 |
| 45 | 16 | 164 | 1 | 8.8000000000000007 | 978 |
| 45 | 16 | 164 | 2 | 11.8 | 978 |
| 45 | 16 | 164 | 1 | 8.1 | 978 |
| 45 | 16 | 164 | 2 | 12.4 | 978 |
| 45 | 16 | 164 | 2 | 9.1 | 978 |
| 45 | 16 | 164 | 2 | 13.3 | 978 |
| 45 | 16 | 164 | 1 | 9.8000000000000007 | 978 |
| 45 | 16 | 164 | 2 | 10.4 | 978 |
| 45 | 16 | 164 | 2 | 13 | 978 |
| 45 | 16 | 164 | 2 | 11.8 | 978 |
| 45 | 16 | 164 | 2 | 13 | 978 |
| 45 | 16 | 164 | 2 | 8.5 | 978 |
| 45 | 16 | 164 | 2 | 8.6 | 978 |
| 45 | 16 | 164 | 2 | 11.4 | 978 |
| 45 | 16 | 164 | 2 | 12.6 | 978 |
| 45 | 16 | 164 | 2 | 8.9 | 978 |
| 45 | 16 | 164 | 2 | 14.2 | 978 |
| 45 | 16 | 164 | 1 | 10.5 | 978 |
| 45 | 16 | 164 | 2 | 11.9 | 978 |
| 45 | 16 | 164 | 2 | 11.6 | 978 |
| 45 | 16 | 164 | 2 | 10.7 | 978 |
| 45 | 16 | 164 | 2 | 10.8 | 978 |
| 53 | 16 | 164 | 1 | 10.6 | 874 |
| 53 | 16 | 164 | 2 | 10 | 874 |
| 53 | 16 | 164 | NA | NA | 874 |
| 53 | 16 | 164 | 1 | 11.8 | 874 |
| 53 | 16 | 164 | 2 | NA | 874 |
| 53 | 16 | 164 | 2 | 12.4 | 874 |
| 53 | 16 | 164 | 1 | 7.9 | 874 |
| 53 | 16 | 164 | 1 | 8.4 | 874 |
| 53 | 16 | 164 | 2 | 11.7 | 874 |
| 53 | 16 | 164 | 2 | 10.6 | 874 |
| 53 | 16 | 164 | 2 | 12.5 | 874 |

Adjustments to measured Hb based on altitude will be done based on the following [\[World](#)

Health Organization, 2011b]:

Table 3: Altitude adjustments to measured haemoglobin concentrations

| Altitude (metres above sea level) | Measured haemoglobin adjustment (g/L) |
|---|--|
| < 1000 | 0 |
| 1000 | -2 |
| 1500 | -5 |
| 2000 | -8 |
| 2500 | -13 |
| 3000 | -19 |
| 3500 | -27 |
| 4000 | -35 |
| 4500 | -45 |

2.2 Serum ferritin and c-reactive protein

Normal serum ferritin levels range from 12 $\mu\text{g/L}$ to 150 $\mu\text{g/L}$. Following are the cut-offs for serum ferritin concentration that indicate either iron depletion or iron overload [World Health Organization and Centers for Disease Control and Prevention, 2007, Gorstein et al., 2007, Wegmüller et al., 2020, World Health Organization, 2011a].

Table 4: Relative extent of iron stores on the basis of serum ferritin concentration ($\mu\text{g/L}$)

| | Serum ferritin ($\mu\text{g/L}$) | | | |
|--|------------------------------------|--------|------------------|--------|
| | Less than 5 years | | 5 years or older | |
| | Male | Female | Male | Female |
| Depleted iron stores | < 12 | < 12 | < 15 | < 15 |
| Depleted iron stores in the presence of infection | < 30 | < 30 | - | - |
| Severe risk of iron overload (adults) | - | - | > 200 | > 150 |

Serum ferritin will be used to assess iron deficiency for children less than 5 and for any other individual above 5 years old. For children less than 5 years old, a cut-off for serum ferritin value of < 12 $\mu\text{g/L}$ indicates iron deficiency while for those older than 5 years old, a cut-off of < 15 $\mu\text{g/L}$ is used.

However, it has been recommended that serum ferritin values be adjusted based on inflammation status ideally using both of the acute phase proteins - C-reactive protein (CRP) and α_1 -acid glycoprotein (AGP) to yield the most unbiased estimates of iron deficiency. However, the Sudan Micronutrient Survey only assessed CRP in the samples. The recommended adjustments when

only one of the active phase proteins is available is to use an appropriate multiplier to the serum ferritin value depending on inflammation status of the respondent as described below:

Table 5: Cut-offs to determine inflammation

| Active Phase Protein | Cut-off |
|----------------------|----------|
| CRP | > 5 mg/L |
| AGP | > 1 g/L |

If a respondent is classified as being in an active inflammation process, then serum ferritin is adjusted accordingly. If inflammation is assessed using CRP only, the serum ferritin is adjusted by 0.65 [Thurnham et al., 2010].

2.3 Calcium

The range of normal values for serum calcium is age-dependent as shown below [Lietman et al., 2010]:

Table 6: Representative normal values for age for concentration of serum total calcium

| Target Group | Age | Serum total calcium (mg/dL) |
|--------------|------------|-----------------------------|
| Infants | 0-3 months | 8.8 - 11.3 |
| | 1-5 years | 9.4 - 10.8 |
| Children | 6-12 years | 9.4 - 10.3 |
| Men | 20 years | 9.1 - 10.2 |
| | 50 years | 8.9 - 10.0 |
| | 70 years | 8.8 - 9.9 |
| Women | 20 years | 8.8 - 10.0 |
| | 50 years | 8.8 - 10.0 |
| | 70 years | 8.8 - 10.0 |

We propose to use these normal ranges by age to determine whether a specific respondent group is hypocalcemic or below the normal range for their age or hypercalcemic or above the normal range for their age.

2.4 Iodine

Currently, cut-offs for urinary iodine are available for school-age children and older (6 years and older), pregnant women, and for lactating women and children aged less than 2 years.

Following are the various criteria for assessing iodine status in school-age children and older [World Health Organization, 2013]:

Table 7: Epidemiologic criteria for assessing iodine nutrition based on median urinary iodine concentration in school-age children and older

| Median urinary iodine (g/L) | Iodine intake | Iodine status |
|-----------------------------|--------------------|--|
| < 20 | Insufficient | Severe iodine deficiency |
| 20 - 49 | Insufficient | Moderate iodine deficiency |
| 50 - 99 | Insufficient | Mild iodine deficiency |
| 100 - 199 | Adequate | Adequate iodine nutrition |
| 200 - 299 | Above requirements | May pose a slight risk of more than adequate iodine intake in these populations |
| 300 | Excessive | Risk of adverse health consequences (iodine-induced hyperthyroidism, autoimmune thyroid disease) |

Following are the various criteria for assessing iodine status in pregnant women, lactating women and children aged less than 2 years [World Health Organization, 2013]:

Table 8: Epidemiologic criteria for assessing iodine nutrition based on median urinary iodine concentration in pregnant women, lactating women, and children aged less than 2 years

| Median urinary iodine (g/L) | Iodine intake |
|--|--------------------|
| Pregnant women | |
| < 150 | Insufficient |
| 150 - 249 | Adequate |
| 250 - 499 | Above requirements |
| 500 or more | Excessive |
| Lactating women and children aged less than 2 years | |
| < 100 | Insufficient |
| 100 or more | Adequate |

3 Micronutrient indicators

Given the biomarkers variables described above, we propose the following indicator sets.

3.1 Anaemia prevalence indicators

The anaemia indicators are:

Table 9: Anaemia indicators

| Indicator variable | Indicator Name |
|--------------------|--|
| AN1A | Mild anaemia in children 6-59 months |
| AN2B | Mild anaemia in non-pregnant carers |
| AN3C | Mild anaemia in pregnant carers |
| AN1A | Moderate anaemia in children 6-59 months |
| AN2B | Moderate anaemia in non-pregnant carers |
| AN3C | Moderate anaemia in pregnant carers |
| AN1A | Severe anaemia in children 6-59 months |
| AN2B | Severe anaemia in non-pregnant carers |
| AN3C | Severe anaemia in pregnant carers |

The anaemia indicators are calculated using data on **AGE**, **PREGNANCY** status and HB measurement (in g/L) of the respondent and on the **ALTITUDE** (in metres) of the location where the respondent resides.

3.1.1 AN1: Mild anaemia

3.1.1.1 AN1A: Mild Anaemia in children 6-59 months old

AN1A is TRUE IF

```
{
  AGE of respondent is between 6-59 months old AND
  {
    ALTITUDE < 1000 metres AND HB >= 100 g/L AND HB <= 109 g/L OR

    ALTITUDE >= 1000 metres AND ALTITUDE < 1500 metres AND
    HB is >= 102 g/L AND HB <= 111 g/L OR

    ALTITUDE >= 1500 metres AND ALTITUDE < 2000 metres AND
    HB >= 105 g/L AND HB <= 114 g/L OR

    ALTITUDE >= 2000 metres AND ALTITUDE < 2500 metres AND
    HB >= 108 g/L AND HB <= 117 g/L OR

    ALTITUDE >= 2500 metres AND ALTITUDE < 3000 metres AND
    HB >= 113 g/L AND HB <= 122 g/L OR

    ALTITUDE >= 3000 metres AND ALTITUDE is < 3500 metres AND
    HB >= 119 g/L AND HB <= 128 g/L OR

    ALTITUDE >= 3500 metres AND ALTITUDE < 4000 metres AND
    HB >= 127 g/L AND HB <= 136 g/L OR

    ALTITUDE >= 4000 metres AND ALTITUDE < 4500 metres AND
    HB >= 135 g/L AND HB <= 144 g/L OR

    ALTITUDE = 4500 metres and HB >= 145 g/L AND HB <= 154 g/L
  }
}
```

3.1.1.2 ANC1B: Mild anaemia in non-pregnant carers

```
ANC1B is TRUE IF
{
  AGE of respondent between 15 and 49 years AND NOT PREGNANT AND
  {
    ALTITUDE < 1000 metres AND HB >= 110 g/L AND HB <= 119 g/L OR

    ALTITUDE >= 1000 metres AND ALTITUDE < 1500 metres AND
    HB >= 112 g/L AND HB <= 121 g/L OR

    ALTITUDE >= 1500 metres AND ALTITUDE < 2000 metres AND
    HB >= 115 g/L AND HB <= 124 g/L OR

    ALTITUDE >= 2000 metres AND ALTITUDE < 2500 metres AND
    HB >= 118 g/L AND HB <= 127 g/L OR

    ALTITUDE >= 2500 metres AND ALTITUDE < 3000 metres AND
    HB >= 123 g/L AND HB <= 132 g/L OR

    ALTITUDE >= 3000 metres AND ALTITUDE < 3500 metres AND
    HB >= 129 g/L AND HB <= 138 g/L OR

    ALTITUDE >= 3500 metres AND ALTITUDE < 4000 metres AND
    HB >= 137 g/L AND HB <= 146 g/L OR

    ALTITUDE >= 4000 metres AND ALTITUDE < 4500 metres AND
    HB >= 145 g/L AND HB <= 154 g/L OR

    ALTITUDE = 4500 metres AND HB >= 155 g/L AND HB <= 164 g/L
  }
}
```

3.1.1.3 AN1C: Mild anaemia in pregnant carers

```
AN1C is TRUE IF
{
  AGE of respondent between 15 and 49 years old AND PREGNANT AND
  {
    ALTITUDE < 1000 metres AND HB >= 100 g/L AND HB <= 109 g/L OR

    ALTITUDE >= 1000 metres AND ALTITUDE < 1500 metres AND
    HB >= 102 g/L AND HB <= 111 g/L OR

    ALTITUDE >= 1500 metres AND ALTITUDE < 2000 metres AND
    HB >= 105 g/L AND HB <= 114 g/L OR

    ALTITUDE >= 2000 metres AND ALTITUDE < 2500 metres AND
    HB >= 108 g/L AND HB <= 117 g/L OR

    ALTITUDE >= 2500 metres AND ALTITUDE < 3000 metres AND
    HB >= 113 g/L AND HB <= 122 g/L OR

    ALTITUDE >= 3000 metres AND ALTITUDE < 3500 metres AND
    HB >= 119 g/L AND HB <= 128 g/L OR

    ALTITUDE >= 3500 metres AND ALTITUDE < 4000 metres AND
    HB >= 127 g/L AND HB <= 136 g/L OR

    ALTITUDE >= 4000 metres AND ALTITUDE < 4500 metres AND
    HB >= 135 g/L AND HB <= 144 g/L OR

    ALTITUDE = 4500 metres AND HB >= 145 g/L AND HB <= 154 g/L
  }
}
```

3.1.2 AN2: Moderate anaemia

3.1.2.1 AN2A: Moderate anaemia in children 6-59 months

AN2A is TRUE IF

```
{
  AGE of respondent is between 6-59 months old AND
  {
    ALTITUDE < 1000 metres AND HB >= 70 g/L AND HB <= 99 g/L OR

    ALTITUDE >= 1000 metres AND ALTITUDE < 1500 metres AND
    HB >= 72 g/L AND HB <= 101 g/L OR

    ALTITUDE >= 1500 metres AND ALTITUDE < 2000 metres AND
    HB >= 75 g/L AND HB <= 104 g/L OR

    ALTITUDE >= 2000 metres AND ALTITUDE < 2500 metres AND
    HB >= 78 g/L AND HB <= 107 g/L OR

    ALTITUDE >= 2500 metres AND ALTITUDE < 3000 metres AND
    HB >= 83 g/L AND HB <= 112 g/L OR

    ALTITUDE >= 3000 metres AND ALTITUDE < 3500 metres AND
    HB >= 89 g/L AND HB <= 118 g/L OR

    ALTITUDE >= 3500 metres AND ALTITUDE < 4000 metres AND
    HB >= 97 g/L AND HB <= 126 g/L OR

    ALTITUDE >= 4000 metres AND ALTITUDE < 4500 metres AND
    HB >= 105 g/L AND HB <= 134 g/L OR

    ALTITUDE is = 4500 metres AND HB >= 115 g/L AND <= 144 g/L
  }
}
```

3.1.2.2 AN2B: Moderate anaemia in non-pregnant carers

```
AN2B is TRUE IF
{
  AGE of respondent is between 15 and 49 years AND NOT PREGNANT AND
  {
    ALTITUDE < 1000 metres AND HB >= 80 g/L AND HB <= 109 g/L OR

    ALTITUDE >= 1000 metres AND ALTITUDE < 1500 metres AND
    HB >= 82 g/L AND HB <= 111 g/L OR

    ALTITUDE >= 1500 metres AND ALTITUDE < 2000 metres AND
    HB >= 85 g/L AND HB <= 114 g/L OR

    ALTITUDE >= 2000 metres AND ALTITUDE < 2500 metres AND
    HB >= 88 g/L AND HB <= 117 g/L OR

    ALTITUDE >= 2500 metres AND ALTITUDE < 3000 metres AND
    HB >= 93 g/L AND HB <= 122 g/L OR

    ALTITUDE >= 3000 metres AND ALTITUDE < 3500 metres AND
    HB >= 99 g/L AND HB <= 128 g/L OR

    ALTITUDE >= 3500 metres AND ALTITUDE < 4000 metres AND
    HB >= 107 g/L AND HB <= 136 g/L OR

    ALTITUDE >= 4000 metres AND ALTITUDE < 4500 metres AND
    HB >= 115 g/L AND HB <= 144 g/L OR

    ALTITUDE is = 4500 metres AND HB >= 125 g/L and HB <= 154 g/L
  }
}
```

3.1.3 AN2C: Moderate anaemia in pregnant carers

```
AN2C is TRUE IF
{
  AGE of respondent is between 15 and 49 years old AND PREGNANT AND
  {
    ALTITUDE < 1000 metres AND HB >= 70 g/L AND HB <= 99 g/L OR

    ALTITUDE >= 1000 metres AND ALTITUDE < 1500 metres AND
    HB >= 72 g/L AND HB <= 101 g/L OR

    ALTITUDE >= 1500 metres AND ALTITUDE < 2000 metres AND
    HB >= 75 g/L AND HB <= 104 g/L OR

    ALTITUDE >= 2000 metres AND ALTITUDE < 2500 metres AND
    HB >= 78 g/L AND HB <= 107 g/L OR

    ALTITUDE >= 2500 metres AND ALTITUDE < 3000 metres AND
    HB >= 83 g/L AND HB <= 112 g/L OR

    ALTITUDE >= 3000 metres AND ALTITUDE < 3500 metres AND
    HB >= 89 g/L AND HB <= 118 g/L OR

    ALTITUDE >= 3500 metres AND ALTITUDE < 4000 metres AND
    HB >= 97 g/L AND HB <= 126 g/L OR

    ALTITUDE >= 4000 metres AND ALTITUDE < 4500 metres AND
    HB >= 105 g/L AND HB <= 134 g/L OR

    ALTITUDE is = 4500 metres AND HB >= 115 g/L AND <= 144 g/L
  }
}
```

3.1.4 AN3: Severe anaemia

3.1.4.1 AN3A: Severe anaemia in children 6-59 months

AN3 is TRUE IF

```
{  
  AGE of respondent is between 6-59 months old AND  
  {  
    ALTITUDE < 1000 metres AND HB < 70 g/L OR  
  
    ALTITUDE >= 1000 metres AND ALTITUDE < 1500 metres AND HB < 72 g/L OR  
  
    ALTITUDE >= 1500 metres AND ALTITUDE < 2000 metres AND HB < 75 g/L OR  
  
    ALTITUDE >= 2000 metres AND ALTITUDE < 2500 metres AND HB < 78 g/L OR  
  
    ALTITUDE >= 2500 metres AND ALTITUDE < 3000 metres AND HB < 83 g/L OR  
  
    ALTITUDE >= 3000 metres AND ALTITUDE < 3500 metres AND HB < 89 g/L OR  
  
    ALTITUDE >= 3500 metres AND ALTITUDE < 4000 metres AND HB < 97 g/L OR  
  
    ALTITUDE >= 4000 metres AND ALTITUDE < 4500 metres AND HB < 105 g/L OR  
  
    ALTITUDE is = 4500 metres AND HB < 115 g/L  
  }  
}
```


3.1.4.2 ANC3B: Severe anameia in non-pregnant carers

ANC3B is TRUE IF

```
{
  AGE of respondent is between 15 and 49 years AND NOT PREGNANT AND
  {
    ALTITUDE < 1000 metres AND HB < 80 g/L OR

    ALTITUDE >= 1000 metres AND ALTITUDE < 1500 metres AND HB < 82 g/L OR

    ALTITUDE >= 1500 metres AND ALTITUDE < 2000 metres AND HB < 85 g/L OR

    ALTITUDE >= 2000 metres AND ALTITUDE < 2500 metres AND HB < 88 g/L OR

    ALTITUDE >= 2500 metres AND ALTITUDE < 3000 metres AND HB < 93 g/L OR

    ALTITUDE >= 3000 metres AND ALTITUDE < 3500 metres AND HB < 99 g/L OR

    ALTITUDE >= 3500 metres AND ALTITUDE < 4000 metres AND HB < 107 g/L OR

    ALTITUDE >= 4000 metres AND ALTITUDE < 4500 metres AND HB < 115 g/L OR

    ALTITUDE is = 4500 metres AND HB < 125 g/L
  }
}
```

3.1.4.3 ANC3C: Severe anaemia in pregnant carers

AN3 is TRUE IF

```
{
  AGE of respondent is between 15 and 49 years old AND PREGNANT AND
  {
    ALTITUDE < 1000 metres AND HB < 70 g/L OR

    ALTITUDE >= 1000 metres AND ALTITUDE < 1500 metres AND HB < 72 g/L OR

    ALTITUDE >= 1500 metres AND ALTITUDE < 2000 metres AND HB < 75 g/L OR

    ALTITUDE >= 2000 metres AND ALTITUDE < 2500 metres AND HB < 78 g/L OR

    ALTITUDE >= 2500 metres AND ALTITUDE < 3000 metres AND HB < 83 g/L OR

    ALTITUDE >= 3000 metres AND ALTITUDE < 3500 metres AND HB < 89 g/L OR

    ALTITUDE >= 3500 metres AND ALTITUDE < 4000 metres AND HB < 97 g/L OR

    ALTITUDE >= 4000 metres AND ALTITUDE < 4500 metres AND HB < 105 g/L OR

    ALTITUDE is = 4500 metres AND HB < 115 g/L
  }
}
```

3.2 Serum iron stores indicators

The serum iron stores indicators are:

Table 10: Body iron stores indicators

| Indicator variable | Indicator name |
|--------------------|---|
| IR1A | Iron deficiency in children 6-59 months |
| IR1B | Iron deficiency in non-pregnant carers |
| IR1C | Iron deficiency in pregnant carers |
| IR2A | Iron overload in non-pregnant carers |
| IR2B | Iron overload in pregnant carers |

Serum iron stores indicators require data on **AGE**, **PREGNANCY** status, **CRP** and serum **FERRITIN** concentration.

3.2.1 IR1: Iron deficiency indicators

3.2.1.1 IR1A: Iron deficiency in children 6-59 months

```
IR1A is TRUE IF
{
  AGE of respondent is between 6-59 months old AND
  {
    CRP <= 5 mg/L AND serum FERRITIN < 12 µg/L OR

    CRP > 5 mg/L AND serum FERRITIN < 19.8 µg/L
  }
}
```

3.2.1.2 IR1B: Iron deficiency in non-pregnant carers

```
IR1B is TRUE IF
{
  AGE of respondent is between 15 and 49 years old AND NOT PREGNANT AND
  {
    CRP <= 5 mg/L AND serum FERRITIN < 15 µg/L OR

    CRP > 5 mg/L AND serum FERRITIN < 24.75 µg/L
  }
}
```

3.2.1.3 IR1C: Iron deficiency in pregnant carers

IR1C is TRUE IF

```
{  
  AGE of respondent is between 15 and 49 years old AND  
  {  
    CRP <= 5 mg/L AND serum FERRITIN < 15 µg/L OR  
  
    CRP > 5 mg/L AND serum FERRITIN < 24.75 µg/L  
  }  
}
```

3.2.2 IR2: Iron overload indicators

3.2.2.1 IR2A: Iron overload in non-pregnant carers

IR2A is **TRUE** if either...

```
{
  AGE of respondent is between 15 and 49 years old AND NOT PREGNANT AND
  {
    CRP <= 5 mg/L AND serum FERRITIN > 150 µg/L OR

    CRP > 5 mg/L AND serum FERRITIN > 247.5 µg/L
  }
}
```

3.2.2.2 IR2B: Iron overload in pregnant carers

IR2B is **TRUE** if either...

```
{
  AGE of respondent is between 15 and 49 years old AND PREGNANT AND
  {
    CRP <= 5 mg/L AND serum FERRITIN > 150 µg/L OR

    CRP > 5 mg/L AND serum FERRITIN > 247.5 µg/L
  }
}
```

3.3 Acute inflammation indicators

The acute inflammation indicators are:

Table 11: Acute inflammation indicators

| Indicator variable | Indicator name |
|--------------------|--|
| AI1 | Acute inflammation in children 6-59 months |
| AI2 | Acute inflammation in non-pregnant carers |
| AI3 | Acute inflammation in pregnant carers |

Acute inflammation (AI) indicators require serum **CRP** concentration.

3.3.1 AI1: Acute inflammation in children 6-59 months

AI1 is **TRUE** IF AGE is between 6 and 59 months old AND CRP > 5 mg/L

3.3.2 AI2: Acute inflammation in non-pregnant carers

AI2 is **TRUE** IF AGE is between 15 and 49 years old AND
NOT PREGNANT AND CRP > 5 mg/L

3.3.3 AI3: Acute inflammation in pregnant carers

AI3 is **TRUE** IF AGE is between 15 and 49 years old AND
PREGNANT AND CRP > 5 mg/L

3.4 Calcium stores indicators

The calcium stores indicators are:

Table 12: Serum calcium stores indicators

| Indicator variable | Indicator name |
|--------------------|--|
| CA1A | Hypocalcemia in children 12-59 months old |
| CA1B | Hypocalcemia in non-pregnant carers |
| CA1C | Hypocalcemia in pregnant carers |
| CA2A | Hypercalcemia in children 12-59 months old |
| CA2B | Hypercalcemia in non-pregnant carers |
| CA2C | Hypercalcemia in pregnant carers |

The serum calcium stores indicators require **AGE**, **SEX**, **PREGNANCY** status and serum **CALCIUM** concentration.

3.4.1 CA1: Hypocalcemia indicators

3.4.1.1 CA1A: Hypocalcemia in children 12-59 months

```
CA1A is TRUE IF
{
  AGE of respondent is between 12 and 59 months AND
  serum CALCIUM < 9.4 mg/dL
}
```

3.4.1.2 CA1B: Hypocalcemia in non-pregnant carers

```
CA1B is TRUE IF
{
  AGE of respondent > 12 years and <= 70 years AND
  NOT PREGNANT AND serum CALCIUM < 8.8 mg/dL
}
```

3.4.1.3 CA1C: Hypocalcemia in pregnant carers

```
CA1C is TRUE IF
{
  AGE of respondent > 12 years and <= 70 years AND
  PREGNANT and serum CALCIUM < 8.8 mg/dL
}
```

3.4.2 CA2: Hypercalcemia indicators

3.4.2.1 CA2A: Hypercalcemia in children 12-59 months

```
CA2A is TRUE IF
{
  AGE of respondent is between 12 and 59 months AND
  serum CALCIUM > 10.8 mg/dL
}
```

3.4.2.2 CA2B: Hypercalcemia in non-pregnant carers

```
CA2B is TRUE IF
{
  AGE of respondent > 12 years and <= 70 years AND
  NOT PREGNANT and serum CALCIUM > 10.0 mg/dL
}
```

3.4.2.3 CA2C: Hypercalcemia in pregnant carers

```
CA2C is TRUE IF
{
  AGE of respondent > 12 years and <= 70 years AND
  PREGNANT and serum CALCIUM > 10.0 mg/dL
}
```

3.5 Iodine concentration indicators

The iodine concentration indicators are:

Table 13: Iodine concentration indicators

| Indicator variable | Indicator name |
|--------------------|---|
| ID1A | Insufficient iodine in children less than 24 months old |
| ID1B | Insufficient iodine in lactating carers |
| ID1C | Insufficient iodine in pregnant carers |
| ID2 | Mild iodine deficiency in non-pregnant non-lactating carers |
| ID3 | Moderate iodine deficiency in non-pregnant non-lactating carers |
| ID4 | Severe iodine deficiency in non-pregnant non-lactating carers |
| ID5A | Excessive iodine in pregnant carers |
| ID5B | Excessive iodine in non-pregnant non-lactating carers |

The iodine concentration indicators require **AGE**, **SEX**, **PREGNANCY** status and **LACTATION** status and urinary **IODINE** concentration.

3.5.1 ID1: Iodine insufficiency indicator

3.5.1.1 ID1A: Insufficient iodine in children 6-59 months

ID1A is **TRUE** IF AGE is less than 24 months old AND urinary IODINE < 100 g/L

3.5.1.2 ID1B: Insufficient iodine in lactating carers

ID1B is **TRUE** IF AGE is between 15 and 49 years old AND
LACTATING AND urinary ****IODINE**** < 100 g/L

3.5.1.3 ID1C: Insufficient iodine in pregnant carers

ID1C is **TRUE** IF AGE is between 15 and 49 years old AND
PREGNANT AND urinary IODINE < 150 g/L

3.5.2 ID2: Mild iodine deficiency in non-pregnant non-lactating

ID2 is **TRUE** IF AGE is between 15-49 years old AND
NOT PREGNANT AND NOT LACTATING AND
urinary IODINE >= 50 g/L AND urinary IODINE <= 99 g/L

3.5.3 ID3: Moderate iodine deficiency in non-pregnant non-lactating

ID3 is **TRUE** IF AGE is between 15-49 years old AND
NOT PREGNANT AND NOT LACTATING AND
urinary IODINE >= 20 g/L AND urinary IODINE <= 49 g/L

3.5.4 ID4: Severe iodine deficiency in non-pregnant non-lactating

ID4 is **TRUE** IF AGE is between 15-49 years old AND
NOT PREGNANT AND NOT LACTATING AND
urinary IODINE < 20 g/L.

3.5.5 ID5: Excessive iodine

3.5.5.1 ID5A: Excessive iodine in pregnant carers

ID5A is **TRUE** if AGE is between **15-49** years old AND
NOT PREGNANT AND NOT LACTATING AND
urinary IODINE **>= 300** g/L

3.5.5.2 ID5B: Excessive iodine in non-pregnant non-lactating carers

ID5B is **TRUE** IF AGE is between **15-49** years old AND
PREGNANT AND urinary IODINE **>= 500** g/L

4 Summary

Given these indicator definition, we propose the following list of indicators to be analysed and reported for the Sudan Micronutrient Survey.

Table 14: List of proposed indicators for Sudan National Micronutrient Survey

| Category | Indicator.variable | Indicator.name | Type |
|--------------------|--------------------|---|------------|
| Anaemia | AN1A | Mild anaemia in children 6-59 months | Proportion |
| | AN2B | Mild anaemia in non-pregnant carers | Proportion |
| | AN3C | Mild anaemia in pregnant carers | Proportion |
| | AN1A | Moderate anaemia in children 6-59 months | Proportion |
| | AN2B | Moderate anaemia in non-pregnant carers | Proportion |
| | AN3C | Moderate anaemia in pregnant carers | Proportion |
| | AN1A | Severe anaemia in children 6-59 months | Proportion |
| | AN2B | Severe anaemia in non-pregnant carers | Proportion |
| | AN3C | Severe anaemia in pregnant carers | Proportion |
| Iron deficiency | IR1A | Iron deficiency in children 6-59 months | Proportion |
| | IR1B | Iron deficiency in non-pregnant carers | Proportion |
| | IR1C | Iron deficiency in pregnant carers | Proportion |
| Iron overload | IR2A | Iron overload in non-pregnant carers | Proportion |
| | IR2B | Iron overload in pregnant carers | Proportion |
| Inflammation | AI1 | Acute inflammation in children 6-59 months | Proportion |
| | AI2 | Acute inflammation in non-pregnant carers | Proportion |
| | AI3 | Acute inflammation in pregnant carers | Proportion |
| Calcium deficiency | CA1A | Hypocalcemia in children 12-59 months old | Proportion |
| | CA1B | Hypocalcemia in non-pregnant carers | Proportion |
| | CA1C | Hypocalcemia in pregnant carers | Proportion |
| Calcium overload | CA2A | Hypercalcemia in children 12-59 months old | Proportion |
| | CA2B | Hypercalcemia in non-pregnant carers | Proportion |
| | CA2C | Hypercalcemia in pregnant carers | Proportion |
| Iodine deficiency | ID1A | Insufficient iodine in children less than 24 months old | Proportion |
| | ID1B | Insufficient iodine in lactating carers | Proportion |
| | ID1C | Insufficient iodine in pregnant carers | Proportion |
| | ID2 | Mild iodine deficiency in non-pregnant non-lactating carers | Proportion |
| | ID3 | Moderate iodine deficiency in non-pregnant non-lactating carers | Proportion |
| | ID4 | Severe iodine deficiency in non-pregnant non-lactating carers | Proportion |
| | ID5A | Excessive iodine in pregnant carers | Proportion |
| Iodine excess | ID5B | Excessive iodine in non-pregnant non-lactating carers | Proportion |

4.1 Results presentation

Estimates with corresponding confidence intervals for these indicators will be presented in tables. Estimates for these indicators will also be presented as choropleth maps at the state level for which estimates will be representative of.

In addition, we will summarise the biomarker values using the usual 6 figure summary (minimum value, first quantile, median, mean, third quartile and maximum value) and then present the distribution of the biomarker values as a violin plot with the range of normal values shown. An example violin plot for Hb is shown below.




Figure 2: Example violin plot for Hb in children 6-59 months sample by state

For absurdly high or low outliers such as that in State 17, we propose to winsorize with the aim of discarding as little data as possible by bringing down high values or bringing up low values enough that no bias is introduced into the estimate of mean values.

References

- CGIAR-Consortium for Spatial Information. SRTM 90m DEM Digital Elevation Database. URL <http://srtm.csi.cgiar.org>.
- J Gorstein, K M Sullivan, I Parvanta, and F Begin. Indicators and Methods for Cross-Sectional Surveys of Vitamin and Mineral Status of Populations. Technical report, Ottawa, Canada and Atlanta, Georgia, USA, May 2007.
- Steven A Lietman, Emily L Germain-Lee, and Michael A Levine. Hypercalcemia in children and adolescents. *Current Opinion in Pediatrics*, 22(4):508–515, August 2010.
- David I Thurnham, Linda D McCabe, Sumanto Haldar, Frank T Wieringa, Christine A Northrop-Clewes, and George P McCabe. Adjusting plasma ferritin concentrations to remove the effects of subclinical inflammation in the assessment of iron deficiency: a meta-analysis. *The American journal of clinical nutrition*, 92(3):546–555, July 2010.
- Rita Wegmüller, Helena Bentil, James P Wirth, Nicolai Petry, Sherry A Tanumihardjo, Lindsay Allen, Thomas N Williams, Lilian Selenje, Abraham Mahama, Esi Amoahful, Matilda Steiner-Asiedu, Seth Adu-Afarwuah, and Fabian Rohner. Anemia, micronutrient deficiencies, malaria, hemoglobinopathies and malnutrition in young children and non-pregnant women in Ghana: Findings from a national survey. *PloS one*, 15(1):e0228258–19, January 2020.
- World Health Organization. Serum ferritin concentrations for the assessment of iron status and iron deficiency in populations. Technical report, Geneva, 2011a.
- World Health Organization. Haemoglobin concentrations for the diagnosis of anaemia and assessment of severity. Technical report, World Health Organization, Geneva, 2011b.
- World Health Organization. Urinary iodine concentrations for determining iodine status deficiency in populations. *Vitamin and Mineral Nutrition Information System*, pages 1–5, 2013.
- World Health Organization and Centers for Disease Control and Prevention. Assessing iron status of populations: including literature reviews. In *Joint World Health Organization and Centers for Disease Control and Prevention Technical Consultation on the Assessment of Iron Status at the Population Level Geneva, Switzerland*, pages 1–112. World Health Organization, 2007.