Scattering Amplitudes: From QFT to Modern Methods

Ernest Yeung

April 5, 2026

Contents

I Quantum Field Theory (QFT Foundations) 7

1 Quantum Field Theory (QFT) Foundations 9
1.1 LSZ Reduction Formula . . . 9
1.2 Path Integrals . . . 9
1.3 Feynman Rules for Gauge Theories . . . 9
1.4 Non-Abelian Gauge Theory . . . 9

2 Helicity States and Weyl Spinors 11
2.1 Lorentz Group Representations . . . 11
2.2 Left-Handed Weyl Spinors ψ_α . . . 11
2.3 Right-Handed Weyl Spinors $\tilde{\psi}_{\dot{\alpha}}$ . . . 12
2.4 The ε Symbol: $\mathrm{SL}(2,\mathbb{C})$ Metric . . . 12
2.4.1 Raising and lowering . . . 12
2.5 Lorentz-Invariant Spinor Products . . . 13
2.6 The σ^μ Symbol: Vector–Spinor Dictionary . . . 13
2.6.1 Key identities . . . 13
2.6.2 Momentum bi-spinor . . . 13
2.7 Helicity from the Little Group . . . 14
2.8 The Weyl Lagrangian . . . 14
2.8.1 Equations of motion . . . 14
2.8.2 Gamma matrices: Weyl representation . . . 14
2.9 Majorana and Dirac Fermions . . . 15
2.10 Summary: Spinor Toolkit . . . 15

II Spinor-Helicity Formalism 17

3 Spinor-Helicity Formalism 19
3.1 The Factorization of Null Momenta . . . 19
3.2 Spinor Brackets . . . 19
3.3 Affine (Inhomogeneous) Spinor Coordinates . . . 20
3.4 Schouten Identity . . . 20
3.5 Polarization Vectors . . . 20
3.6 Momentum Twistors . . . 20

4 Maximally Helicity Violating (MHV) Amplitudes 21
4.1 Why "Maximally Helicity Violating"? . . . 21
4.2 The Parke-Taylor Formula . . . 21
4.3 Helicity Configuration Table . . . 22
4.4 Single-Minus Amplitudes Are Nonzero . . . 22
4.5 BCJ Relations . . . 23

4.6 Double Copy . . . 23

5 On-Shell Recursion (BCFW) 25
5.1 The BCFW Shift . . . 25
5.2 The Recursion Formula . . . 25
5.3 Why It Works . . . 25
5.4 Application to Single-Minus . . . 25
5.5 Soft Limits . . . 26

III Computational Tools 27

6 Computational Tools 29
6.1 Recommended Tool Stack . . . 29
6.2 lips . . . 29
6.3 SymPy . . . 29
6.4 CADABRA2 . . . 29
6.5 FORM . . . 30

IV Research Frontiers 31

7 Research Frontiers 33
7.1 The Amplituhedron . . . 33
7.2 Celestial Amplitudes . . . 33
7.3 Single-Minus Open Problems . . . 33
7.4 Twistor String Theory . . . 33

V Literature Review 35

8 Literature Review and Paper Notes 37
8.1 A Brief Introduction to Modern Amplitude Methods . . . 37
8.1.1 Key Concepts Covered . . . 37
8.1.2 Important Results & Equations . . . 37
8.1.3 Why This Paper Matters for Our Project . . . 38
8.2 Single-Minus Gluon Tree Amplitudes Are Nonzero . . . 39
8.2.1 The Discovery . . . 39
8.2.2 Key Equations . . . 39
8.2.3 Proof Strategy . . . 39
8.2.4 Open Directions (from the paper) . . . 40
8.2.5 Why This Paper Matters for Our Project . . . 40

VI Bridges: Modern Literature to Srednicki 41

9 Dixon–Srednicki Bridge: From Textbook QFT to Modern Amplitudes 43
9.1 Color Decomposition (Dixon §2 ↔ Srednicki Ch.79–80) . . . 43
9.1.1 SU(N_c) Generators and Dixon's Non-Standard Normalization . . . 43
9.1.2 Deriving Dixon (2.2): Structure Constants from the Commutator . . . 44
9.1.3 The SU(N_c) Fierz Identity (Dixon (2.3)) . . . 45
9.1.4 Proving Tracelessness via the Fierz Identity . . . 45
9.1.5 Srednicki 80.17 and Color-Ordered Amplitudes . . . 47

9.2 Spinor Variables (Dixon §3.1 ↔ Srednicki Ch.48, 50) . . . 47
9.2.1 Trading k^μ for Spinors: The Core Idea . . . 47
9.2.2 Notation Comparison Table . . . 48
9.2.3 Positive Energy Projector (Dixon (3.5)) . . . 48
9.2.4 Reconstructing 4-Momentum from Spinors (Dixon (3.6) ↔ Srednicki 50.15) 48
9.2.5 Explicit Spinor Representations . . . 49
9.3 Spinor Product Identities (Dixon §3 ↔ Srednicki Ch.50) . . . 49
9.3.1 Anti-Symmetry (Dixon (3.13) ↔ Srednicki) . . . 49
9.3.2 Squaring (Dixon (3.14) ↔ Srednicki (60.4)) . . . 50
9.3.3 Momentum Conservation (Dixon (3.15) ↔ Srednicki (60.18)) . . . 50
9.3.4 Schouten Identity (Dixon (3.16) ↔ Srednicki (50.21)/(60.17)) . . . 51
9.3.5 The Complex Square Root Property and Collinear Singularities . . . 52
9.4 Four-Point Example (Dixon §3.2 ↔ Srednicki Ch.60) . . . 52
9.4.1 The Setup: $e^+e^- \to q\bar{q}$ via a Photon . . . 52
9.4.2 Deriving Dixon (3.24): The Four-Point Amplitude . . . 52
9.4.3 Deriving Dixon (3.25): The Parity-Conjugate Amplitude . . . 54
9.5 Helicity Formalism for Massless Vectors (Dixon §3.3 ↔ Srednicki Ch.81) . . . 55
9.5.1 Gluon Polarization Vectors as Spinor Bilinears . . . 55
9.5.2 Slashed Polarization Vectors (Srednicki (60.15)) . . . 55
9.5.3 Key Simplification: Reference Momentum Choice . . . 55
9.5.4 Srednicki Ch. 81 and Color-Ordered Feynman Rules . . . 55
9.6 Soft Gluon Limit (Dixon §4.1) . . . 56
9.6.1 Definition of the Soft Limit . . . 56
9.6.2 Soft Factor Derivation (Dixon (4.1)) . . . 56
9.6.3 Soft Limit Singularity Structure . . . 57
9.6.4 Connection to Weinberg Soft Theorem and SMGA Paper . . . 57
9.7 Summary: Dixon–Srednicki Dictionary . . . 58

Part I

Quantum Field Theory (QFT Foundations)

Chapter 1

Quantum Field Theory (QFT) Foundations

1.1 LSZ Reduction Formula

1.2 Path Integrals

1.3 Feynman Rules for Gauge Theories

1.4 Non-Abelian Gauge Theory

Chapter 2

Helicity States and Weyl Spinors

This chapter builds the spinor algebra that underlies everything in Parts II and III. The physical target is simple: massless particles carry definite *helicity* $h = \pm\frac{1}{2}$ (or ± 1 for gluons), and the two-component Weyl spinors λ_α and $\tilde{\lambda}_{\dot{\alpha}}$ are precisely the objects that represent those helicity eigenstates. We establish this connection carefully here so that Chapter 3 can use the spinor-bracket formalism without re-deriving its foundations.

2.1 Lorentz Group Representations

The Lorentz algebra $\mathfrak{so}(1,3)$ is (complexified) isomorphic to $\mathfrak{su}(2)_L \oplus \mathfrak{su}(2)_R$ via the non-Hermitian combinations

$$N_i \equiv \tfrac{1}{2}(J_i - iK_i), \qquad N_i^\dagger \equiv \tfrac{1}{2}(J_i + iK_i), \tag{2.1}$$

where J_i are rotation generators and K_i are boost generators. They satisfy $[N_i, N_j] = i\varepsilon_{ijk}N_k$, $[N_i^\dagger, N_j^\dagger] = i\varepsilon_{ijk}N_k^\dagger$, $[N_i, N_j^\dagger] = 0$, so irreducible representations are labelled by two half-integers $(n,\, n')$.

(n, n')	Srednicki	Dim	Field	Index type
$(0,0)$	$(1,1)$	1	scalar ϕ	none
$(\frac{1}{2},0)$	$(2,1)$	2	left-handed Weyl ψ_α	undotted
$(0,\frac{1}{2})$	$(1,2)$	2	right-handed Weyl $\tilde{\psi}_{\dot{\alpha}}$	dotted
$(\frac{1}{2},\frac{1}{2})$	$(2,2)$	4	vector A^μ	spacetime

The key point for helicity: under a rotation by angle θ about the momentum axis (the little group of a massless particle), a $(\frac{1}{2},0)$ field picks up phase $e^{-i\theta/2}$ (helicity $-\frac{1}{2}$) and a $(0,\frac{1}{2})$ field picks up $e^{+i\theta/2}$ (helicity $+\frac{1}{2}$).

2.2 Left-Handed Weyl Spinors ψ_α

A left-handed Weyl field $\psi_\alpha(x)$ lives in the $(\frac{1}{2},0)$ representation. Under an infinitesimal Lorentz transformation $\delta\omega_{\mu\nu}$:

$$\delta\psi_\alpha = \tfrac{i}{2}\,\delta\omega_{\mu\nu}\,(S_{\mathrm{L}}^{\mu\nu})_\alpha{}^\beta\,\psi_\beta. \tag{2.2}$$

The six generators are explicitly (Srednicki eqs. 34.9–34.10):

$$(S_{\mathrm{L}}^{ij})_\alpha{}^\beta = \tfrac{1}{2}\varepsilon^{ijk}\sigma_k, \qquad (S_{\mathrm{L}}^{k0})_\alpha{}^\beta = \tfrac{i}{2}\sigma_k, \tag{2.3}$$

where σ_k are the Pauli matrices:

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \tag{2.4}$$

These satisfy the Lorentz algebra $[S_L^{\mu\nu}, S_L^{\rho\sigma}] = i(g^{\nu\rho} S_L^{\mu\sigma} - \cdots)$. The boost generators S_L^{k0} are anti-Hermitian (the Lorentz group is non-compact), reflecting that boosts are not unitary.

2.3 Right-Handed Weyl Spinors $\tilde{\psi}_{\dot{\alpha}}$

Hermitian conjugation swaps $N_i \leftrightarrow N_i^\dagger$, sending $(\frac{1}{2}, 0) \to (0, \frac{1}{2})$. So

$$(\psi_\alpha)^\dagger \equiv \tilde{\psi}_{\dot{\alpha}}(x) \tag{2.5}$$

is a *right-handed* Weyl field with a *dotted* index. Its generators are (Srednicki eq. 34.17):

$$\boxed{(S_{\mathrm{R}}^{\mu\nu})_{\dot{\alpha}}{}^{\dot{\beta}} = -[(S_{\mathrm{L}}^{\mu\nu})_\alpha{}^\beta]^*} \tag{2.6}$$

Concretely this reverses the sign of boosts (consistent with $L \leftrightarrow R$ under parity) and from eq. (35.22):

$$S_{\mathrm{R}}^{\mu\nu} = -\tfrac{i}{4}(\bar{\sigma}^\mu \sigma^\nu - \bar{\sigma}^\nu \sigma^\mu). \tag{2.7}$$

Convention note. We adopt Srednicki's metric $g^{\mu\nu} = \mathrm{diag}(+1, -1, -1, -1)$ and $\varepsilon^{12} = \varepsilon_{21} = +1$ throughout. The amplitudes literature often uses the same conventions; we flag any deviation when it appears.

2.4 The ε Symbol: SL$(2, \mathbb{C})$ Metric

From $(\frac{1}{2}, 0) \otimes (\frac{1}{2}, 0) = (0, 0)_A \oplus (1, 0)_S$ there is a unique (up to scale) invariant antisymmetric symbol. With Srednicki's normalization (eq. 34.22 / 35.1):

$$\varepsilon^{12} = \varepsilon_{21} = +1, \qquad \varepsilon^{21} = \varepsilon_{12} = -1, \tag{2.8}$$

so in matrix form:

$$\varepsilon_{\alpha\beta} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \qquad \varepsilon^{\alpha\beta} = \begin{pmatrix} 0 & +1 \\ -1 & 0 \end{pmatrix}, \qquad \varepsilon_{\alpha\beta}\varepsilon^{\beta\gamma} = \delta_\alpha{}^\gamma. \tag{2.9}$$

The same structure holds for dotted indices.

2.4.1 Raising and lowering

$$\psi^\alpha = \varepsilon^{\alpha\beta}\psi_\beta, \qquad \psi_\alpha = \varepsilon_{\alpha\beta}\psi^\beta. \tag{2.10}$$

Sign trap (Srednicki eq. 34.27):

$$\psi^a \chi_a = \varepsilon^{ab}\psi_b\chi_a = -\varepsilon^{ba}\psi_b\chi_a = -\psi_b\chi^b. \tag{2.11}$$

Hence $\psi^\alpha\chi_\alpha = -\psi_\alpha\chi^\alpha$: *index position matters and carries a sign.*

2.5 Lorentz-Invariant Spinor Products

The fundamental Lorentz-invariant bilinears are:

$$\psi\chi \equiv \psi^\alpha \chi_\alpha = \varepsilon^{\alpha\beta}\psi_\alpha\chi_\beta, \tag{2.12}$$

$$\tilde{\psi}\tilde{\chi} \equiv \tilde{\psi}_{\dot\alpha}\tilde{\chi}^{\dot\alpha} = \varepsilon_{\dot\alpha\dot\beta}\tilde{\psi}^{\dot\alpha}\tilde{\chi}^{\dot\beta}. \tag{2.13}$$

For Grassmann-valued fields, $\psi\chi = +\chi\psi$ (two minus signs cancel; Srednicki eq. 35.25), and $(\psi\chi)^\dagger = \tilde{\chi}\tilde{\psi}$.

These are the precursors of the spinor brackets $\langle ij\rangle$ and $[ij]$ introduced in Chapter 3: when $\psi_\alpha = \lambda_{i\alpha}$ and $\chi_\alpha = \lambda_{j\alpha}$ are the spinors of massless momenta p_i, p_j, then $\langle ij\rangle = \varepsilon^{\alpha\beta}\lambda_{i\alpha}\lambda_{j\beta}$ and $[ij] = \varepsilon_{\dot\alpha\dot\beta}\tilde{\lambda}_i^{\dot\alpha}\tilde{\lambda}_j^{\dot\beta}$.

2.6 The σ^μ Symbol: Vector–Spinor Dictionary

A field in the $(\frac{1}{2}, \frac{1}{2})$ representation carries one undotted and one dotted index. The dictionary to 4-vectors is (Srednicki eq. 34.28–34.30):

$$A_{\alpha\dot\alpha} = \sigma^\mu_{\alpha\dot\alpha} A_\mu, \qquad \sigma^\mu_{\alpha\dot\alpha} = (I, \vec{\sigma}). \tag{2.14}$$

Explicitly:

$$\sigma^0 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \ \sigma^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \ \sigma^2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \ \sigma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \tag{2.15}$$

Raising both spinor indices via the ε symbols gives (Srednicki eq. 35.19–35.20):

$$\bar{\sigma}^{\mu\,\dot\alpha\alpha} \equiv \varepsilon^{\alpha\beta}\varepsilon^{\dot\alpha\dot\beta}\,\sigma^\mu_{\beta\dot\beta} = (I, -\vec{\sigma}). \tag{2.16}$$

2.6.1 Key identities

$$\sigma^\mu_{\alpha\dot\alpha}\,\sigma_{\mu\beta\dot\beta} = -2\,\varepsilon_{\alpha\beta}\,\varepsilon_{\dot\alpha\dot\beta}, \tag{35.4}$$

$$\varepsilon^{\alpha\beta}\varepsilon^{\dot\alpha\dot\beta}\,\sigma^\mu_{\alpha\dot\alpha}\,\sigma^\nu_{\beta\dot\beta} = -2\,g^{\mu\nu}, \tag{35.5}$$

$$\mathrm{tr}(\sigma^\mu\bar{\sigma}^\nu) \equiv \sigma^\mu_{\alpha\dot\alpha}\,\bar{\sigma}^{\nu\,\dot\alpha\alpha} = 2\,g^{\mu\nu}, \tag{2.17}$$

$$\sigma^\mu\bar{\sigma}^\nu + \sigma^\nu\bar{\sigma}^\mu = -2\,g^{\mu\nu}\,I_2. \tag{35-Clifford}$$

The last line is the Weyl-spinor analogue of the Dirac Clifford algebra. The generators are its antisymmetric part:

$$S_{\mathrm{L}}^{\mu\nu} = \tfrac{i}{4}(\sigma^\mu\bar{\sigma}^\nu - \sigma^\nu\bar{\sigma}^\mu). \tag{35.21}$$

2.6.2 Momentum bi-spinor

For any 4-momentum p^μ, define

$$p_{\alpha\dot\alpha} \equiv \sigma^\mu_{\alpha\dot\alpha}\,p_\mu = \begin{pmatrix} p^0 + p^3 & p^1 + ip^2 \\ p^1 - ip^2 & p^0 - p^3 \end{pmatrix}. \tag{2.18}$$

Then $\det(p_{\alpha\dot\alpha}) = (p^0)^2 - |\mathbf{p}|^2 = p^2$, so $p^2 = 0$ iff the matrix has rank 1 — the factorization $p_{\alpha\dot\alpha} = \lambda_\alpha\tilde{\lambda}_{\dot\alpha}$ is equivalent to masslessness. This is the starting point of Chapter 3.

2.7 Helicity from the Little Group

For a massless particle with momentum p^μ, the little group (the subgroup of the Lorentz group that fixes p^μ) is $ISO(2)$, whose compact part is $U(1)$. Under the $U(1)$ rotation by angle θ, a field of helicity h transforms as $e^{-2ih\theta}$.

In terms of spinors: the little-group generator rescales

$$\lambda_\alpha \to e^{i\theta/2}\lambda_\alpha, \qquad \tilde{\lambda}_{\dot\alpha} \to e^{-i\theta/2}\tilde{\lambda}_{\dot\alpha}, \tag{2.19}$$

i.e. the continuous parameter $t = e^{i\theta/2}$ acts as

$$\lambda \to t\lambda, \qquad \tilde{\lambda} \to t^{-1}\tilde{\lambda}. \tag{2.20}$$

A state created by λ_α has helicity $-\frac{1}{2}$ (left-handed, positive-chirality), and one created by $\tilde{\lambda}_{\dot\alpha}$ has helicity $+\frac{1}{2}$ (right-handed, negative-chirality).

For a spin-1 gluon the helicity-± 1 polarization vectors scale as $\varepsilon^\pm_\mu \to t^{\mp 2}\varepsilon^\pm_\mu$ under (2.20); this determines the form of $\varepsilon^\pm$ in terms of spinors (see §3.5).

2.8 The Weyl Lagrangian

For a single left-handed Weyl field ψ_α, the Lorentz-invariant hermitian Lagrangian is (Srednicki eq. 36.2):

$$\boxed{\mathcal{L} = i\tilde{\psi}_{\dot\alpha}\,\bar{\sigma}^{\mu\,\dot\alpha\alpha}\,\partial_\mu\psi_\alpha - \tfrac{1}{2}m\,\varepsilon^{\alpha\beta}\psi_\alpha\psi_\beta - \tfrac{1}{2}m\,\varepsilon_{\dot\alpha\dot\beta}\tilde{\psi}^{\dot\alpha}\tilde{\psi}^{\dot\beta}.} \tag{2.21}$$

The kinetic term is real modulo a total divergence. The mass term is a *Majorana mass*: it breaks the $U(1)$ symmetry $\psi \to e^{i\alpha}\psi$ and is non-zero because ψ_α are Grassmann-valued.

2.8.1 Equations of motion

Varying with respect to $\tilde{\psi}^{\dot\alpha}$:

$$-i\,\bar{\sigma}^{\mu\,\dot\alpha\alpha}\,\partial_\mu\psi_\alpha + m\,\tilde{\psi}^{\dot\alpha} = 0. \tag{36.4}$$

For $m = 0$ this is the **Weyl equation** for a massless left-handed spinor. It is the field equation whose plane-wave solutions give the spinors $\lambda_\alpha(p)$ used in amplitude calculations.

2.8.2 Gamma matrices: Weyl representation

Assembling ψ_α and $\tilde{\psi}_{\dot\alpha}$ into a 4-component Majorana spinor $\Psi = (\psi_\alpha, \tilde{\psi}^{\dot\alpha})^T$, the equations of motion take the Dirac form $(-i\gamma^\mu\partial_\mu + m)\Psi = 0$ with (Srednicki eq. 36.7):

$$\gamma^\mu = \begin{pmatrix} 0 & \sigma^\mu_{\alpha\dot\beta} \\ \bar{\sigma}^{\mu\,\dot\alpha\beta} & 0 \end{pmatrix}, \qquad \{\gamma^\mu, \gamma^\nu\} = -2g^{\mu\nu}I_4. \tag{36.7, 36.9}$$

The chiral projectors are $P_L = \frac{1}{2}(I - \gamma^5) = \mathrm{diag}(I_2, 0)$ and $P_R = \frac{1}{2}(I + \gamma^5) = \mathrm{diag}(0, I_2)$ with $\gamma^5 = i\gamma^0\gamma^1\gamma^2\gamma^3 = \mathrm{diag}(-I_2, +I_2)$.

2.9 Majorana and Dirac Fermions

Property	Majorana	Dirac
Weyl fields	1 (ψ)	2 (χ, ξ)
$U(1)$ symmetry	No	Yes
Conserved charge	None	$Q = \int j^0 \, d^3x$
Self-conjugate ($\Psi^C = \Psi$)	Yes	No
Lagrangian	$\frac{i}{2}\bar{\Psi}\gamma^\mu\partial_\mu\Psi - \frac{m}{2}\bar{\Psi}\Psi$	$i\bar{\Psi}\gamma^\mu\partial_\mu\Psi - m\bar{\Psi}\Psi$
Real d.o.f.	2	4

In amplitude calculations with external gluons (massless spin-1), the internal spinors are always effectively massless Weyl spinors; the Majorana/Dirac distinction matters for external fermion lines.

2.10 Summary: Spinor Toolkit

Object	Expression / Rule
Left-handed field	ψ_α (undotted index, helicity $-\frac{1}{2}$)
Right-handed field	$\tilde{\psi}_{\dot{\alpha}} = (\psi_\alpha)^\dagger$ (dotted, $+\frac{1}{2}$)
Rotation generator	$(S_L^{ij})_\alpha{}^\beta = \frac{1}{2}\varepsilon^{ijk}\sigma_k$
Boost generator	$(S_L^{k0})_\alpha{}^\beta = \frac{i}{2}\sigma_k$
R-generators	$S_R^{\mu\nu} = -[S_L^{\mu\nu}]^*$
ε normalization	$\varepsilon^{12} = \varepsilon_{21} = +1$
Raise/lower	$\psi^\alpha = \varepsilon^{\alpha\beta}\psi_\beta, \quad \psi_\alpha = \varepsilon_{\alpha\beta}\psi^\beta$
Sign identity	$\psi^\alpha\chi_\alpha = -\psi_\alpha\chi^\alpha$
Angle bracket	$\psi\chi = \varepsilon^{\alpha\beta}\psi_\alpha\chi_\beta = +\chi\psi$
Square bracket	$\tilde{\psi}\tilde{\chi} = \varepsilon_{\dot{\alpha}\dot{\beta}}\tilde{\psi}^{\dot{\alpha}}\tilde{\chi}^{\dot{\beta}}$
σ^μ	$(I, \vec{\sigma}), \quad \bar{\sigma}^\mu = (I, -\vec{\sigma})$
Trace	$\mathrm{tr}(\sigma^\mu\bar{\sigma}^\nu) = 2g^{\mu\nu}$
Clifford	$\sigma^\mu\bar{\sigma}^\nu + \sigma^\nu\bar{\sigma}^\mu = -2g^{\mu\nu}I_2$
Momentum bi-spinor	$p_{\alpha\dot{\alpha}} = \sigma^\mu_{\alpha\dot{\alpha}}p_\mu, \quad \det = p^2$
Little group	$\lambda \to t\lambda$, $\tilde{\lambda} \to t^{-1}\tilde{\lambda}$
Weyl Lagrangian	$i\tilde{\psi}\bar{\sigma}^\mu\partial_\mu\psi - \frac{m}{2}\psi\psi - \frac{m}{2}\tilde{\psi}\tilde{\psi}$
Weyl equation ($m = 0$)	$i\bar{\sigma}^{\mu\,\dot{\alpha}\alpha}\partial_\mu\psi_\alpha = 0$

Next: Chapter 3 uses the momentum bi-spinor factorization $p_{\alpha\dot{\alpha}} = \lambda_\alpha\tilde{\lambda}_{\dot{\alpha}}$ (which follows directly from $p^2 = 0$ and the rank-1 condition above) to build the spinor-bracket formalism and polarization vectors for massless gauge bosons.

Part II

Spinor-Helicity Formalism

Chapter 3

Spinor-Helicity Formalism

3.1 The Factorization of Null Momenta

For massless particles, the momentum p^μ satisfies $p^2 = 0$. In the Van der Waerden notation, the momentum bi-spinor is defined via the Pauli matrices $\sigma^\mu = (\mathbf{1}, \vec{\sigma}_{\text{Pauli}})$:

$$p_{\alpha\dot{\alpha}} = p^\mu (\sigma_\mu)_{\alpha\dot{\alpha}}. \tag{3.1}$$

Explicitly in 2×2 matrix form:

$$p_{\alpha\dot{\alpha}} = \begin{pmatrix} p^0 + p^3 & p^1 + ip^2 \\ p^1 - ip^2 & p^0 - p^3 \end{pmatrix}. \tag{3.2}$$

For massless p ($p^2 = 0$), $\det(p_{\alpha\dot{\alpha}}) = (p^0)^2 - |\mathbf{p}|^2 = 0$, so this matrix has rank 1 and factorizes as:

$$p_{\alpha\dot{\alpha}} = \lambda_\alpha \tilde{\lambda}_{\dot{\alpha}} \tag{3.3}$$

where λ_α and $\tilde{\lambda}_{\dot{\alpha}}$ are commuting (bosonic) Weyl spinors (Srednicki Sec. 20.3, 42.1). This factorization is unique up to the little group scaling:

$$\lambda_\alpha \to t\lambda_\alpha, \quad \tilde{\lambda}_{\dot{\alpha}} \to t^{-1}\tilde{\lambda}_{\dot{\alpha}}, \quad t \in \mathbb{C}^*. \tag{3.4}$$

For real momenta $t \in U(1)$; for complex momenta λ and $\tilde{\lambda}$ are independent. A left-handed (positive helicity $+\frac{1}{2}$) Weyl spinor scales as $\lambda \to t\lambda$; a right-handed one as $\tilde{\lambda} \to t^{-1}\tilde{\lambda}$.

3.2 Spinor Brackets

The fundamental Lorentz-invariant contractions are (Srednicki eq. 42.16):

$$\langle ij \rangle = \langle ij \rangle = \epsilon^{\alpha\beta} \lambda_{i\alpha} \lambda_{j\beta} = \lambda_i^0 \lambda_j^1 - \lambda_i^1 \lambda_j^0 \tag{3.5}$$

$$[ij] = [ij] = \epsilon^{\dot{\alpha}\dot{\beta}} \tilde{\lambda}_{i\dot{\alpha}} \tilde{\lambda}_{j\dot{\beta}} \tag{3.6}$$

with $\epsilon^{01} = -\epsilon^{10} = 1$ (Levi-Civita). These are antisymmetric: $\langle ij \rangle = -\langle ji \rangle$, $[ij] = -[ji]$.

The dot product of two massless momenta is encoded by:

$$2\, p_i \cdot p_j = \text{tr}(\not{p}_i \not{p}_j) = \langle ij \rangle [ji]. \tag{3.7}$$

In particular $p_i^2 = 0$ implies $\langle ii \rangle = [ii] = 0$.

3.3 Affine (Inhomogeneous) Spinor Coordinates

For explicit calculations (as in arXiv:2602.12176v1, Eq. (2)), one fixes the little-group gauge and writes the spinors in affine coordinates on $\mathbb{CP}^1$:

$$|i\rangle \equiv \lambda_i = \begin{pmatrix} 1 \\ z_i \end{pmatrix}, \qquad |i] \equiv \tilde{\lambda}_i = \omega_i \begin{pmatrix} 1 \\ \tilde{z}_i \end{pmatrix}, \tag{3.8}$$

where $z_i, \tilde{z}_i \in \mathbb{C}$ are inhomogeneous (affine) coordinates on the Riemann sphere and $\omega_i \in \mathbb{C}^*$ carries the little-group weight (so $\omega_i \to t^{-1}\omega_i$ under $\lambda \to t\lambda$). In this gauge the spinor brackets read:

$$\langle ij\rangle = z_j - z_i, \tag{3.9}$$

$$[ij] = \omega_i\omega_j\,(\tilde{z}_j - \tilde{z}_i). \tag{3.10}$$

The dot product identity (3.7) then becomes $2p_i \cdot p_j = (z_j - z_i)\,\omega_i\omega_j(\tilde{z}_i - \tilde{z}_j)$. This coordinate patch is particularly convenient for the half-collinear kinematics relevant to single-minus amplitudes (§4.4).

3.4 Schouten Identity

For any three spinors λ_i, λ_j, λ_k:

$$\langle ij\rangle\lambda_{k\alpha} + \langle jk\rangle\lambda_{i\alpha} + \langle ki\rangle\lambda_{j\alpha} = 0 \tag{3.11}$$

Equivalently:

$$\langle ij\rangle\langle kl\rangle = \langle ik\rangle\langle jl\rangle + \langle il\rangle\langle kj\rangle \tag{3.12}$$

3.5 Polarization Vectors

For a gluon of momentum p and helicity $\pm$:

$$\varepsilon^+_\mu = \frac{\langle r|\gamma_\mu|p]}{\sqrt{2}\langle rp\rangle} \tag{3.13}$$

$$\varepsilon^-_\mu = \frac{\langle p|\gamma_\mu|r]}{\sqrt{2}[pr]} \tag{3.14}$$

where r is an arbitrary reference spinor.

3.6 Momentum Twistors

For planar amplitudes, momentum twistors $\mathcal{Z}_i^A = (\lambda_i^\alpha, \mu_i^{\dot\alpha})$ provide a dual description where:

$$x_i^{\alpha\dot\alpha} - x_{i+1}^{\alpha\dot\alpha} = p_i^{\alpha\dot\alpha} = \lambda_i^\alpha\tilde{\lambda}_i^{\dot\alpha} \tag{3.15}$$

The dual conformal symmetry becomes manifest in these variables.

Chapter 4

Maximally Helicity Violating (MHV) Amplitudes

4.1 Why "Maximally Helicity Violating"?

For a tree-level n-gluon amplitude $A_n(h_1, \ldots, h_n)$ with $h_i = \pm 1$:

- Let m_- denote the number of negative-helicity gluons.
- Naive Yang–Mills vertices with all same helicity ($+\cdots+$ or $-\cdots-$) are strongly suppressed — amplitudes with $m_- = 0$ or $m_- = n$ vanish at tree level (and at all loops).
- Amplitudes with $m_- = 1$ (single-minus) were long conjectured to vanish; see §4.4.
- In twistor space (Witten 2004), tree amplitudes are supported on curves of degree $d = m_- - 1 + \ell$, so at tree level $\ell = 0$ the non-zero sector requires $0 \le m_- \le 4$ generically.

MHV = maximally helicity violating corresponds to $m_- = 2$, i.e. $m_+ = n - 2$. This is the maximum number of "wrong-helicity" gluons that gives a generically non-zero amplitude, in the sense that every cyclic ordering is non-zero. The name reflects that having only 2 negative-helicity gluons out of n is the extreme violation of a naive helicity-conservation intuition.

4.2 The Parke-Taylor Formula

For n-gluon scattering with legs j and k carrying negative helicity, the tree-level color-ordered amplitude is:

$$\boxed{A_n^{\mathrm{MHV}}(\ldots, j^-, \ldots, k^-, \ldots) = i\frac{\langle jk\rangle^4}{\langle 12\rangle\langle 23\rangle\cdots\langle n1\rangle}} \tag{4.1}$$

(momentum-conserving delta function $\delta^4(P)$ implicit). For the canonical choice $(j, k) = (1, 2)$ this reads $i\,\langle 12\rangle^4/(\langle 12\rangle\langle 23\rangle\cdots\langle n1\rangle)$.

This compact formula replaces $O(n!)$ Feynman diagrams with a single expression.

Little group weight check. Under $\lambda_i \to t_i\lambda_i$, $\tilde{\lambda}_i \to t_i^{-1}\tilde{\lambda}_i$, a helicity-$h_i$ gluon requires the amplitude to scale as $t_i^{-2h_i}$.

- Negative-helicity legs j, k: numerator $\langle jk\rangle^4 \sim t_j^4 t_k^4$, denom contributes $t_j^{-1}t_k^{-1}$ (one factor each), net $t_j^4/t_j^{-1} = t_j^{4+1}$... more carefully: $\langle jk\rangle^4$ in numerator gives weight $+4$ for j and $+4$ for k; each $\langle\cdot j\cdot\rangle$ in denominator gives -1; there are exactly 2 such factors, so net weight for j is $4 - 2 = +2 = -2h_j$ with $h_j = -1$. ✓
- Positive-helicity legs: appear twice in the denominator (once as $\langle i{,}i{+}1\rangle$ and once as $\langle i{-}1{,}i\rangle$), giving weight $-2 = -2(+1)$. ✓

4.3 Helicity Configuration Table

Helicity Configuration	Tree Amplitude	Status
$(-,-,+,+,\ldots,+)$	$\neq 0$	MHV
$(-,+,+,+,\ldots,+)$	$\neq 0$	MHV (Parke-Taylor)
$(+,-,+,+,\ldots,+)$	$\neq 0$	MHV (cyclic)
$(-,-,-,+,\ldots,+)$	$\neq 0$	NMHV
$(-,+,+,+,+,+)$	$\neq 0$	Single-minus (2026 result)
$(+,+,+,+,+,+)$	$= 0$	Vanishing

4.4 Single-Minus Amplitudes Are Nonzero

Recent work (Guevara, Lupsasca, Skinner, Strominger, Weil, arXiv:2602.12176) proves that single-minus ($m_- = 1$) tree amplitudes are non-zero for all $n \geq 4$.

$$A_n(1^-, 2^+, \ldots, n^+) \neq 0 \quad \forall\, n \geq 4. \tag{4.2}$$

This breaks the longstanding conjecture that only $m_- \geq 2$ configurations were generically nonvanishing.

Half-collinear locus (arXiv:2602.12176 Eq. (13)). The single-minus amplitude is supported on the *half-collinear locus*:

$$\langle ij \rangle = 0 \quad \forall\, i, j \in \{1, \ldots, n\}. \tag{4.3}$$

This means all λ_i are proportional, i.e. $\lambda_i = c_i\, \lambda_1$ for some $c_i \in \mathbb{C}$, so all n momenta are collinear. The $\delta(\langle 1a \rangle)$ factors in the ansatz below enforce this support explicitly for $a = 2, \ldots, n$.

Explicit single-minus ansatz (arXiv:2602.12176 Eq. (15)). The ansatz is:

$$\begin{aligned} \mathcal{A}_n(1^-, 2^+, \ldots, n^+) = i^{2-n}\, \frac{\langle r1 \rangle^{n+1}}{\langle r2 \rangle \langle r3 \rangle \cdots \langle rn \rangle}\, A_{1\cdots n} \prod_{a=2}^{n} \delta(\langle 1a \rangle) \\ \times\ \delta^2\Big(\sum_{i=1}^{n} \langle ri \rangle\, \tilde{\lambda}_i\Big) \end{aligned} \tag{4.4}$$

where r is the reference spinor, $A_{1\cdots n}$ is a scalar prefactor, and $\delta^2(\cdots)$ is a 2-component ($\dot{\alpha}$) delta function imposing a residual momentum constraint. In affine coordinates (§3.3), $\langle 1a \rangle = z_a - z_1$, so the delta functions set $z_a = z_1$ for all a.

MHV comparison (arXiv:2602.12176 Eq. (11)). For comparison, the MHV amplitude in the same notation with negative-helicity legs r, s reads:

$$\mathcal{A}_n^{\mathrm{MHV}} = i\, \langle rs \rangle^4\, \mathrm{PT}_{\mathrm{cyc}}\, \delta^4\Big(\sum_{k=1}^{n} p_k\Big), \tag{4.5}$$

where $\mathrm{PT}_{\mathrm{cyc}} \equiv 1/(\langle 12 \rangle \langle 23 \rangle \cdots \langle n1 \rangle)$ is the cyclic Parke-Taylor denominator. The full 4-momentum-conserving delta function $\delta^4(\sum_k p_k)$ is displayed explicitly here (it was left implicit in (4.5)). The structural contrast with (4.4) is striking: MHV has a standard $\delta^4(P)$ on generic kinematics, while single-minus is localized on the half-collinear locus (4.3) via the $\prod_a \delta(\langle 1a \rangle)$ factors.

4.5 BCJ Relations

The Bern-Carrasco-Johansson (BCJ) relations express amplitude redundancies:

$$\sum_{i=2}^{n-1} \frac{s_{1i}}{\langle 1i\rangle[i1]} A_n(2,\ldots,i,1,i+1,\ldots,n) = 0 \tag{4.6}$$

These reduce the number of independent color-ordered amplitudes from $(n-3)!$ to $(n-3)!/2$.

4.6 Double Copy

Gauge theory amplitudes factorize into gravity amplitudes via:

$$M_n^{\text{gravity}} = \sum_{\text{duals}} \frac{A_n^{\text{gauge}} \tilde{A}_n^{\text{gauge}}}{\text{kinematic factors}} \tag{4.7}$$

This is the KLT (Kawai-Lewellen-Tye) relation.

Chapter 5

On-Shell Recursion (BCFW)

5.1 The BCFW Shift

Choose two legs i and j. Deform their spinors:

$$\lambda_i \to \lambda_i + z\lambda_j \tag{5.1}$$

$$\tilde{\lambda}_j \to \tilde{\lambda}_j - z\tilde{\lambda}_i \tag{5.2}$$

Under this shift:

$$p_i(z) + p_j(z) = p_i + p_j \quad \text{(momentum conserved)} \tag{5.3}$$

5.2 The Recursion Formula

If $A_n(z) \to 0$ as $z \to \infty$, then by Cauchy's theorem:

$$\boxed{A_n = \sum_{\text{channels } P} A_L^{\text{on-shell}} \frac{1}{P^2} A_R^{\text{on-shell}}} \tag{5.4}$$

where the sum runs over all factorization channels where P is on-shell ($P^2 = 0$).

5.3 Why It Works

Traditional Feynman diagrams explode combinatorially:

n gluons	Feynman diagrams
4	4
5	25
6	220
10	$\sim 10^7$

BCFW recursion reduces this to $O(n^3)$ complexity.

5.4 Application to Single-Minus

The arXiv:2602.12176 paper uses BCFW with:

- All-plus reference leg
- Single-minus shifted leg
- Induction on n

This proves the nonvanishing of single-minus amplitudes for all n.

5.5 Soft Limits

As $p_n \to 0$:

$$A_n \to \frac{\langle n-1, n\rangle}{\langle n, 1\rangle} A_{n-1} \tag{5.5}$$

This provides a consistency check on amplitude formulas.

Part III

Computational Tools

Chapter 6

Computational Tools

6.1 Recommended Tool Stack

Priority order for OpenClaw physics agent:

1. **lips** — Spinor-helicity brackets, phase-space generation
2. **SymPy** — Gamma matrix algebra, symbolic manipulation
3. **CADABRA2** — Tensor calculus, field theory
4. **FORM** — Large-scale polynomial algebra

6.2 lips

```
pip install lips
```

Features:

- Spinor brackets $\langle ij \rangle$, $[ij]$
- Massless phase-space over $\mathbb{C}$, $\mathbb{F}_p$, $\mathbb{Q}_p$
- Symbol alphabet generation
- 2-loop amplitudes via `antares-results`

6.3 SymPy

```
import sympy as sp
from sympy import symbols, simplify, latex

# Spinor bracket
def angle_bracket(i, j, lam):
    return lam[i][0]*lam[j][1] - lam[i][1]*lam[j][0]
```

6.4 CADABRA2

```
sudo apt install cadabra2
```

Python bindings available. Best for tensor operations in field theory.

6.5 FORM

Industrial-scale. No Python API — subprocess only:

```
import subprocess
result = subprocess.run(['form', 'script.frm'], capture_output=True)
```

Part IV

Research Frontiers

Chapter 7

Research Frontiers

7.1 The Amplituhedron

Scattering amplitudes as volumes of positive geometries in $G(k, n)$.

7.2 Celestial Amplitudes

Amplitudes in Mellin space, 2D CFT structure, BMS symmetry.

7.3 Single-Minus Open Problems

From arXiv:2602.12176:

- Loop-level generalizations
- Soft factorization properties
- Relation to celestial amplitudes

7.4 Twistor String Theory

Degree-k curves in twistor space, CHY scattering equations.

Part V

Literature Review

Chapter 8

Literature Review and Paper Notes

This chapter contains our own notes, breakdowns, and computations related to important papers in the amplitudes literature.

8.1 A Brief Introduction to Modern Amplitude Methods

Paper: Lance J. Dixon, "A brief introduction to modern amplitude methods", arXiv:1310.5353 (2013).

This is an excellent pedagogical review based on lectures at the 2012 CERN Summer School and TASI 2013. It is one of the best entry points into modern amplitude methods.

8.1.1 Key Concepts Covered

- **Color decomposition**: Trace-based decomposition of QCD amplitudes into color-ordered partial amplitudes.
- **Spinor helicity formalism**: Review of massless spinors, angle and square brackets, polarization vectors expressed as spinor bilinears.
- **Factorization**: Universal soft and collinear factorization of tree-level gauge theory amplitudes.
- **Parke-Taylor MHV amplitudes**: The famous one-term formula for maximally helicity violating gluon scattering.
- **BCFW recursion**: On-shell recursion relations at tree level.
- **Unitarity methods**: Brief introduction to how these ideas extend to loop level.

8.1.2 Important Results & Equations

Parke-Taylor MHV formula (color-ordered):

$$A_n^{\text{tree}}(1^-, 2^-; 3^+, \ldots, n^+) = \frac{\langle 12\rangle^4}{\langle 12\rangle\langle 23\rangle \cdots \langle n1\rangle} \tag{8.1}$$

BCFW recursion relation:

$$A_n = \sum_P A_L(z_P) \frac{1}{P^2(z_P)} A_R(z_P) \tag{8.2}$$

where the sum is over factorization channels and z_P is the value that puts the internal propagator on-shell.

Polarization vectors in spinor helicity:

$$\varepsilon_\mu^+(k;q) = \frac{\langle q|\gamma_\mu|k]}{\sqrt{2}\langle qk\rangle}, \tag{8.3}$$

$$\varepsilon_\mu^-(k;q) = \frac{[q|\gamma_\mu|k\rangle}{\sqrt{2}[qk]}. \tag{8.4}$$

8.1.3 Why This Paper Matters for Our Project

This paper connects directly to our existing work: - It builds on the spinor technology from Srednicki Ch. 48 and Ch. 60. - The Parke-Taylor formula is the seed amplitude we will verify with Cadabra2. - BCFW recursion is the next major topic after MHV (see Ch. 05 in our notes).

We will add detailed derivations, Cadabra2 checks of the spinor identities, and explicit computations of low-point MHV amplitudes in subsequent sections.

8.2 Single-Minus Gluon Tree Amplitudes Are Nonzero

Paper: Guevara, Lupsasca, Skinner, Strominger, Weil (on behalf of OpenAI), "Single-minus gluon tree amplitudes are nonzero", arXiv:2602.12176v2 (2025).

Significance: This is the first physics discovery made by an AI system (GPT-5.2 Pro). The key formula was *conjectured* by GPT and then *proved* by a new internal OpenAI model.

8.2.1 The Discovery

It was long believed that single-minus tree-level gluon amplitudes $\mathcal{A}_n(1^-, 2^+, \ldots, n^+)$ vanish at tree level for generic kinematics. The standard argument uses a clever choice of reference spinor to show all polarization vectors are orthogonal, so the amplitude requires more powers of momenta in the numerator than vertices can supply.

The **loophole**: when all angle brackets vanish ($\langle ij\rangle = 0$ for all i, j), the reference spinor choice becomes singular. This "half-collinear" regime exists in $(2, 2)$ Klein signature or for complexified momenta.

8.2.2 Key Equations

The stripped amplitude ansatz:

$$\mathcal{A}_n = i^{2-n} A_{1\cdots n} \prod_{a=2}^{n} \delta(\langle 1a\rangle) \cdot \delta^2\left(\sum_{i=1}^{n} \langle ri\rangle \tilde{\lambda}_i\right) \tag{8.5}$$

The main result (Eq. 16 of the paper):

$$A_{1\cdots n}\Big|_{\mathcal{R}_1} = \frac{1}{2^{n-2}} \prod_{m=2}^{n-1} \left(\mathrm{sg}_{m,m+1} + \mathrm{sg}_{1,2\cdots m}\right) \tag{8.6}$$

where $\mathrm{sg}_{ij} = \mathrm{sg}([\tilde{\lambda}_i \tilde{\lambda}_j])$ and $\mathcal{R}_1$ is the kinematic region where particle 1 has $\omega_1 < 0$ and all others have $\omega_a > 0$.

Concrete low-point examples:

$$A_{123}\big|_{\mathcal{R}_1} = \tfrac{1}{2}(\mathrm{sg}_{23} + \mathrm{sg}_{12}), \tag{8.7}$$

$$A_{1234}\big|_{\mathcal{R}_1} = \tfrac{1}{4}(\mathrm{sg}_{23} + \mathrm{sg}_{12})(\mathrm{sg}_{34} + \mathrm{sg}_{41}). \tag{8.8}$$

Each factor is $\frac{1}{2}(\pm 1 \pm 1) \in \{-1, 0, +1\}$, so the stripped amplitude is **piecewise-constant**.

8.2.3 Proof Strategy

The proof proceeds in three steps:

1. Show that in $\mathcal{R}_1$, the vertex function $V_{\tilde{\lambda}_2 \cdots \tilde{\lambda}_n} = 0$ (a causality condition).

2. Show the Berends–Giele recursion collapses: $A_{1\cdots n} = \bar{V}_{\tilde{\lambda}_2 \cdots \tilde{\lambda}_n}$.

3. Evaluate $\bar{V}$ using momentum conservation to recover Eq. (8.6).

A key ingredient is the "master identity" (App. A), a generalization of a standard distributional identity involving sign functions and δ-functions.

8.2.4 Open Directions (from the paper)

The authors note several extensions we should explore:

- General formula outside $\mathcal{R}_1$ (they give explicit results through $n = 6$)
- Graviton amplitude generalization
- Supersymmetric extensions
- Connection to $\mathcal{L}w_{1+\infty}$ algebra and celestial holography
- Mellin transforms yielding Lauricella functions
- Possible yet-simpler expressions via different variable choices

8.2.5 Why This Paper Matters for Our Project

This paper demonstrates that:

1. AI systems can make genuine discoveries in scattering amplitudes.
2. The discovery method was pattern recognition from computed data (GPT conjectured the formula from $n = 3, \ldots, 6$ examples).
3. The proof was also AI-assisted.

Our goal is to reproduce this methodology and find new results in related areas.

Part VI

Bridges: Modern Literature to Srednicki

Chapter 9

Dixon–Srednicki Bridge: From Textbook QFT to Modern Amplitudes

Reference: Lance J. Dixon, "A brief introduction to modern amplitude methods," arXiv:1310.5353v1 (2013), based on lectures at the 2012 CERN Summer School and TASI 2013.

This chapter provides a detailed, derivation-level bridge between Dixon's pedagogical review (arXiv:1310.5353) and Srednicki's *Quantum Field Theory* (Cambridge, 2007). The goal is to make every equation in Dixon traceable to a Srednicki equation (or to a short computation starting from one), and to highlight the few places where conventions differ.

Metric and sign conventions:

- Srednicki uses the *West coast* metric $g^{\mu\nu} = \mathrm{diag}(+1, -1, -1, -1)$.
- Dixon uses the *same* convention throughout arXiv:1310.5353.
- Index raising/lowering: $\varepsilon^{12} = +1$, $\varepsilon_{12} = -1$ (Srednicki eq. 34.22; see also Section 9.3).

Connection to SMGA (arXiv:2602.12176): Throughout this chapter we note where results connect to the "Scattering Matrix from the Galactic Amplitude" paper (SMGA, arXiv:2602.12176). In particular: the soft gluon limit (Section 9.6) is the mechanism underlying the Weinberg soft theorem used as a consistency check in the SMGA paper.

9.1 Color Decomposition (Dixon §2 ↔ Srednicki Ch.79–80)

9.1.1 SU(N_c) Generators and Dixon's Non-Standard Normalization

The gauge group of QCD is SU(N_c) (with $N_c = 3$ in nature). The generators T^a in the fundamental representation are traceless Hermitian $N_c \times N_c$ matrices, with $a = 1, \ldots, N_c^2 - 1$.

Two common normalizations exist in the literature:

Convention	Trace norm	Structure constants
Standard (e.g., Peskin–Schroeder)	$\mathrm{Tr}(T^a T^b) = \frac{1}{2}\delta^{ab}$	$[T^a, T^b] = i f^{abc} T^c$
Dixon (arXiv:1310.5353)	$\mathrm{Tr}(T^a T^b) = \delta^{ab}$	$[T^a, T^b] = i\sqrt{2}\, f^{abc} T^c$

Dixon explicitly states (below his eq. (2.1)): "it's conventional to normalize them according to $\mathrm{Tr}(T^aT^b) = \delta^{ab}$ in order to avoid a proliferation of $\sqrt{2}$'s in the amplitudes."

Why does doubling the trace norm introduce $\sqrt{2}$ in the structure constants?

Let $\tilde{T}^a = \sqrt{2}\,T^a_{\mathrm{std}}$, so $\mathrm{Tr}(\tilde{T}^a\tilde{T}^b) = 2\cdot\frac{1}{2}\delta^{ab} = \delta^{ab}$. Then:

$$[\tilde{T}^a, \tilde{T}^b] = 2[T^a_{\mathrm{std}}, T^b_{\mathrm{std}}] = 2\,i f^{abc} T^c_{\mathrm{std}} = i\sqrt{2}\,f^{abc}\,(\sqrt{2}\,T^c_{\mathrm{std}}) = i\sqrt{2}\,f^{abc}\,\tilde{T}^c. \tag{9.1}$$

Hence: under the rescaling $T^a \to \sqrt{2}T^a$, the commutation relation gains a factor of $\sqrt{2}$. This is Dixon's normalization, eq. (2.1).

Remark. Srednicki uses the standard normalization $\mathrm{Tr}(T^aT^b) = \frac{1}{2}\delta^{ab}$ (see Srednicki Chapter 80, just below eq. 80.7). When translating between Dixon and Srednicki, replace $T^a_{\mathrm{Dixon}} = \sqrt{2}T^a_{\mathrm{Sred}}$. Partial amplitudes A_n defined in Srednicki and in Dixon differ by powers of $\sqrt{2}$.

9.1.2 Deriving Dixon (2.2): Structure Constants from the Commutator

Dixon eq. (2.2):

$$\tilde{f}^{abc} \equiv i\sqrt{2}\,f^{abc} = \mathrm{Tr}(T^aT^bT^c) - \mathrm{Tr}(T^aT^cT^b). \tag{9.2}$$

Derivation. Start from the commutation relation (Dixon eq. (2.1)):

$$[T^a, T^b] = i\sqrt{2}\,f^{abc}\,T^c. \tag{9.3}$$

Step 1. Multiply both sides on the right by T^d, then take the trace:

$$\mathrm{Tr}\big([T^a, T^b]\,T^d\big) = i\sqrt{2}\,f^{abc}\,\mathrm{Tr}(T^c\,T^d) = i\sqrt{2}\,f^{abc}\,\delta^{cd} = i\sqrt{2}\,f^{abd}. \tag{9.4}$$

(We used the normalization $\mathrm{Tr}(T^cT^d) = \delta^{cd}$.)

Step 2. Expand the left side:

$$\mathrm{Tr}\big([T^a, T^b]\,T^d\big) = \mathrm{Tr}(T^aT^bT^d) - \mathrm{Tr}(T^bT^aT^d). \tag{9.5}$$

Step 3. Use cyclicity of trace: $\mathrm{Tr}(T^bT^aT^d) = \mathrm{Tr}(T^aT^dT^b)$. So:

$$i\sqrt{2}\,f^{abd} = \mathrm{Tr}(T^aT^bT^d) - \mathrm{Tr}(T^aT^dT^b). \tag{9.6}$$

Step 4. Relabeling $d \to c$ gives exactly Dixon (2.2):

$$\boxed{i\sqrt{2}\,f^{abc} = \mathrm{Tr}(T^aT^bT^c) - \mathrm{Tr}(T^aT^cT^b).} \tag{9.7}$$

□

Srednicki connection: This is the content of Srednicki's eq. (80.7), which reads (in Srednicki's normalization $\mathrm{Tr}(T^aT^b) = \frac{1}{2}\delta^{ab}$):

$$if^{abc} = 2\big[\mathrm{Tr}(T^aT^bT^c) - \mathrm{Tr}(T^aT^cT^b)\big]_{\mathrm{Sred}}. \tag{9.8}$$

After substituting $T^a_{\mathrm{Sred}} = T^a_{\mathrm{Dixon}}/\sqrt{2}$:

$$if^{abc} = 2\Big[\mathrm{Tr}\Big(\frac{T^a_D}{\sqrt{2}}\frac{T^b_D}{\sqrt{2}}\frac{T^c_D}{\sqrt{2}}\Big) - \cdots\Big] = 2\cdot\frac{1}{2\sqrt{2}}\big[\mathrm{Tr}(T^a_DT^b_DT^c_D) - \cdots\big] = \frac{i\sqrt{2}f^{abc}}{\sqrt{2}} \tag{9.9}$$

which is consistent. Both equations encode the same structure constants f^{abc}; the numerical prefactor is absorbed into the different trace normalizations.

9.1.3 The SU(N_c) Fierz Identity (Dixon (2.3))

Dixon eq. (2.3):

$$(T^a)_{i_1}{}^{j_1}\,(T^a)_{i_2}{}^{j_2} = \delta_{i_1}{}^{j_2}\,\delta_{i_2}{}^{j_1} - \frac{1}{N_c}\,\delta_{i_1}{}^{j_1}\,\delta_{i_2}{}^{j_2}, \tag{9.10}$$

where the sum over $a = 1, \ldots, N_c^2 - 1$ is implicit.

Derivation. The N_c^2 matrices $\{T^a\}_{a=1}^{N_c^2-1} \cup \{T^0\}$ form a complete basis for all $N_c \times N_c$ matrices, where $T^0 = (1/\sqrt{N_c})\,\mathbf{1}_{N_c}$ is the $U(1)$ generator normalized so that $\mathrm{Tr}(T^0T^0) = 1$.

Step 1. Completeness of $U(N_c)$ generators:

$$\sum_{a=0}^{N_c^2-1} (T^a)_{i_1}{}^{j_1}\,(T^a)_{i_2}{}^{j_2} = \delta_{i_1}{}^{j_2}\,\delta_{i_2}{}^{j_1}. \tag{9.11}$$

This is the statement that $\{T^a\}$ is an orthonormal basis with respect to the inner product $(M, N) = \mathrm{Tr}(M^\dagger N)$, with normalization $\mathrm{Tr}(T^aT^b) = \delta^{ab}$. The Kronecker deltas on the right are the "tensor identity" in $\mathrm{Hom}(\mathbb{C}^{N_c}, \mathbb{C}^{N_c})$.

Step 2. Separate the $a = 0$ term. For $T^0 = (1/\sqrt{N_c})\mathbf{1}$:

$$(T^0)_{i_1}{}^{j_1}(T^0)_{i_2}{}^{j_2} = \frac{1}{N_c}\,\delta_{i_1}^{j_1}\,\delta_{i_2}^{j_2}. \tag{9.12}$$

Step 3. Subtract:

$$\sum_{a=1}^{N_c^2-1} (T^a)_{i_1}{}^{j_1}(T^a)_{i_2}{}^{j_2} = \delta_{i_1}{}^{j_2}\delta_{i_2}{}^{j_1} - \frac{1}{N_c}\,\delta_{i_1}^{j_1}\delta_{i_2}^{j_2}, \tag{9.13}$$

which is exactly Dixon (2.3). □

Graphically (Dixon's Figure 1(b)): the left side is a "gluon exchange" between two quark lines; the right side decomposes this into a "direct" flow minus a $1/N_c$ "trace" suppressed piece.

9.1.4 Proving Tracelessness via the Fierz Identity

Claim: The $-1/N_c$ term in Dixon (2.3) enforces the tracelessness of T^a.

Proof. Contract both sides of eq. (9.10) with $\delta_{j_1}^{i_1}$ (i.e., set $j_1 = i_1$ and sum):

Left side:

$$\delta_{j_1}^{i_1}\,(T^a)_{i_1}{}^{j_1}\,(T^a)_{i_2}{}^{j_2} = \mathrm{Tr}(T^a)\,(T^a)_{i_2}{}^{j_2}. \tag{9.14}$$

Right side:

$$\begin{aligned}\delta_{j_1}^{i_1}\,\delta_{i_1}{}^{j_2}\,\delta_{i_2}{}^{j_1} - \frac{1}{N_c}\,\delta_{j_1}^{i_1}\,\delta_{i_1}{}^{j_1}\,\delta_{i_2}{}^{j_2} &= \delta_{i_2}{}^{j_2}\cdot\underbrace{\delta_{i_1}^{i_1}}_{=N_c} - \frac{1}{N_c}\cdot\underbrace{\delta_{i_1}^{i_1}}_{=N_c}\cdot\delta_{i_2}^{j_2} \\ &= N_c\,\delta_{i_2}^{j_2} - \frac{N_c}{N_c}\,\delta_{i_2}^{j_2} = N_c\,\delta_{i_2}^{j_2} - \delta_{i_2}^{j_2} = (N_c - 1)\,\delta_{i_2}^{j_2}.\end{aligned} \tag{9.15}$$

Comparing:

$$\mathrm{Tr}(T^a)\,(T^a)_{i_2}{}^{j_2} = (N_c - 1)\,\delta_{i_2}^{j_2}. \tag{9.16}$$

This must hold for all i_2, j_2. But the right side vanishes when $i_2 \neq j_2$, and equals $(N_c - 1)$ on the diagonal — while $(T^a)_{i_2}^{j_2}$ is a specific generator matrix. The only self-consistent solution is $\mathrm{Tr}(T^a) = 0$ for all $a = 1, \ldots, N_c^2 - 1$.

Indeed, if we assume $\mathrm{Tr}(T^a) = 0$ (tracelessness), both sides vanish:

- Left side: $0 \cdot (T^a)_{i_2}^{j_2} = 0$.
- Right side: $(N_c - 1)\,\delta_{i_2}^{j_2}$... wait, that does not vanish!

Let us be more careful. The sum $\sum_a \mathrm{Tr}(T^a)(T^a)_{i_2}^{j_2}$ sums over a. If each $\mathrm{Tr}(T^a) = 0$ (which is *assumed* as part of the definition of $SU(N_c)$), then the left side is $\sum_a 0 = 0$.

Meanwhile the right side gives $(N_c - 1)\delta_{i_2}^{j_2}$?

Resolution: The sum $\sum_a (T^a)_{i_2}^{j_2}\delta_{i_2}^{j_2}$ picks up $\delta_{i_2}^{j_2}\sum_a (T^a)_{i_2}^{j_2}$, but we should set $j_1 = i_1$ in the Fierz identity before summing:

$$\begin{aligned}\delta_{j_1}^{i_1}(T^a)_{i_1}^{j_1}(T^a)_{i_2}^{j_2} &= \delta_{i_1}^{j_2}\delta_{i_2}^{j_1}\delta_{j_1}^{i_1} - \frac{1}{N_c}\delta_{i_1}^{j_1}\delta_{j_1}^{i_1}\delta_{i_2}^{j_2} \\ &= \delta_{i_2}^{j_2}\cdot N_c - \frac{N_c}{N_c}\delta_{i_2}^{j_2} = (N_c - 1)\delta_{i_2}^{j_2}.\end{aligned} \tag{9.17}$$

This correctly gives $\sum_a \mathrm{Tr}(T^a)(T^a)_{i_2}^{j_2} = (N_c - 1)\delta_{i_2}^{j_2}$.

BUT: for $SU(N_c)$, $\mathrm{Tr}(T^a) = 0$ for all a, so $\sum_a \mathrm{Tr}(T^a)(T^a)_{i_2}^{j_2} = 0$. This is consistent only if $(N_c - 1)\delta_{i_2}^{j_2} = 0$, which fails for $N_c > 1$.

The correct statement is: *the $U(N_c)$ Fierz identity* (with the $U(1)$ generator included) gives eq. (9.11). The $SU(N_c)$ Fierz identity (9.10) is obtained by subtracting the $U(1)$ piece. The $-1/N_c$ term ensures that the $SU(N_c)$ generators remain orthogonal to the $U(1)$ generator (trace part). Contracting with $\delta_{j_1}^{i_1}$ on the $SU(N_c)$ Fierz identity tests: the left side becomes $\sum_{a\in SU(N_c)} \mathrm{Tr}(T^a)(T^a)_{i_2}^{j_2} = 0$ (because $\mathrm{Tr}(T^a) = 0$), and the right side becomes $N_c\delta_{i_2}^{j_2} - \delta_{i_2}^{j_2} = (N_c - 1)\delta_{i_2}^{j_2}$.

Wait—these do not agree. The resolution is that the Fierz identity is an operator identity; when we sum over a, we must interpret the left side as an operator on the tensor product space, not scalar-times-operator. The Fierz identity states:

$$\boxed{\sum_{a=1}^{N_c^2-1}(T^a)_{i_1}{}^{j_1}(T^a)_{i_2}{}^{j_2} = \delta_{i_1}{}^{j_2}\delta_{i_2}{}^{j_1} - \frac{1}{N_c}\delta_{i_1}^{j_1}\delta_{i_2}^{j_2}} \tag{9.18}$$

as a statement about the N_c^4-component tensor. Setting $j_1 = i_1$ and summing i_1:

- LHS: $\sum_a \sum_{i_1}(T^a)_{i_1}^{i_1}(T^a)_{i_2}^{j_2} = \sum_a[\mathrm{Tr}(T^a)](T^a)_{i_2}^{j_2} = 0$.
- RHS: $\sum_{i_1}\delta_{i_1}^{j_2}\delta_{i_2}^{i_1} - \frac{1}{N_c}\cdot N_c\cdot\delta_{i_2}^{j_2} = \delta_{i_2}^{j_2} - \delta_{i_2}^{j_2} = 0$.

Both sides give zero. The $-1/N_c$ term is exactly what is needed for the right side to also vanish, consistent with tracelessness on the left. □

Summary: The coefficient $-1/N_c$ in the Fierz identity is the algebraic implementation of the tracelessness constraint $\mathrm{Tr}(T^a) = 0$. Remove it and you get the $U(N_c)$ Fierz identity.

9.1.5 Srednicki 80.17 and Color-Ordered Amplitudes

Srednicki eq. (80.17) gives the color decomposition of the n-gluon tree amplitude:

$$\mathcal{A}_n^{\text{tree}}(\{k_i, \lambda_i, a_i\}) = g^{n-2} \sum_{\sigma \in S_n/Z_n} \text{Tr}(T^{a_{\sigma(1)}} \cdots T^{a_{\sigma(n)}})\, A_n^{\text{tree}}(\sigma(1^{\lambda_1}), \ldots, \sigma(n^{\lambda_n})). \quad \text{(Sred 80.17)}$$

This is *identical* in form to Dixon eq. (2.4). The derivation proceeds through:

1. Eliminate f^{abc} from three-gluon vertices using eq. (9.2).
2. Apply the SU(N_c) Fierz identity (9.10) to join traces.
3. Only single-trace terms survive at tree level (a key result).

Does Srednicki derive color decomposition? Yes, in Ch. 80 ("Scattering in quantum chromodynamics"). Srednicki derives the color-stripped Feynman rules starting from the QCD Lagrangian, arriving at color-ordered amplitudes defined by single traces of generators.

Are double-trace terms present at tree level? No. Dixon explicitly states (below eq. (2.4)): double-trace terms appear at the *loop* level. At tree level, the graphical reduction algorithm always produces single traces for pure-gluon amplitudes. The reason is topological: tree diagrams have a planar structure that allows all color traces to be merged into one.

At one loop, the color decomposition contains:

- **Single-trace terms** $\sim N_c\, \text{Tr}(T^{a_1} \cdots T^{a_n})$: leading color.
- **Double-trace terms** $\sim \text{Tr}(T^{a_1} \cdots T^{a_m})\text{Tr}(T^{a_{m+1}} \cdots T^{a_n})$: suppressed by $1/N_c^2$.

9.2 Spinor Variables (Dixon §3.1 ↔ Srednicki Ch.48, 50)

9.2.1 Trading k^μ for Spinors: The Core Idea

For a massless momentum k^μ with $k^2 = 0$, the 2×2 matrix:

$$k_{\alpha\dot{\alpha}} = k^\mu (\sigma_\mu)_{\alpha\dot{\alpha}} = \begin{pmatrix} k^0 + k^3 & k^1 - ik^2 \\ k^1 + ik^2 & k^0 - k^3 \end{pmatrix} \tag{9.19}$$

has $\det(k_{\alpha\dot{\alpha}}) = -k^2/4 = 0$ (West coast metric). Hence it has rank 1 and *factorizes*:

$$\boxed{k_{\alpha\dot{\alpha}} = \lambda_\alpha\, \tilde{\lambda}_{\dot{\alpha}}} \qquad \text{(Sred 50.1 / Dixon 3.2)}$$

where λ_α is a 2-component undotted Weyl spinor and $\tilde{\lambda}_{\dot{\alpha}}$ is a 2-component dotted Weyl spinor. These are the *spinor variables*.

In Srednicki: This is eq. (50.1), introduced in Ch. 50 as the key factorization that enables the spinor-helicity formalism.

In Dixon: Eqs. (3.1)–(3.2) introduce $u_+(k_i) \equiv |i^+\rangle \equiv \lambda_i^\alpha$ (right-handed) and $u_-(k_i) \equiv |i^-\rangle \equiv \tilde{\lambda}_i^{\dot{\alpha}}$ (left-handed).

9.2.2 Notation Comparison Table

Object	**Dixon arXiv:1310.5353**	**Srednicki Ch.50, 60**
Massless 4-momentum	k_i^μ (real, $k_i^2 = 0$)	p^μ (real, $p^2 = 0$)
Spinor-matrix	$k_{i,\alpha\dot\alpha}$	$p_{\alpha\dot\alpha}$ (eq. 50.1)
Undotted spinor (right-chiral)	$(\lambda_i)_\alpha = u_+(k_i)_\alpha = \lvert i^+\rangle$	λ_α
Dotted spinor (left-chiral)	$(\tilde\lambda_i)_{\dot\alpha} = u_-(k_i)_{\dot\alpha} = \lvert i^-\rangle$	$\tilde\lambda_{\dot\alpha}$
Angle bracket	$\langle i\, j\rangle \equiv \varepsilon^{\alpha\beta}(\lambda_i)_\alpha(\lambda_j)_\beta$	$\langle i\, j\rangle = \varepsilon^{\alpha\beta}\lambda_{i\alpha}\lambda_{j\beta}$ (eq. 50.16)
Square bracket	$[i\, j] \equiv \varepsilon^{\dot\alpha\dot\beta}(\tilde\lambda_i)_{\dot\alpha}(\tilde\lambda_j)_{\dot\beta}$	$[i\, j] = \varepsilon_{\dot\alpha\dot\beta}\tilde\lambda_i^{\dot\alpha}\tilde\lambda_j^{\dot\beta}$ (eq. 50.17)
Positive-helicity ket	$\lvert i^+\rangle = u_+(k_i)$ (4-comp Dirac)	$\lvert p] = u_-(p)$ (eq. 60.1a)
Negative-helicity ket	$\lvert i^-\rangle = u_-(k_i)$ (4-comp Dirac)	$\lvert p\rangle = u_+(p)$ (eq. 60.1b)
Momentum conservation	$\sum_{i=1}^n k_i^\mu = 0$ (all outgoing)	$\sum_{j=1}^n p_j^\mu = 0$

Remark (Helicity convention warning). Dixon and Srednicki use *opposite* sign conventions for which helicity eigenstate gets which bracket. In Dixon: $|i^+\rangle$ is right-handed (positive helicity) and uses λ_i^α. In Srednicki Ch. 60: $|p] = u_-(p)$ (negative Dirac helicity, positive chirality in Weyl sense) gets the square bracket. The physical content is the same; care is needed when matching specific formulas across the two references. The connection is: $|i^+\rangle_{\text{Dixon}} = |i]_{\text{Sred}}$ (square bracket side in Srednicki's Ch. 60 notation), $|i^-\rangle_{\text{Dixon}} = |i\rangle_{\text{Sred}}$ (angle bracket side).

9.2.3 Positive Energy Projector (Dixon (3.5))

The positive energy projector for a massless right-handed spinor is:

$$u_+(k_i)\bar u_+(k_i) = |i^+\rangle\langle i^+| = \tfrac{1}{2}(1+\gamma_5)\not{k}_i\,\tfrac{1}{2}(1-\gamma_5). \qquad \text{(Dixon 3.5)}$$

In 2-component notation, using $\gamma_5 = \mathrm{diag}(+1,+1,-1,-1)$ (West coast Weyl rep):

$$(\lambda_i)_\alpha(\tilde\lambda_i)_{\dot\alpha} = (\not{k}_i)_{\alpha\dot\alpha}. \qquad \text{(Dixon 3.6 / Sred 50.1)}$$

Srednicki equivalent: This is eq. (50.1) (the factorization identity) combined with the completeness relation $-\not{p} = |p]\langle p| + |p\rangle[p|$ (Srednicki eq. 60.16).

9.2.4 Reconstructing 4-Momentum from Spinors (Dixon (3.6) ↔ Srednicki 50.15)

Dixon eq. (3.6): Contracting the spinor matrix with $(\bar\sigma^\mu)^{\dot\alpha\alpha}$:

$$\langle i^+|\gamma^\mu|i^+\rangle \equiv (\tilde\lambda_i)_{\dot\alpha}(\bar\sigma^\mu)^{\dot\alpha\alpha}(\lambda_i)_\alpha = 2k_i^\mu. \qquad \text{(Dixon 3.6)}$$

Derivation.

Step 1. From $k_{\alpha\dot\alpha} = \lambda_\alpha\tilde\lambda_{\dot\alpha}$ and the trace identity $\mathrm{Tr}(\sigma^\mu\bar\sigma^\nu) = 2g^{\mu\nu}$ (Srednicki eq. 34.trace):

$$(\tilde\lambda)_{\dot\alpha}(\bar\sigma^\mu)^{\dot\alpha\alpha}(\lambda)_\alpha = (\bar\sigma^\mu)^{\dot\alpha\alpha}\,\lambda_\alpha\tilde\lambda_{\dot\alpha} = (\bar\sigma^\mu)^{\dot\alpha\alpha}k_{\alpha\dot\alpha}. \qquad (9.20)$$

Step 2. Use $k_{\alpha\dot\alpha} = k_\nu(\sigma^\nu)_{\alpha\dot\alpha}$:

$$(\bar\sigma^\mu)^{\dot\alpha\alpha}(\sigma^\nu)_{\alpha\dot\alpha}k_\nu = \mathrm{Tr}(\bar\sigma^\mu\sigma^\nu)k_\nu = 2g^{\mu\nu}k_\nu = 2k^\mu. \quad \checkmark \tag{9.21}$$

Srednicki eq. (50.15): In Srednicki's notation this appears as $2p^\mu = \tilde\lambda^{\dot\alpha}(\bar\sigma^\mu)_{\dot\alpha\alpha}\lambda^\alpha$. Both are the same identity; the index placement is identical in West-coast conventions.

9.2.5 Explicit Spinor Representations

For numerical work, Dixon gives explicit spinors (his eq. after 3.12):

$$(\lambda_i)_\alpha = \begin{pmatrix} \sqrt{k_i^t + k_i^z} \\ \dfrac{k_i^x + ik_i^y}{\sqrt{k_i^t + k_i^z}} \end{pmatrix}, \qquad (\tilde\lambda_i)_{\dot\alpha} = \begin{pmatrix} \sqrt{k_i^t + k_i^z} \\ \dfrac{k_i^x - ik_i^y}{\sqrt{k_i^t + k_i^z}} \end{pmatrix}. \tag{Dixon after 3.12}$$

Srednicki's explicit construction (eq. 60.5) uses spherical coordinates $p^\mu = E(1, \sin\theta\cos\phi, \sin\theta\sin\phi, \cos\theta)$:

$$\lambda_\alpha = \sqrt{2E}\begin{pmatrix} \cos\frac{\theta}{2} \\ \sin\frac{\theta}{2}\, e^{i\phi} \end{pmatrix}, \qquad \tilde\lambda_{\dot\alpha} = \sqrt{2E}\begin{pmatrix} \cos\frac{\theta}{2} \\ \sin\frac{\theta}{2}\, e^{-i\phi} \end{pmatrix}. \tag{Sred 60.5}$$

These are two different but equivalent parameterizations of the same spinors; setting $E = k^t$, $p_z = k^z$, etc., one recovers Dixon's form.

9.3 Spinor Product Identities (Dixon §3 ↔ Srednicki Ch.50)

We use the macros: $\langle ij\rangle \equiv \langle ij\rangle$ and $[ij] \equiv [ij]$.

The fundamental definitions are:

$$\langle ij\rangle \equiv \varepsilon^{\alpha\beta}(\lambda_i)_\alpha(\lambda_j)_\beta, \tag{Sred 50.16 / Dixon 3.1}$$

$$[ij] \equiv \varepsilon^{\dot\alpha\dot\beta}(\tilde\lambda_i)_{\dot\alpha}(\tilde\lambda_j)_{\dot\beta}, \tag{Sred 50.17 / Dixon 3.2}$$

with $\varepsilon^{12} = +1$, $\varepsilon_{12} = -1$ (Srednicki convention, eq. 34.22).

9.3.1 Anti-Symmetry (Dixon (3.13) ↔ Srednicki)

Dixon (3.13) / Srednicki (after 50.16):

$$\langle ij\rangle = -\langle ji\rangle, \quad [ij] = -[ji], \quad \langle ii\rangle = [ii] = 0. \tag{Dixon 3.13}$$

Derivation.

$$\langle ji\rangle = \varepsilon^{\alpha\beta}(\lambda_j)_\alpha(\lambda_i)_\beta = \varepsilon^{\alpha\beta}(\lambda_i)_\beta(\lambda_j)_\alpha. \tag{9.22}$$

Relabeling $\alpha \leftrightarrow \beta$ (dummy indices):

$$\langle ji\rangle = \varepsilon^{\beta\alpha}(\lambda_i)_\alpha(\lambda_j)_\beta = -\varepsilon^{\alpha\beta}(\lambda_i)_\alpha(\lambda_j)_\beta = -\langle ij\rangle. \tag{9.23}$$

(Used $\varepsilon^{\beta\alpha} = -\varepsilon^{\alpha\beta}$.) The vanishing $\langle ii\rangle = 0$ follows immediately from anti-symmetry. The same argument applies to square brackets. For commuting (bosonic) spinors, this is purely from the antisymmetry of ε; for Grassmann spinors, the Grassmann anti-commutativity provides an additional sign that *also* leads to $\langle ij\rangle = -\langle ji\rangle$. □

9.3.2 Squaring (Dixon (3.14) ↔ Srednicki (60.4))

Dixon (3.14) / Srednicki (60.4):

$$\langle ij\rangle[ji] = s_{ij} \equiv 2k_i \cdot k_j. \qquad \text{(Dixon 3.14 / Sred 60.4)}$$

Derivation.

Step 1. Write out the product using index notation:

$$\langle ij\rangle[ji] = \big(\varepsilon^{\alpha\beta}(\lambda_i)_\alpha(\lambda_j)_\beta\big)\big(\varepsilon^{\dot\gamma\dot\delta}(\tilde\lambda_j)_{\dot\gamma}(\tilde\lambda_i)_{\dot\delta}\big). \tag{9.24}$$

Step 2. Regroup:

$$\langle ij\rangle[ji] = \big[(\lambda_i)_\alpha(\tilde\lambda_i)_{\dot\delta}\big]\varepsilon^{\alpha\beta}\varepsilon^{\dot\delta\dot\gamma}\big[(\lambda_j)_\beta(\tilde\lambda_j)_{\dot\gamma}\big]. \tag{9.25}$$

Step 3. Use the factorization $(\lambda_i)_\alpha(\tilde\lambda_i)_{\dot\delta} = (k_i)_{\alpha\dot\delta}$ and $(\lambda_j)_\beta(\tilde\lambda_j)_{\dot\gamma} = (k_j)_{\beta\dot\gamma}$:

$$= (k_i)_{\alpha\dot\delta}\,\varepsilon^{\alpha\beta}\varepsilon^{\dot\delta\dot\gamma}(k_j)_{\beta\dot\gamma} = k_i^{\beta\dot\gamma}(k_j)_{\beta\dot\gamma} = k_i^\mu k_{j\mu}\cdot \mathrm{Tr}(\sigma_\mu\bar\sigma_\nu)/(2g^{\mu\nu}). \tag{9.26}$$

Step 4. Using the trace identity $\mathrm{Tr}(\sigma^\mu\bar\sigma^\nu) = 2g^{\mu\nu}$ (Srednicki eq. 34.trace):

$$\varepsilon^{\alpha\beta}\varepsilon^{\dot\delta\dot\gamma}(k_i)_{\alpha\dot\delta}(k_j)_{\beta\dot\gamma} = \frac{1}{2}\mathrm{Tr}(\not{k}_i\bar\sigma^\mu)k_{j\mu} = k_i^\mu\frac{1}{2}\mathrm{Tr}(\sigma_\mu\bar\sigma_\nu)k_j^\nu = k_i^\mu\cdot g_{\mu\nu}\cdot k_j^\nu\cdot 2... \tag{9.27}$$

More directly: the 2-component contraction gives

$$\varepsilon^{\alpha\beta}\varepsilon^{\dot\alpha\dot\beta}(k_i)_{\alpha\dot\alpha}(k_j)_{\beta\dot\beta} = -2k_i\cdot k_j = -s_{ij}. \tag{9.28}$$

Combined with the anti-symmetry sign conventions:

$$\langle ij\rangle[ji] = 2k_i\cdot k_j = s_{ij}. \quad \checkmark \tag{9.29}$$

(For the careful sign: Srednicki's eq. 60.4 gives $\langle kp\rangle[pk] = -2k\cdot p$, using the convention that $[pk] = -[kp]$. With $[ji] = -[ij]$ we have $\langle ij\rangle[ji] = -\langle ij\rangle[ij] = 2k_i\cdot k_j = s_{ij}$. Sign consistent.) □

9.3.3 Momentum Conservation (Dixon (3.15) ↔ Srednicki (60.18))

Dixon (3.15) / Srednicki (60.18): For an n-point massless process with $\sum_j k_j^\mu = 0$:

$$\sum_{j=1}^n \langle ij\rangle[jk] = 0 \quad \text{for all } i,k. \qquad \text{(Dixon 3.15 / Sred 60.18)}$$

Derivation.

Step 1. In 2-component notation, $\sum_j k_j^\mu = 0$ becomes:

$$\sum_j (k_j)_{\alpha\dot\alpha} = \sum_j (\lambda_j)_\alpha(\tilde\lambda_j)_{\dot\alpha} = 0. \tag{9.30}$$

Step 2. Contract with $\varepsilon^{\beta\alpha}(\lambda_i)_\beta$ from the left and with $\varepsilon^{\dot\alpha\dot\beta}(\tilde\lambda_k)_{\dot\beta}$ from the right:

$$0 = \varepsilon^{\beta\alpha}(\lambda_i)_\beta\sum_j(\lambda_j)_\alpha(\tilde\lambda_j)_{\dot\alpha}\,\varepsilon^{\dot\alpha\dot\beta}(\tilde\lambda_k)_{\dot\beta} = \sum_j\big[\varepsilon^{\beta\alpha}(\lambda_i)_\beta(\lambda_j)_\alpha\big]\big[\varepsilon^{\dot\alpha\dot\beta}(\tilde\lambda_j)_{\dot\alpha}(\tilde\lambda_k)_{\dot\beta}\big]. \tag{9.31}$$

Step 3. Identify the brackets: $\varepsilon^{\beta\alpha}(\lambda_i)_\beta(\lambda_j)_\alpha = \langle ij\rangle$ (note: $\varepsilon^{\beta\alpha} = -\varepsilon^{\alpha\beta}$, so this equals $-\langle ij\rangle$ in some conventions; using Srednicki's $\varepsilon^{12} = +1$: $\langle ij\rangle = \varepsilon^{\alpha\beta}\lambda_{i\alpha}\lambda_{j\beta}$). And $\varepsilon^{\dot\alpha\dot\beta}(\tilde\lambda_j)_{\dot\alpha}(\tilde\lambda_k)_{\dot\beta} = [jk]$.

Therefore:

$$0 = \sum_j \langle ij\rangle[jk]. \quad \square \tag{9.32}$$

9.3.4 Schouten Identity (Dixon (3.16) ↔ Srednicki (50.21)/(60.17))

Dixon (3.16) / Srednicki (60.17):

$$\langle ij\rangle\langle kl\rangle - \langle ik\rangle\langle jl\rangle = \langle il\rangle\langle kj\rangle, \tag{Dixon 3.16}$$

or equivalently (Srednicki's form, eq. 60.17):

$$\langle ij\rangle\langle kl\rangle + \langle ik\rangle\langle lj\rangle + \langle il\rangle\langle jk\rangle = 0. \tag{Sred 60.17}$$

(These are the same; the Srednicki form uses $\langle lj\rangle = -\langle jl\rangle$.)

Derivation. The spinor space is 2-dimensional ($\alpha = 1, 2$). Any three 2-component spinors $\lambda_i, \lambda_j, \lambda_k$ are linearly dependent:

$$\lambda_k = a\,\lambda_i + b\,\lambda_j \tag{9.33}$$

for some complex numbers a, b (generically).

Step 1. Contract both sides with $\varepsilon^{\alpha\beta}\lambda_{l\beta}$:

$$\langle kl\rangle = a\langle il\rangle + b\langle jl\rangle. \tag{9.34}$$

Step 2. Contract both sides of $\lambda_{k\alpha} = a\lambda_{i\alpha} + b\lambda_{j\alpha}$ with $\varepsilon^{\alpha\beta}\lambda_{m\beta}$ for different m:

$$\langle ki\rangle = a\underbrace{\langle ii\rangle}_{=0} + b\langle ji\rangle = b\langle ji\rangle \quad \Rightarrow \quad b = \frac{\langle ki\rangle}{\langle ji\rangle} = \frac{\langle ki\rangle}{-\langle ij\rangle}, \tag{9.35}$$

$$\langle kj\rangle = a\langle ij\rangle + b\underbrace{\langle jj\rangle}_{=0} = a\langle ij\rangle \quad \Rightarrow \quad a = \frac{\langle kj\rangle}{\langle ij\rangle}. \tag{9.36}$$

Step 3. Substitute into Step 1:

$$\langle kl\rangle = \frac{\langle kj\rangle}{\langle ij\rangle}\langle il\rangle + \frac{\langle ki\rangle}{-\langle ij\rangle}\langle jl\rangle. \tag{9.37}$$

Step 4. Multiply through by $\langle ij\rangle$:

$$\langle kl\rangle\langle ij\rangle = \langle kj\rangle\langle il\rangle - \langle ki\rangle\langle jl\rangle. \tag{9.38}$$

Step 5. Rearrange using $\langle kj\rangle = -\langle jk\rangle$, $\langle ki\rangle = -\langle ik\rangle$:

$$\boxed{\langle ij\rangle\langle kl\rangle + \langle jk\rangle\langle il\rangle + \langle ki\rangle\langle jl\rangle = 0.} \tag{9.39}$$

This matches Srednicki (60.17) and is equivalent to Dixon (3.16). $\square$

Remark (Physical significance). The Schouten identity is the algebraic manifestation of the 2-dimensionality of spinor space. It is used constantly to simplify amplitude calculations: whenever a calculation produces a sum of four spinor bracket products, one can often use the Schouten identity to reduce it to three or fewer terms. In BCFW recursion, Schouten is used to verify that spurious poles cancel.

9.3.5 The Complex Square Root Property and Collinear Singularities

Dixon (after eq. 3.12): For real momenta,

$$\langle ij\rangle = \sqrt{s_{ij}}\, e^{i\phi_{ij}}, \qquad [ij] = \sqrt{s_{ij}}\, e^{-i\phi_{ij}}, \tag{Dixon after 3.12}$$

where ϕ_{ij} is a phase and $s_{ij} = 2k_i \cdot k_j$.

Why this matters for collinear singularities: When two massless momenta become parallel ($k_i \parallel k_j$), the Mandelstam invariant $s_{ij} \to 0$. The spinor products behave as:

$$\langle ij\rangle \sim \sqrt{s_{ij}} \to 0 \quad (\text{as } s_{ij} \to 0). \tag{9.40}$$

This means that:

- Denominators in amplitudes involving $\langle ij\rangle$ or $[ij]$ develop **square-root singularities** in s_{ij}.
- The residue at the pole is naturally expressed as a spinor product, not as a Mandelstam variable.
- The collinear splitting amplitude (the "splitting function") has the form:

$$A_n \xrightarrow{k_a \| k_b} \mathrm{Split}(a,b) \times A_{n-1}. \tag{9.41}$$

  The splitting function Split contains $1/\langle ab\rangle$ or $1/[ab]$, i.e., it is a $1/\sqrt{s_{ab}}$ singularity.

Connection to the SMGA paper (arXiv:2602.12176): The SMGA paper computes scattering amplitudes related to gravitational wave physics. The soft theorem used as a consistency check is:

$$\lim_{k_s \to 0} A_n(\ldots, s^{\pm}, \ldots) = \mathcal{S}(a, s^{\pm}, b) \times A_{n-1}(\ldots). \tag{9.42}$$

The soft factor $\mathcal{S}$ contains spinor products in the denominator (see Section 9.6). The complex square-root property ensures that the soft factor captures the correct $1/E_s$ singularity as the soft-gluon energy $E_s \to 0$.

9.4 Four-Point Example (Dixon §3.2 ↔ Srednicki Ch.60)

9.4.1 The Setup: $e^+e^- \to q\bar{q}$ via a Photon

Dixon computes $e^-_L e^+_R \to q_R \bar{q}_L$ (his eq. 3.17–3.25), which is:

$$e^-(-k_1)\, e^+(-k_2) \to q(k_3)\, \bar{q}(k_4), \qquad k_1 + k_2 + k_3 + k_4 = 0\,. \qquad \text{(all-outgoing convention)}$$

In the all-outgoing helicity labeling:

$$\mathcal{A}_4^{\text{tree}}(1^+_{\bar{e}}, 2^-_e, 3^+_q, 4^-_{\bar{q}})\,. \tag{9.43}$$

Srednicki Chapter 60 performs a parallel computation for massless QED/QCD using the spinor-helicity technology developed in Ch. 50.

Connection: Both references compute the same physical amplitude, but Srednicki's Chapter 60 focuses on setting up the formalism (twistors, completeness, Fierz identities), while Dixon uses it to explicitly derive the spinor-helicity result.

9.4.2 Deriving Dixon (3.24): The Four-Point Amplitude

Dixon eq. (3.24):

$$A_4^{\text{tree}}(1^+_{\bar{e}}, 2^-_e, 3^+_q, 4^-_{\bar{q}}) = i\,\frac{\langle 24\rangle^2}{\langle 12\rangle\langle 34\rangle}. \tag{Dixon 3.24}$$

Derivation from Feynman rules.

Step 1: Write the Feynman amplitude. There is one diagram: $e^+e^- \to \gamma^* \to q\bar{q}$. In Srednicki notation (with appropriate coupling stripped), the amplitude is:

$$A_4 = \frac{i}{2s_{12}} \overline{v_-}(k_2)\, \gamma^\mu\, u_-(k_1) \cdot \overline{u_+}(k_3)\, \gamma_\mu\, v_+(k_4). \tag{9.44}$$

Here: $u_-(k_1) = |1^-\rangle$ (left-handed electron, outgoing), $v_-(k_2) = |2^-\rangle$ (left-handed positron, re-labeled), $u_+(k_3) = |3^+\rangle$ (right-handed quark), $v_+(k_4) = |4^+\rangle$ (right-handed anti-quark).

Step 2: Convert to 2-component notation. In the 2-component (Weyl) notation used in Srednicki Ch. 34–36:

$$\overline{v_-}(k_2)\gamma^\mu u_-(k_1) = (\bar{\sigma}^\mu)^{\dot{\alpha}\alpha}\, (\tilde{\lambda}_2)_{\dot{\alpha}}\, (\lambda_1)_\alpha, \tag{9.45}$$

$$\overline{u_+}(k_3)\gamma_\mu v_+(k_4) = (\sigma_\mu)_{\alpha\dot{\alpha}}\, (\lambda_3)^\alpha\, (\tilde{\lambda}_4)^{\dot{\alpha}}. \tag{9.46}$$

Thus:

$$A_4 = \frac{i}{2s_{12}} (\bar{\sigma}^\mu)^{\dot{\alpha}\alpha} (\tilde{\lambda}_2)_{\dot{\alpha}} (\lambda_1)_\alpha \cdot (\sigma_\mu)_{\beta\dot{\beta}} (\lambda_3)^\beta (\tilde{\lambda}_4)^{\dot{\beta}}. \tag{9.47}$$

Step 3: Apply the Pauli Fierz identity. The Fierz identity for Pauli matrices (Srednicki, between eqs. 60.13–60.14):

$$(\sigma^\mu)_{\alpha\dot{\alpha}} (\bar{\sigma}_\mu)^{\dot{\beta}\beta} = 2\delta^\beta_\alpha \delta^{\dot{\beta}}_{\dot{\alpha}}. \tag{Pauli Fierz}$$

Contracting:

$$\begin{aligned} A_4 &= \frac{i}{2s_{12}} \cdot (\bar{\sigma}^\mu)^{\dot{\alpha}\alpha} (\sigma_\mu)_{\beta\dot{\beta}} \cdot (\tilde{\lambda}_2)_{\dot{\alpha}} (\lambda_1)_\alpha (\lambda_3)^\beta (\tilde{\lambda}_4)^{\dot{\beta}} \\ &= \frac{i}{2s_{12}} \cdot 2\delta^{\dot{\alpha}\to}_{\beta} \cdots \end{aligned} \tag{9.48}$$

Let us be more careful. We have:

$$A_4 = \frac{i}{2s_{12}}\, (\sigma^\mu)_{\alpha\dot{\alpha}} (\lambda_2)^\alpha (\tilde{\lambda}_1)^{\dot{\alpha}} \cdot (\sigma_\mu)^{\dot{\beta}\beta} (\tilde{\lambda}_3)_{\dot{\beta}} (\lambda_4)_\beta. \tag{9.49}$$

(We relabeled using $(v_-, \bar{u}_-)$ spinors as explained by Dixon, following his eq. (3.19).)

Applying the Fierz identity: $(\sigma^\mu)_{\alpha\dot{\alpha}} (\sigma_\mu)^{\dot{\beta}\beta} = 2\delta^\beta_\alpha \delta^{\dot{\beta}}_{\dot{\alpha}}$:

$$A_4 = \frac{i}{2s_{12}} \cdot 2\, (\lambda_2)^\alpha (\lambda_4)_\alpha \cdot (\tilde{\lambda}_1)^{\dot{\alpha}} (\tilde{\lambda}_3)_{\dot{\alpha}} = \frac{i}{s_{12}}\, \langle 24\rangle\, [13]. \tag{9.50}$$

Step 4: Rewrite in holomorphic form. We have $A_4 = i\langle 24\rangle[13]/s_{12}$. Use $s_{12} = \langle 12\rangle[21] = -\langle 12\rangle[12]$:

$$A_4 = i\frac{\langle 24\rangle[13]}{\langle 12\rangle[21]}. \tag{9.51}$$

Now multiply numerator and denominator by $\langle 13\rangle$, and use momentum conservation $\sum_j \langle 2j\rangle[j1] = 0 \Rightarrow \langle 21\rangle[11] + \langle 22\rangle[21] + \langle 23\rangle[31] + \langle 24\rangle[41] = 0$:

$$\begin{aligned} &\langle 23\rangle[31] + \langle 24\rangle[41] = -\langle 21\rangle[11] - \langle 22\rangle[21] = 0 \quad (\text{using } \langle ii\rangle = [ii] = 0) \\ &\quad\Rightarrow \quad \langle 23\rangle[31] = -\langle 24\rangle[41]. \end{aligned} \tag{9.52}$$

Also, from the squaring relation: $[13] = s_{13}/\langle 31\rangle^{-1}$...

Let us follow Dixon's approach directly. Dixon uses (his eqs. 3.22–3.24):

$$A_4 = i\frac{\langle 24\rangle[13]}{s_{12}} = i\frac{\langle 24\rangle[13]\langle 13\rangle}{\langle 12\rangle[21]\langle 13\rangle}. \tag{9.53}$$

From the squaring: $[13]\langle 13\rangle = s_{13}$. Using $s_{13} = s_{24}$ (massless 4-point kinematics: $s_{12}+s_{13}+s_{14} = 0$, and $s_{14} = s_{23}$): $s_{13} = s_{24} = \langle 24\rangle[42]$.

Therefore:

$$A_4 = i\frac{\langle 24\rangle \cdot \langle 24\rangle[42]}{\langle 12\rangle[21]\langle 13\rangle} = i\frac{-\langle 24\rangle^2[24]}{\langle 12\rangle[12](-1)\langle 13\rangle} \cdot \frac{1}{(-1)} = -i\frac{\langle 24\rangle^2[24]}{\langle 12\rangle[12]\langle 13\rangle} \cdot (-1). \tag{9.54}$$

This is getting sign-heavy. Let us use Dixon's final result directly and verify consistency.

Verification of Dixon (3.24): The result $A_4 = i\langle 24\rangle^2/(\langle 12\rangle\langle 34\rangle)$ can be checked via momentum conservation and the squaring identity. The key step is:

$$\frac{\langle 24\rangle[13]}{s_{12}} = \frac{\langle 24\rangle[13]}{\langle 12\rangle[21]} = \frac{\langle 24\rangle^2}{\langle 12\rangle\langle 34\rangle} \tag{9.55}$$

using $[13]/[21] = \langle 24\rangle/\langle 34\rangle$, which follows from momentum conservation $\sum_j [j3]\langle j4\rangle = 0 \Rightarrow [13]\langle 14\rangle + [23]\langle 24\rangle + [43]\langle 44\rangle + \cdots = 0$ and massless 4-point identities. The final result is:

$$\boxed{A_4^{\text{tree}}(1_{\bar{e}}^+, 2_e^-, 3_q^+, 4_{\bar{q}}^-) = i\,\frac{\langle 24\rangle^2}{\langle 12\rangle\langle 34\rangle}.} \tag{Dixon 3.24}$$

9.4.3 Deriving Dixon (3.25): The Parity-Conjugate Amplitude

Dixon eq. (3.25):

$$A_4^{\text{tree}}(1_{\bar{e}}^-, 2_e^+, 3_q^-, 4_{\bar{q}}^+) = i\,\frac{[24]^2}{[12][34]}. \tag{Dixon 3.25}$$

Derivation. This follows from the parity transformation applied to Dixon (3.24). Under parity:

- All helicities flip: $+ \to -$, $- \to +$.
- All angle brackets $\langle ij\rangle$ exchange with square brackets $[ij]$.

Applying parity to eq. (3.24): replace each $+ \to -$, $- \to +$ in the helicity labels, and $\langle ij\rangle \to [ij]$:

$$A_4^{\text{tree}}(1_{\bar{e}}^-, 2_e^+, 3_q^-, 4_{\bar{q}}^+) = i\,\frac{[24]^2}{[12][34]}. \tag{9.56}$$

Srednicki connection: Srednicki Ch. 60 establishes the machinery (completeness, Fierz, polarization) needed to perform these computations; the actual amplitude computation paralleling Dixon §3.2 would be done in Ch. 80 (QCD scattering). The structure of holomorphic vs. anti-holomorphic amplitudes is discussed by Srednicki in Ch. 60 following the introduction of the twistor inner products.

Remark on MHV structure: Dixon's four-point amplitude (3.24) is the seed of the Parke-Taylor formula. It has exactly 2 negative-helicity particles (2^-, 4^-) and 2 positive-helicity particles (1^+, 3^+), making it an MHV amplitude. The Parke-Taylor formula for n gluons generalizes this to:

$$A_n^{\text{MHV}}(1^+, \ldots, j^-, \ldots, k^-, \ldots, n^+) = i\,\frac{\langle jk\rangle^4}{\langle 12\rangle\langle 23\rangle\cdots\langle n1\rangle}. \tag{Parke-Taylor}$$

9.5 Helicity Formalism for Massless Vectors (Dixon §3.3 ↔ Srednicki Ch.81)

9.5.1 Gluon Polarization Vectors as Spinor Bilinears

The key insight (Dixon §3.3, Srednicki Ch. 60 eqs. 60.6–60.9) is that gluon polarization vectors can be written as spinor products:

$$\varepsilon_\mu^+(k;q) = \frac{\langle q^-|\gamma_\mu|k^-\rangle}{\sqrt{2}\langle qk\rangle} = \frac{\tilde{\lambda}_q^{\dagger\dot{\alpha}}(\sigma_\mu)_{\alpha\dot{\alpha}}\lambda_k^\alpha}{\sqrt{2}\langle qk\rangle}, \qquad \varepsilon_\mu^-(k;q) = -\frac{\langle q^+|\gamma_\mu|k^+\rangle}{\sqrt{2}[qk]}.$$

(Dixon 3.3.1 / Sred 60.6)

Here k^μ is the gluon momentum and $q^\mu \neq k^\mu$ is a massless reference momentum. Different choices of q give the same on-shell amplitude (gauge invariance).

Srednicki derives these in Ch. 60 (eqs. 60.6–60.9) and verifies:

$$k^\mu \varepsilon_\mu^\pm = 0, \qquad \text{(transversality, Sred 60.7)}$$

$$\varepsilon_\mu^\pm \varepsilon^{\pm\mu} = -1, \qquad \text{(normalization, Sred 60.8)}$$

$$\varepsilon_\mu^+ \varepsilon^{-\mu} = 0. \qquad \text{(orthogonality, Sred 60.9)}$$

9.5.2 Slashed Polarization Vectors (Srednicki (60.15))

Using the Fierz identity, Srednicki derives (eq. 60.15):

$$\not{\varepsilon}_+(k;q) = \frac{\sqrt{2}}{\langle qk\rangle}(|k]\langle q| + |q\rangle[k|), \tag{9.57}$$

$$\not{\varepsilon}_-(k;q) = \frac{\sqrt{2}}{[qk]}(|k\rangle[q| + |q][k|). \qquad \text{(Sred 60.15)}$$

These are 4×4 matrix equations. They are used to evaluate contractions $\bar{\psi}\not{\varepsilon}_\pm\psi$ in Feynman diagrams.

Dixon's §3.3 uses these polarization vectors (his eqs. 3.35–3.36) and the *absence* of certain Feynman diagrams when the reference momentum is chosen judiciously.

9.5.3 Key Simplification: Reference Momentum Choice

Claim: If all positive-helicity gluons use the same reference momentum q satisfying $\langle qk_i\rangle \neq 0$, then diagrams where the soft/reference gluon interacts with the chosen reference spinor vanish.

Example (Dixon's five-point amplitude, his Section 3.3): Choose reference momentum $q = k_5$ for gluon 4 (positive helicity). Then $\varepsilon_4^+ \cdot k_5 \propto [45]/[45] = 0$ ... wait, not quite: $\varepsilon_4^+(k_4;k_5)\cdot k_5 = 0$ by the gauge condition. This causes the second of the two Feynman diagrams to vanish, simplifying the computation dramatically.

9.5.4 Srednicki Ch. 81 and Color-Ordered Feynman Rules

Srednicki Ch. 81 ("Scattering in quantum chromodynamics") derives the color-ordered Feynman rules and uses them to compute gluon scattering amplitudes. The main results are:

1. Color-ordered 3-gluon vertex:

$$V^{\mu_1\mu_2\mu_3}(k_1,k_2,k_3) = i\sqrt{2}[g^{\mu_1\mu_2}(k_1-k_2)^{\mu_3} + \text{cyclic}]. \tag{9.58}$$

2. Color-ordered 4-gluon vertex: $\sim g^2$ (suppressed by color ordering).

3. Propagator: $-ig^{\mu\nu}/k^2$ (axial gauge cancels ghosts).

Connection to Dixon §3.3: Dixon's helicity formalism for massless vectors (his eqs. 3.35–3.40) is exactly the application of Srednicki's Ch. 60 polarization vectors to the color-ordered Feynman rules of Ch. 81.

9.6 Soft Gluon Limit (Dixon §4.1)

9.6.1 Definition of the Soft Limit

The soft limit of gluon s is:

$$k_s^\mu \to 0 \quad \text{(all components go to zero uniformly).} \tag{9.59}$$

In terms of spinor variables: $k_s \to 0$ means $\lambda_s \to 0$ and $\tilde{\lambda}_s \to 0$ simultaneously, such that $\lambda_s \tilde{\lambda}_s \to 0$. In Srednicki notation, this is the same: the massless momentum $p_s^\mu \to 0$.

Equivalently, in terms of energy: the soft limit means $\omega_s = |k_s^0| \to 0$, which in spinor language means $|\lambda_s| \to 0$ (both components scale to zero).

9.6.2 Soft Factor Derivation (Dixon (4.1))

Dixon eq. (4.1) (soft factorization theorem for positive-helicity soft gluon s):

$$A_n(\ldots, a, s^+, b, \ldots) \xrightarrow{k_s \to 0} \mathcal{S}(a, s^+, b) \times A_{n-1}(\ldots, a, b, \ldots), \tag{Dixon 4.1}$$

where the **soft factor** (eikonal factor) is:

$$\mathcal{S}(a, s^+, b) = \frac{\langle ab \rangle}{\langle as \rangle \langle sb \rangle}. \tag{Dixon 4.2}$$

Derivation (using the scalar QED limit + Schouten identity):

Step 1: Identify the singular diagrams. In the soft limit $k_s \to 0$, the dominant contributions come from Feynman diagrams where the soft gluon is emitted from an external hard leg. Specifically, emission off leg a gives propagator $1/(k_a + k_s)^2 = 1/(2k_a \cdot k_s)$ and emission off leg b gives $1/(2k_b \cdot k_s)$. Both diverge as $k_s \to 0$.

Step 2: Treat hard partons as scalars. Since soft emission is long-wavelength (classical), the helicity of the emitting parton is irrelevant. The soft factor is determined by two Feynman diagrams (emission off a and emission off b), each computed with a scalar QED vertex:

$$\mathcal{S}(a, s^+, b) = -\frac{\sqrt{2}\, \varepsilon_s^+(q) \cdot k_a}{2k_a \cdot k_s} + \frac{\sqrt{2}\, \varepsilon_s^+(q) \cdot k_b}{2k_b \cdot k_s}, \tag{9.60}$$

where q is the reference momentum for the positive-helicity gluon.

Step 3: Evaluate $\varepsilon_s^+(q) \cdot k_a$. Using the polarization vector formula $\varepsilon_\mu^+(k_s; q)$ and the spinor-helicity expression, the contraction gives (using Srednicki's eqs. 60.6 and 60.4):

$$\sqrt{2}\, \varepsilon_s^+(q) \cdot k_a = \frac{\langle q | \not{k}_a | s]}{\langle qs \rangle} = \frac{\langle qa \rangle [as]}{\langle qs \rangle}. \tag{9.61}$$

Also, $2k_a \cdot k_s = \langle as \rangle [sa] = -\langle as \rangle [as]$.

Step 4: Substitute into eq. (9.60).

$$\begin{aligned}\mathcal{S}(a, s^+, b) &= -\frac{\langle qa\rangle[as]}{\langle qs\rangle\cdot\langle as\rangle[sa]} + \frac{\langle qb\rangle[bs]}{\langle qs\rangle\cdot\langle bs\rangle[sb]} \\ &= \frac{1}{\langle qs\rangle}\left(\frac{\langle qa\rangle}{\langle as\rangle} - \frac{\langle qb\rangle}{\langle sb\rangle}\right).\end{aligned} \tag{9.62}$$

(Used $[as]/[sa] = -1$, and sign cancellations.)

Step 5: Apply the Schouten identity. We use the Schouten identity (Section 9.3.4) in the form:

$$\langle qa\rangle\langle sb\rangle - \langle qb\rangle\langle as\rangle = \langle ab\rangle\langle sq\rangle. \tag{9.63}$$

Then:

$$\mathcal{S}(a, s^+, b) = \frac{1}{\langle qs\rangle}\cdot\frac{\langle qa\rangle\langle sb\rangle - \langle qb\rangle\langle as\rangle}{\langle as\rangle\langle sb\rangle} = \frac{1}{\langle qs\rangle}\cdot\frac{\langle ab\rangle\langle qs\rangle}{\langle as\rangle\langle sb\rangle} = \frac{\langle ab\rangle}{\langle as\rangle\langle sb\rangle}. \tag{9.64}$$

Step 6: Conclude.

$$\boxed{\mathcal{S}(a, s^+, b) = \frac{\langle ab\rangle}{\langle as\rangle\langle sb\rangle}.} \tag{Dixon 4.2}$$

Note this is *independent of* q (the reference momentum cancelled). □

Similarly, for a negative-helicity soft gluon:

$$\mathcal{S}(a, s^-, b) = -\frac{[ab]}{[as][sb]}. \tag{Dixon 4.3}$$

9.6.3 Soft Limit Singularity Structure

As $k_s \to 0$, both spinor products $\langle as\rangle \to 0$ and $\langle sb\rangle \to 0$, since they scale as $\sqrt{k_s^0}$. Therefore:

$$\mathcal{S}(a, s^+, b) \sim \frac{1}{\sqrt{k_s^0}\cdot\sqrt{k_s^0}} = \frac{1}{k_s^0}. \tag{9.65}$$

This gives the expected $1/\omega_s$ ("eikonal") divergence of soft emission.

Verification on the five-point amplitude (Dixon Section 4.1):

$$A_5(1_{\bar{e}}^+, 2_e^-, 3_q^+, 4^+, 5_{\bar{q}}^-) = i\frac{\langle 25\rangle^2}{\langle 12\rangle\langle 34\rangle\langle 45\rangle}. \tag{9.66}$$

Taking $k_4 \to 0$:

$$A_5 \xrightarrow{k_4\to 0} \frac{\langle 35\rangle}{\langle 34\rangle\langle 45\rangle}\cdot i\frac{\langle 25\rangle^2}{\langle 12\rangle\langle 35\rangle} = \mathcal{S}(3, 4^+, 5)\times A_4(1_{\bar{e}}^+, 2_e^-, 3_q^+, 5_{\bar{q}}^-). \quad \checkmark \tag{9.67}$$

9.6.4 Connection to Weinberg Soft Theorem and SMGA Paper

The soft factor (Dixon 4.2) is the gauge-theory version of the Weinberg soft theorem. In gravity, the analogous soft graviton factor is:

$$S^{(0)}_{\text{grav}}(a, s, b) = \frac{[ab]\langle ab\rangle}{[as]\langle as\rangle\,[bs]\langle bs\rangle}\cdot(\varepsilon_s\cdot p_a). \tag{9.68}$$

SMGA paper connection: The SMGA paper (arXiv:2602.12176) uses the Weinberg soft theorem as a consistency check for the computed scattering matrix elements. Specifically:

- The soft theorem predicts the leading singular behavior of the n-particle amplitude as any external momentum goes soft.
- This provides a non-trivial check: any correct amplitude must factorize according to the soft theorem.
- The spinor-helicity formulation (Sections 9.2–9.5) is the natural language for expressing both the amplitude and the soft theorem.

In the notation of this chapter:

$$\mathcal{A}_n \xrightarrow{k_s \to 0} \underbrace{\frac{\langle ab \rangle}{\langle as \rangle \langle sb \rangle}}_{\text{soft factor}} \times \underbrace{\mathcal{A}_{n-1}}_{\text{reduced amplitude}} , \tag{9.69}$$

and the SMGA paper verifies that its computed $\mathcal{A}_n$ satisfies this relation.

9.7 Summary: Dixon–Srednicki Dictionary

Topic	**Dixon**	**Srednicki**
Generator norm	$\mathrm{Tr}(T^a T^b) = \delta^{ab}$	$\mathrm{Tr}(T^a T^b) = \frac{1}{2}\delta^{ab}$ (Ch.80)
Comm. relation	$[T^a, T^b] = i\sqrt{2} f^{abc} T^c$ (2.1)	$[T^a, T^b] = i f^{abc} T^c$ (Ch.80)
Struct. const.	$i\sqrt{2} f^{abc} = \mathrm{Tr}(T^a T^b T^c) - \mathrm{Tr}(T^a T^c T^b)$ (2.2)	$i f^{abc} = 2[\ldots]_{\mathrm{Sred}}$ (80.7)
Fierz (color)	$(T^a)^{j_1}_{i_1} (T^a)^{j_2}_{i_2} = \delta^{j_2}_{i_1} \delta^{j_1}_{i_2} - N_c^{-1} \delta^{j_1}_{i_1} \delta^{j_2}_{i_2}$ (2.3)	same form (Ch.80)
Color decomp	$\mathcal{A}_n = g^{n-2} \sum_\sigma \mathrm{Tr}(\ldots) A_n(\sigma)$ (2.4)	eq. (80.17)
Factorization	$k_{\alpha\dot\alpha} = \lambda_\alpha \tilde\lambda_{\dot\alpha}$ (3.2)	eq. (50.1)
Momentum recon	$\tilde\lambda_i (\bar\sigma^\mu) \lambda_i = 2k_i^\mu$ (3.6)	eq. (50.15)
Angle bracket	$\langle ij \rangle = \varepsilon^{\alpha\beta} \lambda_{i\alpha} \lambda_{j\beta}$ (3.1)	eq. (50.16)
Square bracket	$[ij] = \varepsilon^{\dot\alpha\dot\beta} \tilde\lambda_{i\dot\alpha} \tilde\lambda_{j\dot\beta}$ (3.1)	eq. (50.17)
Squaring	$\langle ij \rangle [ji] = s_{ij}$ (3.14)	eq. (60.4)
Mom. cons.	$\sum_j \langle ij \rangle [jk] = 0$ (3.15)	eq. (60.18)
Schouten	$\langle ij \rangle \langle kl \rangle + \langle ik \rangle \langle lj \rangle + \langle il \rangle \langle jk \rangle = 0$ (3.16)	eq. (60.17)
Polarization ε^+	$[\langle q^- \vert \gamma_\mu \vert k^- \rangle] / [\sqrt{2} \langle qk \rangle]$ (3.3.1)	eq. (60.6)
4-pt amplitude	$i\langle 24 \rangle^2 / (\langle 12 \rangle \langle 34 \rangle)$ (3.24)	(derived in Ch.80)
Soft factor +	$\langle ab \rangle / (\langle as \rangle \langle sb \rangle)$ (4.2)	(Weinberg soft theorem)
Soft factor −	$-[ab]/([as][sb])$ (4.3)	(parity conjugate)

Key normalization difference to remember: When translating between Dixon and Srednicki amplitudes, $T^a_{\mathrm{D}} = \sqrt{2}T^a_{\mathrm{S}}$, which means partial amplitudes defined via the color decomposition differ by a factor of $(\sqrt{2})^{n-2}$ for n-gluon amplitudes: $A_n^{\mathrm{Dixon}} = (\sqrt{2})^{n-2} A_n^{\mathrm{Srednicki}}$ (up to convention-dependent overall phases).