

1 Types

$$\alpha, \beta ::= \tau \mid \alpha \rightarrow \beta$$

1.1 First Element of a List Satisfying a Property

$$\text{here } \frac{P(x)}{\text{First } P \ (x : xs)} \quad \text{there } \frac{\neg P(x) \quad \text{First } P \ xs}{\text{First } P \ (x : xs)}$$

A context is a list of pairs of atoms and types:

$$\text{Context} \equiv [(a, \alpha)]$$

, where for any atom a and context Γ we denote as $a \in \Gamma$ when:

$$\text{First } ((_ \equiv _ b) \circ \pi_1) \ \Gamma$$

Given evidence that an atom a is a context Γ , of type $a \in \Gamma$. Note this evidence always points to its first occurrence by definition of `First` predicate. We denote the function that returns the associated type for the atom a by simple concatenation:

$$_ _ : (\Gamma : \text{Context}) \rightarrow a \in \Gamma \rightarrow \alpha$$

To make the notation more classic, we will just concatenate the context directly with the atom, not the evidence that the atom is in the context, but in all applications of this function we will ask in the premises the required evidence.

We say a context Γ is included in other Δ , denoted as $\Gamma \subseteq \Delta$, if:

$$\forall a \in \Gamma, a \in \Delta \wedge \Gamma x \equiv \Delta x$$

Extending or overriding a context Γ with a new association atom a with a type α is denoted by $\Gamma, a : \alpha$.

1.2 Typing Relation

$$\begin{array}{l} \vdash_v \frac{x \in \Gamma}{\Gamma \vdash x : \Gamma x} \quad \vdash_a \frac{\Gamma \vdash M : \alpha \rightarrow \beta \quad \Gamma \vdash N : \alpha}{\Gamma \vdash MN : \beta} \\ \vdash_\lambda \frac{\Gamma, x : \alpha \vdash M : \beta}{\Gamma \vdash \lambda x M : \alpha \rightarrow \beta} \end{array}$$

1.3 Weakening Lemmas

Lemma 1 (Weakening Lemma).

$$\Gamma \subseteq \Delta \wedge \Gamma \vdash M : \alpha \Rightarrow \Delta \vdash M : \alpha$$

Proof. Induction on the typing relation.

var case: Hypothesis : $\vdash_v \frac{x \in \Gamma}{\Gamma \vdash x : \Gamma x}$ Thesis : $\Delta \vdash x : \Delta x$

$$\vdash_v \frac{\frac{x \in \Gamma \quad \Gamma \subseteq \Delta}{x \in \Delta \wedge \Gamma x = \Delta x}}{\Delta \vdash x : \Delta x = \Gamma x}$$

app. case: Hypothesis: $\vdash_a \frac{\Gamma \vdash M : \alpha \rightarrow \beta \quad \Gamma \vdash N : \alpha}{\Gamma \vdash MN : \beta}$ Thesis: $\Delta \vdash MN : \beta$

$$ih \frac{\frac{\Gamma \subseteq \Delta \quad \Gamma \vdash M : \alpha \rightarrow \beta}{\Delta \vdash M : \alpha \rightarrow \beta} \quad ih \frac{\Gamma \subseteq \Delta \quad \Gamma \vdash N : \alpha}{\Delta \vdash N : \beta}}{\vdash_a \Delta \vdash MN : \beta}$$

abs. case: Hypothesis: $\vdash_\lambda \frac{\Gamma, x : \alpha \vdash M : \beta}{\Gamma \vdash \lambda x M : \alpha \rightarrow \beta}$

Thesis: $\Delta \vdash \lambda x M : \alpha \rightarrow \beta$

$$ih \frac{\frac{\Gamma \subseteq \Delta}{\Gamma, x : \beta \subseteq \Delta, x : \beta} \quad \Gamma, x : \beta \vdash M : \alpha}{\vdash_\lambda \frac{\Delta, x : \beta \vdash M : \alpha}{\Delta \vdash \lambda x M : \alpha}}$$

□

Lemma 2 (Weakening Fresh Atoms).

$$x \# M \wedge \Gamma \vdash M : \alpha \Rightarrow \Gamma, x : \beta \vdash M : \alpha$$

Proof. Induction on the typing relation.

var case: Hypothesis : $\vdash_v \frac{y \in \Gamma}{\Gamma \vdash y : \Gamma y}$ Thesis : $\Gamma, x : \beta \vdash y : \Gamma y$

$$\vdash_v \frac{\frac{\frac{x \# y}{x \neq y} \quad \frac{\Gamma \vdash y : \Gamma y}{y \in \Gamma}}{y \in (\Gamma, x : \beta) \wedge (\Gamma, x : \beta)y = \Gamma y}}{\Gamma, x : \beta \vdash y : (\Gamma, x : \beta)y = \Gamma y}$$

app. case: Hypothesis: $\vdash_a \frac{\Gamma \vdash M : \gamma \rightarrow \alpha \quad \Gamma \vdash N : \gamma}{\Gamma \vdash MN : \alpha}$

Thesis: $\Gamma, x : \beta \vdash MN : \alpha$

$$ih \frac{\frac{\frac{x \# MN}{x \# M} \quad \Gamma \vdash M : \gamma \rightarrow \alpha}{\Gamma, x : \beta \vdash M : \gamma \rightarrow \alpha} \quad ih \frac{\frac{x \# MN}{x \# N} \quad \Gamma \vdash N : \gamma}{\Gamma, x : \beta \vdash N : \gamma}}{\Gamma, x : \beta \vdash MN : \alpha}$$

abs. case: Hypothesis: $\vdash_\lambda \frac{\Gamma, y : \gamma \vdash M : \alpha}{\Gamma \vdash \lambda y M : \gamma \rightarrow \alpha}$

Thesis: $\Gamma, x : \beta \vdash \lambda y M : \gamma \rightarrow \alpha$

$$\begin{array}{l}
x = y \text{ subcase: } \text{lem. 1} \frac{\Gamma, x : \gamma \subseteq \Gamma, x : \beta, x : \gamma \quad \Gamma, x : \gamma \vdash M : \alpha}{\vdash \lambda \frac{\Gamma, x : \beta, x : \gamma \vdash M : \alpha}{\Gamma, x : \beta \vdash \lambda x M : \gamma \rightarrow \alpha}} \\
\\
x \neq y \text{ subcase: } \text{ih} \frac{\frac{x \# \lambda y M \quad x \neq y}{x \# M} \quad \Gamma, y : \gamma \vdash M : \alpha}{\Gamma, y : \gamma, x : \beta \vdash M : \alpha} \quad x \neq y \\
\vdash \lambda \frac{\Gamma, x : \beta, y : \gamma \vdash M : \alpha}{\Gamma, x : \beta \vdash \lambda y M : \gamma \rightarrow \alpha}
\end{array}$$

□

1.4 Strengthening Lemma

Lemma 3 (Strengthening Fresh Variables).

$$x \# M \wedge \Gamma, x : \beta \vdash M : \alpha \Rightarrow \Gamma \vdash M : \alpha$$

Proof. Induction on the typing relation.

$$\text{var case: Hypothesis : } \vdash_v \frac{y \in \Gamma, x : \beta}{\Gamma, x : \beta \vdash y : (\Gamma, y : \beta)y} \quad \text{Thesis : } \Gamma \vdash y : (\Gamma, x : \beta)y$$

$$\vdash_v \frac{\frac{x \# y}{x \neq y} \quad y \in \Gamma, x : \beta}{y \in \Gamma \wedge (\Gamma, x : \beta)y = \Gamma y} \\
\Gamma \vdash y : \Gamma y = (\Gamma, x : \beta)y$$

$$\text{app. case: Hypothesis: } \vdash_a \frac{\Gamma, x : \beta \vdash M : \gamma \rightarrow \alpha \quad \Gamma, x : \beta \vdash N : \gamma}{\Gamma, x : \beta \vdash MN : \alpha}$$

Thesis: $\Gamma \vdash MN : \alpha$

$$\text{ih} \frac{\frac{x \# MN}{x \# M} \quad \Gamma, x : \beta \vdash M : \gamma \rightarrow \alpha}{\Gamma \vdash M : \gamma \rightarrow \alpha} \quad \text{ih} \frac{\frac{x \# MN}{x \# N} \quad \Gamma, x : \beta \vdash N : \gamma}{\Gamma \vdash N : \gamma} \\
\Gamma \vdash MN : \alpha$$

$$\text{abs. case: Hypothesis: } \vdash_\lambda \frac{\Gamma, x : \beta, y : \gamma \vdash M : \alpha}{\Gamma, x : \beta \vdash \lambda y M : \gamma \rightarrow \alpha}$$

Thesis: $\Gamma \vdash \lambda y M : \gamma \rightarrow \alpha$

$$x = y \text{ subcase: } \text{lem. 1} \frac{\Gamma, x : \gamma \subseteq \Gamma, x : \beta, x : \gamma \quad \Gamma, x : \gamma \vdash M : \alpha}{\vdash \lambda \frac{\Gamma, x : \beta, x : \gamma \vdash M : \alpha}{\Gamma, x : \beta \vdash \lambda x M : \gamma \rightarrow \alpha}}$$

$$x \neq y \text{ subcase: } \text{ih} \frac{\frac{x \# \lambda y M \quad x \neq y}{x \# M} \quad \Gamma, y : \gamma \vdash M : \alpha}{\Gamma, y : \gamma, x : \beta \vdash M : \alpha} \quad x \neq y \\
\vdash \lambda \frac{\Gamma, x : \beta, y : \gamma \vdash M : \alpha}{\Gamma, x : \beta \vdash \lambda y M : \gamma \rightarrow \alpha}$$

□

1.5 Equivariant

Lemma 4 (Typing Relation is Equivariant).

$$\Gamma \vdash M : \alpha \Rightarrow (x \ y)\Gamma \vdash (x \ y)M : \alpha$$

Proof. Induction on the typing relation.

$$\begin{array}{l} \text{var case: Hypothesis : } \vdash_v \frac{z \in \Gamma}{\Gamma \vdash z : \Gamma z} \quad \text{Thesis : } (x \ y)\Gamma \vdash (x \ y)z : \Gamma z \\ \vdash_v \frac{\frac{z \in \Gamma}{(x \ y)z \in (x \ y)\Gamma \wedge \Gamma z = ((x \ y)\Gamma)((x \ y)z)}}{(x \ y)\Gamma \vdash (x \ y)z : ((x \ y)\Gamma)((x \ y)z) = \Gamma z} \end{array}$$

$$\begin{array}{l} \text{app. case: Hypothesis: } \vdash_a \frac{\Gamma \vdash M : \beta \rightarrow \alpha \quad \Gamma \vdash N : \beta}{\Gamma \vdash MN : \alpha} \\ \text{Thesis: } (x \ y)\Gamma \vdash (x \ y)(MN) : \alpha \\ \text{ih } \frac{\Gamma \vdash M : \beta \rightarrow \alpha}{(x \ y)\Gamma \vdash (x \ y)M : \beta \rightarrow \alpha} \quad \text{ih } \frac{\Gamma \vdash N : \gamma}{(x \ y)\Gamma \vdash (x \ y)N : \gamma} \\ \frac{(x \ y)\Gamma \vdash ((x \ y)M)((x \ y)N) : \alpha}{(x \ y)\Gamma \vdash (x \ y)(MN) : \alpha} \end{array}$$

$$\begin{array}{l} \text{abs. case: Hypothesis: } \vdash_\lambda \frac{\Gamma, z : \beta \vdash M : \alpha}{\Gamma \vdash \lambda z M : \beta \rightarrow \alpha} \\ \text{Thesis: } (x \ y)\Gamma \vdash (x \ y)(\lambda z M) : \beta \rightarrow \alpha \\ \text{ih } \frac{\Gamma, z : \beta \vdash M : \alpha}{(x \ y)(\Gamma, z : \beta) \vdash (x \ y)M : \alpha} \\ \vdash_\lambda \frac{\frac{((x \ y)\Gamma), (x \ y)z : \beta \vdash (x \ y)M : \alpha}{(x \ y)\Gamma \vdash \lambda((x \ y)z)((x \ y)M) : \beta \rightarrow \alpha}}{(x \ y)\Gamma \vdash (x \ y)(\lambda z M) : \beta \rightarrow \alpha} \end{array}$$

□

1.6 α -compatible

Lemma 5 (Typing relation is α -compatible).

$$\Gamma \vdash M : \alpha \wedge M \sim_\alpha N \Rightarrow \Gamma \vdash N : \alpha$$

Proof. Permutation induction on M .

var case: Direct.

app. case: Direct.

abs. case: Hypotheses: $\vdash \lambda \frac{\Gamma, x : \alpha \vdash M : \beta}{\Gamma \vdash \lambda x M : \alpha \rightarrow \beta} \sim_{\alpha} \lambda \frac{\exists xs, \forall z \notin xs, (x z)M \sim_{\alpha} (y z)N}{\lambda x M \sim_{\alpha} \lambda y N}$
 Inductive Hypothesis:

$$\forall u, v, N, \Gamma, \gamma, \left. \begin{array}{l} \Gamma \vdash (u v)M : \gamma \\ (u v)M \sim_{\alpha} (u v)N \end{array} \right\} \wedge \Rightarrow \Gamma \vdash (u v)N : \gamma$$

Thesis: $\Gamma \vdash \lambda y N : \beta \rightarrow \alpha$

Let be $z \notin \{y\} \cup xs \cup \text{dom}(\Gamma) \cup \text{fv}(N)$.

$$\begin{array}{c} \text{lem. 4} \frac{\Gamma, x : \alpha \vdash M : \beta}{(x z)(\Gamma, x : \alpha) \vdash (x z)M : \beta} \quad \frac{z \notin \text{dom}(\Gamma)}{(x z)\Gamma, z : \alpha \subseteq \Gamma, z : \alpha} \quad \frac{z \notin xs}{(x z)M \sim_{\alpha} (y z)N} \\ \text{lem. 1} \frac{(x z)\Gamma, z : \alpha \vdash (x z)M : \beta}{\text{ih} \frac{\Gamma, z : \alpha \vdash (x z)M : \beta}{\Gamma, z : \alpha \vdash (y z)N : \beta}} \quad \frac{(y z)(\Gamma, z : \alpha) \vdash (y z)(y z)N : \beta}{(y z)\Gamma, y : \alpha \vdash N : \beta} \\ \text{lem. 2} \frac{z \# N \quad (y z)\Gamma, y : \alpha \vdash N : \beta}{(y z)\Gamma, y : \alpha, z : \alpha \vdash N : \beta} \\ \text{lem. 3} \frac{z \# N \quad \Gamma, y : \alpha, z : \alpha \vdash N : \beta}{\vdash \lambda \frac{\Gamma, y : \alpha \vdash N : \beta}{\Gamma \vdash \lambda y N : \alpha \rightarrow \beta}} \end{array}$$

□

1.7 Substitution Lemma for Typing Relation

Lemma 6 (Substitution Lemma for Typing Relation).

$$\Gamma, x : \beta \vdash M : \alpha \wedge \Gamma \vdash N : \beta \Rightarrow \Gamma \vdash M[x := N] : \alpha$$

Proof. α -induction principle on M term to avoid x and free variables in N as bound variable in the abstraction case. The predicate is α -compatible on M as direct consequence of the substitution lemma for α -convertible terms.

$$\text{var case: Hypothesis : } \vdash_v \frac{y \in (\Gamma, x : \beta)}{\Gamma, x : \beta \vdash y : (\Gamma, x : \beta)y}$$

$$\text{Thesis : } \Gamma \vdash y[x := N] : (\Gamma, x : \beta)y$$

$$\text{case } x = y: \text{ hyp. } \frac{\Gamma \vdash x[x := N] : (\Gamma, x : \beta)x}{\Gamma \vdash N : \beta}$$

$$\text{case } x \neq y: \vdash_v \frac{y \in (\Gamma, x : \beta) \quad x \neq y}{\Gamma \vdash \underbrace{y}_{=y[x:=N]} : \underbrace{\Gamma y}_{=(\Gamma, x : \beta)y}}$$

app. case: Hypothesis: $\vdash_a \frac{\Gamma, x : \beta \vdash M : \gamma \rightarrow \alpha \quad \Gamma, x : \beta \vdash P : \gamma}{\Gamma, x : \beta \vdash MP : \alpha}$

Thesis: $\Gamma \vdash (MP)[x := N] : \alpha$

ih. $\frac{\Gamma, x : \beta \vdash M : \gamma \rightarrow \alpha}{\Gamma \vdash M[x := N] : \gamma \rightarrow \alpha} \quad \text{ih. } \frac{\Gamma, x : \beta \vdash P : \gamma}{\Gamma \vdash P[x := N] : \gamma}$
 $\vdash_a \frac{\Gamma \vdash M[x := N] : \gamma \rightarrow \alpha \quad \Gamma \vdash P[x := N] : \gamma}{\Gamma \vdash \underbrace{(M[x := N])(P[x := N])}_{=(MP)[x:=N]} : \alpha}$

abs. case: Hypothesis: $\vdash \lambda \frac{\Gamma, x : \beta, y : \gamma \vdash M : \alpha}{\Gamma, x : \beta \vdash \lambda y M : \gamma \rightarrow \alpha}$

Thesis: $\Gamma \vdash (\lambda y M)[x := N] : \gamma \rightarrow \alpha$

We can assume $y \notin \{x\} \cup fv(N)$.

lem. 1 $\frac{\Gamma, x : \beta, y : \gamma \vdash M : \alpha \quad \frac{y \neq x}{\Gamma, x : \beta, y : \gamma \subseteq \Gamma, y : \gamma, x : \beta}}{\Gamma, y : \gamma, x : \beta \vdash M : \alpha} \quad \text{ih. } \frac{\Gamma, y : \gamma \vdash M[x := N] : \alpha}{\Gamma \vdash \lambda y (M[x := N]) : \gamma \rightarrow \alpha}$
 lem. 2 $\frac{\Gamma \vdash N : \beta \quad y \# N}{\Gamma, y : \gamma \vdash N : \beta}$

Finally, by the α -compatible property of typing relation and due to freshness of binder variable we end up with the naive case of substitution case.

lem. 5 $\frac{\Gamma \vdash \lambda y (M[x := N]) : \gamma \rightarrow \alpha \quad \frac{y \notin \{x\} \cup fv(N)}{\lambda y (M[x := N]) \sim_\alpha (\lambda y M)[x := N]}}{\Gamma \vdash (\lambda y M)[x := N] : \gamma \rightarrow \alpha}$

□

2 Subject Reduction

$$\triangleright_\beta \frac{}{(\lambda x M)N \triangleright M[x := N]}$$

Corollary 1 (Typing judge is preserved by β -contraction).

$$\Gamma \vdash M : \alpha, M \triangleright N \Rightarrow \Gamma \vdash N : \alpha$$

Proof. We have to prove that $\Gamma \vdash M[x := N] : \alpha$. By hypotheses $\Gamma \vdash (\lambda x M) : \alpha$, then by typing judge $\Gamma, x : \alpha \vdash M : \beta$ and $\Gamma \vdash N : \alpha$, using last lemma 6 we end up with the desired result. □

We define the β -reduction relation as the contextual clousure of the β -contraction. We denote it as \rightarrow_β .

Corollary 2. $\Gamma \vdash M : \alpha, M \rightarrow_\beta N \Rightarrow \Gamma \vdash N : \alpha$

Proof. Direct induction on the contextual clousure relation. For the β -contraction case we directly use last result. □

Subject reduction is proved by induction on the reflexive-transitive closure of $\rightarrow_\alpha(\rightarrow_\beta \cup \sim_\alpha)$ relation, using directly the last result 2 and the α -compatible of the typing relation lemma 5 for the \rightarrow_α and \sim_α cases respectively.

Lemma 7 (Subject Reduction). $\Gamma \vdash M : \alpha, M \rightarrow_\beta^* N \Rightarrow \Gamma \vdash N : \alpha$