

Algorithms

Growth of functions

Emanuele Rodolà

rodola@di.uniroma1.it



SAPIENZA
UNIVERSITÀ DI ROMA

Θ -notation

Given a function $g(n)$, define the **set of functions**:

$$\Theta(g(n)) = \{f(n) : 0 \leq c_1g(n) \leq f(n) \leq c_2g(n) \text{ for all } n \geq n_0\}$$


where c_1, c_2, n_0 are positive constants.

Θ -notation

Given a function $g(n)$, define the **set of functions**:

$$\Theta(g(n)) = \{f(n) : 0 \leq c_1g(n) \leq f(n) \leq c_2g(n) \text{ for all } n \geq n_0\}$$

where c_1, c_2, n_0 are positive constants.



Θ -notation: Example

Intuition: Ignore the lower-order terms and the coefficient for the higher-order term.

Θ -notation: Example

Intuition: Ignore the **lower-order** terms and the **coefficient** for the higher-order term.

Example:

$$\frac{1}{2}n^2 - 3n \in \Theta(n^2)$$

Θ -notation: Example

Intuition: Ignore the **lower-order** terms and the **coefficient** for the higher-order term.

Example:

$$\frac{1}{2}n^2 - 3n \in \Theta(n^2)$$

If this is true, then we can find positive constants c_1, c_2, n_0 such that:

$$c_1 n^2 \leq \frac{1}{2}n^2 - 3n \leq c_2 n^2 \quad \text{for all } n \geq n_0$$

Θ -notation: Example

Intuition: Ignore the **lower-order** terms and the **coefficient** for the higher-order term.

Example:

$$\frac{1}{2}n^2 - 3n \in \Theta(n^2)$$

If this is true, then we can find positive constants c_1, c_2, n_0 such that:

$$c_1 \leq \frac{1}{2} - \frac{3}{n} \leq c_2 \quad \text{for all } n \geq n_0$$

Θ -notation: Example

Intuition: Ignore the lower-order terms and the coefficient for the higher-order term.

Example:

$$\frac{1}{2}n^2 - 3n \in \Theta(n^2)$$

If this is true, then we can find positive constants c_1, c_2, n_0 such that:

$$c_1 \leq \frac{1}{2} - \frac{3}{n} \leq c_2 \quad \text{for all } n \geq n_0$$

This holds, for instance, with $c_1 = \frac{1}{14}, c_2 = \frac{1}{2}, n_0 = 7$.

Θ -notation: Example

Intuition: Ignore the lower-order terms and the coefficient for the higher-order term.

Example:

$$\frac{1}{2}n^2 - 3n \in \Theta(n^2)$$

If this is true, then we can find positive constants c_1, c_2, n_0 such that:

$$c_1 \leq \frac{1}{2} - \frac{3}{n} \leq c_2 \quad \text{for all } n \geq n_0$$

This holds, for instance, with $c_1 = \frac{1}{14}, c_2 = \frac{1}{2}, n_0 = 7$.

Does it meet the intuition?

- A coefficient that is different from $\frac{1}{2}$ in $\frac{1}{2}n^2$ would just change c_1, c_2 .

Θ -notation: Example

Intuition: Ignore the **lower-order** terms and the **coefficient** for the higher-order term.

Example:

$$\frac{1}{2}n^2 - 3n \in \Theta(n^2)$$

If this is true, then we can find positive constants c_1, c_2, n_0 such that:

$$c_1 \leq \frac{1}{2} - \frac{3}{n} \leq c_2 \quad \text{for all } n \geq n_0$$

This holds, for instance, with $c_1 = \frac{1}{14}, c_2 = \frac{1}{2}, n_0 = 7$.

Does it meet the intuition?

- A coefficient that is different from $\frac{1}{2}$ in $\frac{1}{2}n^2$ would just change c_1, c_2 .
- The mere presence of $\frac{1}{2}$ allows us to find c_1, c_2 , so we can always **dominate** the linear term.

Θ -notation: Example

Intuition: Ignore the **lower-order** terms and the **coefficient** for the higher-order term.

Example:

$$an^2 + bn + c \in \Theta(n^2) \quad \text{with } a > 0$$

Θ -notation: Example

Intuition: Ignore the **lower-order** terms and the **coefficient** for the higher-order term.

Example:

$$an^2 + bn + c \in \Theta(n^2) \quad \text{with } a > 0$$

Or any polynomial with degree d :

$$\sum_{i=0}^d a_i n^i \in \Theta(n^d) \quad \text{with } a_d > 0$$

Θ -notation: Example

Intuition: Ignore the **lower-order** terms and the **coefficient** for the higher-order term.

Example:

$$an^2 + bn + c \in \Theta(n^2) \quad \text{with } a > 0$$

Or any polynomial with degree d :

$$\sum_{i=0}^d a_i n^i \in \Theta(n^d) \quad \text{with } a_d > 0$$

Which includes constant functions:

$$an^0 \in \Theta(n^0) \quad \text{with } a > 0$$

Θ -notation: Example

Intuition: Ignore the lower-order terms and the coefficient for the higher-order term.

Example:

$$an^2 + bn + c \in \Theta(n^2) \quad \text{with } a > 0$$

Or any polynomial with degree d :

$$\sum_{i=0}^d a_i n^i \in \Theta(n^d) \quad \text{with } a_d > 0$$

Which includes constant functions:

$$a \in \Theta(1) \quad \text{with } a > 0$$

O -notation

Given a function $g(n)$, define the **set of functions**:

$$O(g(n)) = \{f(n) : 0 \leq f(n) \leq cg(n) \text{ for all } n \geq n_0\}$$


where c, n_0 are positive constants.

O -notation

Given a function $g(n)$, define the **set of functions**:

$$O(g(n)) = \{f(n) : 0 \leq f(n) \leq cg(n) \text{ for all } n \geq n_0\}$$

where c, n_0 are positive constants.



O -notation: Example

Intuition: Start from the observation that:

$$f(n) \in \Theta(g(n)) \text{ implies } f(n) \in O(g(n))$$

Thus, lower-order terms are also allowed now.

O -notation: Example

Intuition: Start from the observation that:

$$f(n) \in \Theta(g(n)) \text{ implies } f(n) \in O(g(n))$$

Thus, lower-order terms are also allowed now.

Example:

$$6n + 3 \in O(n^2)$$

O -notation: Example

Intuition: Start from the observation that:

$$f(n) \in \Theta(g(n)) \text{ implies } f(n) \in O(g(n))$$

Thus, **lower-order** terms are also allowed now.

Example:

$$6n + 3 \in O(n^2)$$

If this is true, then we can find positive constants c, n_0 such that:

$$6n + 3 \leq cn^2 \quad \text{for all } n \geq n_0$$

O -notation: Example

Intuition: Start from the observation that:

$$f(n) \in \Theta(g(n)) \text{ implies } f(n) \in O(g(n))$$

Thus, **lower-order** terms are also allowed now.

Example:

$$6n + 3 \in O(n^2)$$

If this is true, then we can find positive constants c, n_0 such that:

$$6n + 3 \leq cn^2 \quad \text{for all } n \geq n_0$$

This holds, for instance, with $c = 6 + 3 = 9$ and $n_0 = 1$.

O -notation: Example

Intuition: Start from the observation that:

$$f(n) \in \Theta(g(n)) \text{ implies } f(n) \in O(g(n))$$

Thus, **lower-order** terms are also allowed now.

Example:

$$6n + 3 \in O(n^2)$$

If this is true, then we can find positive constants c, n_0 such that:

$$6n + 3 \leq cn^2 \quad \text{for all } n \geq n_0$$

This holds, for instance, with $c = 6 + 3 = 9$ and $n_0 = 1$.

- Useful to write (asymptotic) **upper bounds**.

O -notation: Example

Intuition: Start from the observation that:

$$f(n) \in \Theta(g(n)) \text{ implies } f(n) \in O(g(n))$$

Thus, **lower-order** terms are also allowed now.

Example:

$$6n + 3 \in O(n^2)$$

If this is true, then we can find positive constants c, n_0 such that:

$$6n + 3 \leq cn^2 \quad \text{for all } n \geq n_0$$

This holds, for instance, with $c = 6 + 3 = 9$ and $n_0 = 1$.

- Useful to write (asymptotic) **upper bounds**.
- We use it to represent the **worst case** performance.

Ω -notation

Given a function $g(n)$, define the **set of functions**:

$$\Omega(g(n)) = \{f(n) : 0 \leq cg(n) \leq f(n) \text{ for all } n \geq n_0\}$$


where c, n_0 are positive constants.

Ω -notation

Given a function $g(n)$, define the **set of functions**:

$$\Omega(g(n)) = \{f(n) : 0 \leq cg(n) \leq f(n) \text{ for all } n \geq n_0\}$$

where c, n_0 are positive constants.



Ω -notation: Example

Intuition: Again, start from the observation that:

$$f(n) \in \Theta(g(n)) \text{ implies } f(n) \in \Omega(g(n))$$

Thus, **higher-order** terms are allowed now.

Ω -notation: Example

Intuition: Again, start from the observation that:

$$f(n) \in \Theta(g(n)) \text{ implies } f(n) \in \Omega(g(n))$$

Thus, **higher-order** terms are allowed now.

Example:

$$7n^2 - 2n + 14 \in \Omega(n)$$

Ω -notation: Example

Intuition: Again, start from the observation that:

$$f(n) \in \Theta(g(n)) \text{ implies } f(n) \in \Omega(g(n))$$

Thus, **higher-order** terms are allowed now.

Example:

$$7n^2 - 2n + 14 \in \Omega(n)$$

If this is true, then we can find positive constants c, n_0 such that:

$$7n^2 - 2n + 14 \geq cn \quad \text{for all } n \geq n_0$$

Ω -notation: Example

Intuition: Again, start from the observation that:

$$f(n) \in \Theta(g(n)) \text{ implies } f(n) \in \Omega(g(n))$$

Thus, **higher-order** terms are allowed now.

Example:

$$7n^2 - 2n + 14 \in \Omega(n)$$

If this is true, then we can find positive constants c, n_0 such that:

$$7n^2 - 2n + 14 \geq cn \quad \text{for all } n \geq n_0$$

This holds, for instance, with $c = 1$ and $n_0 = 1$.

Ω -notation: Example

Intuition: Again, start from the observation that:

$$f(n) \in \Theta(g(n)) \text{ implies } f(n) \in \Omega(g(n))$$

Thus, **higher-order** terms are allowed now.

Example:

$$7n^2 - 2n + 14 \in \Omega(n)$$

If this is true, then we can find positive constants c, n_0 such that:

$$7n^2 - 2n + 14 \geq cn \quad \text{for all } n \geq n_0$$

This holds, for instance, with $c = 1$ and $n_0 = 1$.

- Useful to write (asymptotic) **lower bounds**.

Ω -notation: Example

Intuition: Again, start from the observation that:

$$f(n) \in \Theta(g(n)) \text{ implies } f(n) \in \Omega(g(n))$$

Thus, **higher-order** terms are allowed now.

Example:

$$7n^2 - 2n + 14 \in \Omega(n)$$

If this is true, then we can find positive constants c, n_0 such that:

$$7n^2 - 2n + 14 \geq cn \quad \text{for all } n \geq n_0$$

This holds, for instance, with $c = 1$ and $n_0 = 1$.

- Useful to write (asymptotic) **lower bounds**.
- We use it to bound the **best case**.

Properties

Not too difficult to prove:

$$f(n) \in \Theta(g(n)) \Leftrightarrow f(n) \in O(g(n)) \text{ and } f(n) \in \Omega(g(n))$$

Properties

Not too difficult to prove:

$$f(n) \in \Theta(g(n)) \Leftrightarrow f(n) \in O(g(n)) \text{ and } f(n) \in \Omega(g(n))$$

Transitivity:

$$f(n) = \Theta(g(n)) \text{ and } g(n) = \Theta(h(n)) \Rightarrow f(n) = \Theta(h(n))$$

$$f(n) = O(g(n)) \text{ and } g(n) = O(h(n)) \Rightarrow f(n) = O(h(n))$$

$$f(n) = \Omega(g(n)) \text{ and } g(n) = \Omega(h(n)) \Rightarrow f(n) = \Omega(h(n))$$

Properties

Not too difficult to prove:

$$f(n) \in \Theta(g(n)) \Leftrightarrow f(n) \in O(g(n)) \text{ and } f(n) \in \Omega(g(n))$$

Transitivity:

$$f(n) = \Theta(g(n)) \text{ and } g(n) = \Theta(h(n)) \Rightarrow f(n) = \Theta(h(n))$$

$$f(n) = O(g(n)) \text{ and } g(n) = O(h(n)) \Rightarrow f(n) = O(h(n))$$

$$f(n) = \Omega(g(n)) \text{ and } g(n) = \Omega(h(n)) \Rightarrow f(n) = \Omega(h(n))$$

Reflexivity:

$$f(n) = \Theta(f(n))$$

$$f(n) = O(f(n))$$

$$f(n) = \Omega(f(n))$$

Properties

Not too difficult to prove:

$$f(n) \in \Theta(g(n)) \Leftrightarrow f(n) \in O(g(n)) \text{ and } f(n) \in \Omega(g(n))$$

Transitivity:

$$f(n) = \Theta(g(n)) \text{ and } g(n) = \Theta(h(n)) \Rightarrow f(n) = \Theta(h(n))$$

$$f(n) = O(g(n)) \text{ and } g(n) = O(h(n)) \Rightarrow f(n) = O(h(n))$$

$$f(n) = \Omega(g(n)) \text{ and } g(n) = \Omega(h(n)) \Rightarrow f(n) = \Omega(h(n))$$

Reflexivity:

$$f(n) = \Theta(f(n))$$

$$f(n) = O(f(n))$$

$$f(n) = \Omega(f(n))$$

Symmetry: $f(n) = \Theta(g(n)) \Leftrightarrow g(n) = \Theta(f(n))$

Properties

Not too difficult to prove:

$$f(n) \in \Theta(g(n)) \Leftrightarrow f(n) \in O(g(n)) \text{ and } f(n) \in \Omega(g(n))$$

Transitivity:

$$f(n) = \Theta(g(n)) \text{ and } g(n) = \Theta(h(n)) \Rightarrow f(n) = \Theta(h(n))$$

$$f(n) = O(g(n)) \text{ and } g(n) = O(h(n)) \Rightarrow f(n) = O(h(n))$$

$$f(n) = \Omega(g(n)) \text{ and } g(n) = \Omega(h(n)) \Rightarrow f(n) = \Omega(h(n))$$

Reflexivity:

$$f(n) = \Theta(f(n))$$

$$f(n) = O(f(n))$$

$$f(n) = \Omega(f(n))$$

Symmetry: $f(n) = \Theta(g(n)) \Leftrightarrow g(n) = \Theta(f(n))$

Transpose symmetry: $f(n) = O(g(n)) \Leftrightarrow g(n) = \Omega(f(n))$

Intuition and counter

The following analogy helps as a mnemonic:

$$f(n) = O(g(n)) \approx a \leq b$$

$$f(n) = \Omega(g(n)) \approx a \geq b$$

$$f(n) = \Theta(g(n)) \approx a = b$$

Intuition and counter

The following analogy helps as a mnemonic:

$$f(n) = O(g(n)) \approx a \leq b$$

$$f(n) = \Omega(g(n)) \approx a \geq b$$

$$f(n) = \Theta(g(n)) \approx a = b$$

But two functions are not always **asymptotically comparable**.

$$f(n) = n \qquad g(n) = n^{1+\sin n}$$

Intuition and counter

The following analogy helps as a mnemonic:

$$f(n) = O(g(n)) \approx a \leq b$$

$$f(n) = \Omega(g(n)) \approx a \geq b$$

$$f(n) = \Theta(g(n)) \approx a = b$$

But two functions are not always **asymptotically comparable**.

$$f(n) = n \quad g(n) = n^{1+\sin n}$$

For these f and g , we have:

$$f(n) \notin O(g(n)) \quad \text{and} \quad f(n) \notin \Omega(g(n))$$

Suggested reading

Chapter 3.1 of:

“Introduction to Algorithms – 2nd Ed.”, Cormen et al.

Skip the o -**notation** and ω -**notation** paragraphs.