

# Computer Graphics

Metric spaces

Emanuele Rodolà  
[rodola@di.uniroma1.it](mailto:rodola@di.uniroma1.it)




SAPIENZA  
UNIVERSITÀ DI ROMA

2nd semester a.y. 2018/2019 · March 21, 2019


## Measuring distance: surfaces

How do you measure **distance** between  $x$  and  $y$  in this picture?



## Measuring distance: surfaces

How do you measure **distance** between  $x$  and  $y$  in this picture?



There is not a unique way!

- You can pass through the sphere with a straight line (**Euclidean**)
- You can walk on the surface in a “straight” path (**non-Euclidean**)

## Measuring distance: images

How do you measure *distance* between  $x$  and  $y$  in this picture?



## Measuring distance: images

How do you measure **distance** between  $x$  and  $y$  in this picture?




Again, not a unique way:

- Natural Euclidean distance
- Smallest amount of pixels between  $x$  and  $y$  (**Manhattan** distance)
- $d(x, y) = 0$  if same **color**,  $d(x, y) = 1$  otherwise


# Euclidean distance

The Euclidean distance measures the length of a **straight line** connecting two points:




# Euclidean distance

The Euclidean distance measures the length of a **straight line** connecting two points:




# Euclidean distance

The Euclidean distance measures the length of a **straight line** connecting two points:



# Euclidean distance


The Euclidean distance measures the length of a **straight line** connecting two points:



Apply Pythagoras' theorem:  $d(a, b) = (\sqrt{|x_b - x_a|^2 + |y_b - y_a|^2})^{\frac{1}{2}}$

# Euclidean distance

The Euclidean distance measures the length of a **straight line** connecting two points:



Apply Pythagoras' theorem:  $d(a, b) = (\lvert x_b - x_a \rvert^2 + \lvert y_b - y_a \rvert^2)^{\frac{1}{2}}$

In vector notation:

$$d(\mathbf{a}, \mathbf{b}) = \|\mathbf{a} - \mathbf{b}\|_2$$

where  $\mathbf{a} = \begin{pmatrix} x_a \\ y_a \end{pmatrix}$  and  $\mathbf{b} = \begin{pmatrix} x_b \\ y_b \end{pmatrix}$

$L_p$  distance in  $\mathbb{R}^k$

One can generalize to different power coefficients  $p \geq 1$ :

$$\|\mathbf{x} - \mathbf{y}\|_2 = (|x_1 - y_1|^2 + |x_2 - y_2|^2)^{\frac{1}{2}}$$
$$\Downarrow$$
$$\|\mathbf{x} - \mathbf{y}\|_p = (|x_1 - y_1|^p + |x_2 - y_2|^p)^{\frac{1}{p}}$$

$L_p$  distance in  $\mathbb{R}^k$

One can generalize to different power coefficients  $p \geq 1$ :

$$\begin{aligned}\|\mathbf{x} - \mathbf{y}\|_2 &= (|x_1 - y_1|^2 + |x_2 - y_2|^2)^{\frac{1}{2}} \\ &\Downarrow \\ \|\mathbf{x} - \mathbf{y}\|_p &= (|x_1 - y_1|^p + |x_2 - y_2|^p)^{\frac{1}{p}}\end{aligned}$$

As well as generalize from  $\mathbb{R}^2$  to  $\mathbb{R}^k$ :

$$\|\mathbf{x} - \mathbf{y}\|_p = \left( \sum_{i=1}^k |x_i - y_i|^p \right)^{\frac{1}{p}}$$

## $L_p$ distance in $\mathbb{R}^k$

One can generalize to different power coefficients  $p \geq 1$ :

$$\|\mathbf{x} - \mathbf{y}\|_2 = (|x_1 - y_1|^2 + |x_2 - y_2|^2)^{\frac{1}{2}}$$
$$\Downarrow$$
$$\|\mathbf{x} - \mathbf{y}\|_p = (|x_1 - y_1|^p + |x_2 - y_2|^p)^{\frac{1}{p}}$$

As well as generalize from  $\mathbb{R}^2$  to  $\mathbb{R}^k$ :


$$\|\mathbf{x} - \mathbf{y}\|_p = \left( \sum_{i=1}^k |x_i - y_i|^p \right)^{\frac{1}{p}}$$

This definition gives us the  $L_p$  distance between vectors in  $\mathbb{R}^k$


Examples:

- Euclidean ( $L_2$ ) distance between 3D points
- Manhattan ( $L_1$ ) distance between cities in a map

# $L_p$ unit balls in $\mathbb{R}^2$



## $L_p$ unit balls on meshes



Each **isoline** identifies points at the same distance from the source

Note that these distances are taken in  $\mathbb{R}^3$ , **not** on the surface!

# Metric spaces

The pair (object, distance) forms a [metric space](#).

## Metric spaces

The pair (object, distance) forms a **metric space**. More formally:

A **set**  $\mathcal{M}$  is a metric space if for every pair of points  $x, y \in \mathcal{M}$  there is a **metric** (or distance) function  $d_{\mathcal{M}} : \mathcal{M} \times \mathcal{M} \rightarrow \mathbb{R}_+$  such that:

# Metric spaces

The pair (object, distance) forms a **metric space**. More formally:

A **set**  $\mathcal{M}$  is a metric space if for every pair of points  $x, y \in \mathcal{M}$  there is a **metric** (or distance) function  $d_{\mathcal{M}} : \mathcal{M} \times \mathcal{M} \rightarrow \mathbb{R}_+$  such that:

- $d_{\mathcal{M}}(x, y) = 0 \Leftrightarrow x = y$  (identity of indiscernibles)

# Metric spaces

The pair (object, distance) forms a **metric space**. More formally:

A **set**  $\mathcal{M}$  is a metric space if for every pair of points  $x, y \in \mathcal{M}$  there is a **metric** (or distance) function  $d_{\mathcal{M}} : \mathcal{M} \times \mathcal{M} \rightarrow \mathbb{R}_+$  such that:

- $d_{\mathcal{M}}(x, y) = 0 \Leftrightarrow x = y$  (identity of indiscernibles)
- $d_{\mathcal{M}}(x, y) = d_{\mathcal{M}}(y, x)$  (symmetry)

# Metric spaces

The pair (object, distance) forms a **metric space**. More formally:

A **set**  $\mathcal{M}$  is a metric space if for every pair of points  $x, y \in \mathcal{M}$  there is a **metric** (or distance) function  $d_{\mathcal{M}} : \mathcal{M} \times \mathcal{M} \rightarrow \mathbb{R}_+$  such that:

- $d_{\mathcal{M}}(x, y) = 0 \Leftrightarrow x = y$  (identity of indiscernibles)
- $d_{\mathcal{M}}(x, y) = d_{\mathcal{M}}(y, x)$  (symmetry)
- $d_{\mathcal{M}}(x, y) \leq d_{\mathcal{M}}(y, z) + d_{\mathcal{M}}(z, x)$  for any  $x, y, z \in \mathcal{M}$  (triangle inequality)

# Metric spaces

The pair (object, distance) forms a **metric space**. More formally:

A **set**  $\mathcal{M}$  is a metric space if for every pair of points  $x, y \in \mathcal{M}$  there is a **metric** (or distance) function  $d_{\mathcal{M}} : \mathcal{M} \times \mathcal{M} \rightarrow \mathbb{R}_+$  such that:

- $d_{\mathcal{M}}(x, y) = 0 \Leftrightarrow x = y$  (identity of indiscernibles)
- $d_{\mathcal{M}}(x, y) = d_{\mathcal{M}}(y, x)$  (symmetry)
- $d_{\mathcal{M}}(x, y) \leq d_{\mathcal{M}}(y, z) + d_{\mathcal{M}}(z, x)$  for any  $x, y, z \in \mathcal{M}$  (triangle inequality)

We will specify a metric space as the pair  $(\mathcal{M}, d_{\mathcal{M}})$

Example:

- The sphere with Euclidean distance is  $(\mathbb{S}^2, d_{L_2})$
- The sphere with geodesic distance is  $(\mathbb{S}^2, d_g)$

## Farthest point sampling (FPS)



Given an initial **seed** point, each new sample is placed **as far as possible** from the previous ones.

This is done following a **greedy** algorithm.

## Farthest point sampling (FPS)

Given an initial **seed** point, each new sample is placed **as far as possible** from the previous ones.

This is done following a **greedy** algorithm.



## Farthest point sampling (FPS)

Given an initial **seed** point, each new sample is placed **as far as possible** from the previous ones.

This is done following a **greedy** algorithm.



FPS clearly depends on an underlying definition of distance function, and we can manipulate colors using a **geometric language**

## Examples: Metric spaces

- $\mathcal{X} = \mathbb{R}$ ,  $d_{\mathcal{X}}(x, y) = |x - y|$

## Examples: Metric spaces

- $\mathcal{X} = \mathbb{R}, \quad d_{\mathcal{X}}(x, y) = |x - y|$
- $\mathcal{X} = \mathcal{A} \subset \mathbb{R}^k, \quad d_{\mathcal{X}}(x, y) = \|x - y\|_2$

## Examples: Metric spaces

- $\mathcal{X} = \mathbb{R}, \quad d_{\mathcal{X}}(x, y) = |x - y|$
- $\mathcal{X} = \mathcal{A} \subset \mathbb{R}^k, \quad d_{\mathcal{X}}(x, y) = \|x - y\|_2$
- $\mathcal{X} = \mathbb{R}, \quad d_{\mathcal{X}}(x, y) = \log(|x - y| + 1)$

## Examples: Metric spaces


- $\mathcal{X} = \mathbb{R}, \quad d_{\mathcal{X}}(x, y) = |x - y|$
- $\mathcal{X} = \mathcal{A} \subset \mathbb{R}^k, \quad d_{\mathcal{X}}(x, y) = \|x - y\|_2$
- $\mathcal{X} = \mathbb{R}, \quad d_{\mathcal{X}}(x, y) = \log(|x - y| + 1)$
- $\mathcal{X} = \text{any set}, \quad d_{\mathcal{X}}(x, y) = \begin{cases} 0 & \text{if } x = y \\ 1 & \text{if } x \neq y \end{cases}$

## Examples: Metric spaces

- $\mathcal{X} = \mathbb{R}, \quad d_{\mathcal{X}}(x, y) = |x - y|$
- $\mathcal{X} = \mathcal{A} \subset \mathbb{R}^k, \quad d_{\mathcal{X}}(x, y) = \|x - y\|_2$
- $\mathcal{X} = \mathbb{R}, \quad d_{\mathcal{X}}(x, y) = \log(|x - y| + 1)$
- $\mathcal{X} = \text{any set}, \quad d_{\mathcal{X}}(x, y) = \begin{cases} 0 & \text{if } x = y \\ 1 & \text{if } x \neq y \end{cases}$
- $\mathcal{X} = \mathcal{A} \times \mathcal{B}, \quad d_{\mathcal{X}}((a_1, b_1), (a_2, b_2)) = \sqrt{d_{\mathcal{A}}(a_1, a_2)^2 + d_{\mathcal{B}}(b_1, b_2)^2}$


## Ambient space and restriction

If  $\mathcal{A}$  is a metric space and  $\mathcal{X} \subset \mathcal{A}$ , then  $\mathcal{A}$  is called **ambient space** for  $\mathcal{X}$ .



## Ambient space and restriction

If  $\mathcal{A}$  is a metric space and  $\mathcal{X} \subset \mathcal{A}$ , then  $\mathcal{A}$  is called **ambient space** for  $\mathcal{X}$ .



A metric on  $\mathcal{X}$  can be obtained by the **restriction**  $d_{\mathcal{X}} = d_{\mathcal{A}|\mathcal{X}}$ , such that:

$$d_{\mathcal{X}}(x, y) = d_{\mathcal{A}}(x, y)$$

for all  $x, y \in \mathcal{X}$

## Isometries

Let  $(\mathcal{M}, d_{\mathcal{M}})$  and  $(\mathcal{N}, d_{\mathcal{N}})$  be two metric spaces.

A bijective map  $f : \mathcal{M} \rightarrow \mathcal{N}$  is called an **isometry** if:

$$d_{\mathcal{M}}(x, y) = d_{\mathcal{N}}(f(x), f(y))$$

for any  $x, y \in \mathcal{M}$ .


# Isometries

Let  $(\mathcal{M}, d_{\mathcal{M}})$  and  $(\mathcal{N}, d_{\mathcal{N}})$  be two metric spaces.

A bijective map  $f : \mathcal{M} \rightarrow \mathcal{N}$  is called an **isometry** if:

$$d_{\mathcal{M}}(x, y) = d_{\mathcal{N}}(f(x), f(y))$$

for any  $x, y \in \mathcal{M}$ .




# Isometries

Let  $(\mathcal{M}, d_{\mathcal{M}})$  and  $(\mathcal{N}, d_{\mathcal{N}})$  be two metric spaces.

A bijective map  $f : \mathcal{M} \rightarrow \mathcal{N}$  is called an **isometry** if:


$$d_{\mathcal{M}}(x, y) = d_{\mathcal{N}}(f(x), f(y))$$

for any  $x, y \in \mathcal{M}$ .




If  $d_{\mathcal{M}} = \|\cdot\|_2$  and  $d_{\mathcal{N}} = \|\cdot\|_2$  we say “rigid isometry”

## Example: Non-rigid “quasi”-isometries



## Example: Non-rigid “quasi”-isometries



$$d_{\mathcal{M}}(x, y) \approx d_{\mathcal{N}}(f(x), f(y))$$

(here  $d_{\mathcal{M}}, d_{\mathcal{N}}$  are geodesic distance functions)


## Example: Rotation in color space

Rotating a color histogram does not change its pairwise distances




## Example: Rotation in color space

Rotating a color histogram does not change its pairwise distances




## Example: Rotation in color space

Rotating a color histogram does not change its pairwise distances



## Example: Rotation in color space

Rotating a color histogram does not change its pairwise distances



## Isometry as equivalence

“Being isometric” is an equivalence relation, since it is:


- reflective ( $a = a$ )
- symmetric ( $a = b \Rightarrow b = a$ )
- transitive ( $a = b \wedge b = c \Rightarrow a = c$ )

# Isometry as equivalence

“Being isometric” is an [equivalence](#) relation, since it is:

- reflective ( $a = a$ )
- symmetric ( $a = b \Rightarrow b = a$ )
- transitive ( $a = b \wedge b = c \Rightarrow a = c$ )

In this sense, we think of isometric shapes as being [the same shape](#):




# Isometry as equivalence

“Being isometric” is an [equivalence relation](#), since it is:

- reflective ( $a = a$ )
- symmetric ( $a = b \Rightarrow b = a$ )
- transitive ( $a = b \wedge b = c \Rightarrow a = c$ )

In this sense, we think of isometric images as being [the same image](#):



# Distance

We now have a notion of **equivalence** between shapes/images. Can we also establish a notion of **distance** between shapes/images?

There are many!


# Distance

We now have a notion of **equivalence** between shapes/images. Can we also establish a notion of **distance** between shapes/images?

There are many!


We start by defining the distance from a point  $x$  to a set  $\mathcal{A} \subseteq (\mathcal{X}, d_{\mathcal{X}})$ :

$$\text{dist}_{\mathcal{X}}(x, \mathcal{A}) = \min_{y \in \mathcal{A}} d_{\mathcal{X}}(x, y)$$



## Hausdorff distance


Consider two subsets  $\mathcal{X}, \mathcal{Y} \subset (\mathcal{Z}, d_{\mathcal{Z}})$ .



# Hausdorff distance

Consider two subsets  $\mathcal{X}, \mathcal{Y} \subset (\mathcal{Z}, d_{\mathcal{Z}})$ .


$$\max_{x \in \mathcal{X}} \text{dist}_{\mathcal{Z}}(x, \mathcal{Y})$$



# Hausdorff distance

Consider two subsets  $\mathcal{X}, \mathcal{Y} \subset (\mathcal{Z}, d_{\mathcal{Z}})$ .


$$\max_{x \in \mathcal{X}} \text{dist}_{\mathcal{Z}}(x, \mathcal{Y}), \max_{y \in \mathcal{Y}} \text{dist}_{\mathcal{Z}}(y, \mathcal{X})$$



# Hausdorff distance

Consider two subsets  $\mathcal{X}, \mathcal{Y} \subset (\mathcal{Z}, d_{\mathcal{Z}})$ .


$$\max \left\{ \max_{x \in \mathcal{X}} \text{dist}_{\mathcal{Z}}(x, \mathcal{Y}), \max_{y \in \mathcal{Y}} \text{dist}_{\mathcal{Z}}(y, \mathcal{X}) \right\}$$



## Hausdorff distance

Consider two subsets  $\mathcal{X}, \mathcal{Y} \subset (\mathcal{Z}, d_{\mathcal{Z}})$ .


$$d_{\mathcal{H}}^{\mathcal{Z}}(\mathcal{X}, \mathcal{Y}) = \max \left\{ \max_{x \in \mathcal{X}} \text{dist}_{\mathcal{Z}}(x, \mathcal{Y}), \max_{y \in \mathcal{Y}} \text{dist}_{\mathcal{Z}}(y, \mathcal{X}) \right\}$$



## Hausdorff distance

Consider two subsets  $\mathcal{X}, \mathcal{Y} \subset (\mathcal{Z}, d_{\mathcal{Z}})$ .

$$d_{\mathcal{H}}^{\mathcal{Z}}(\mathcal{X}, \mathcal{Y}) = \max \left\{ \max_{x \in \mathcal{X}} \text{dist}_{\mathcal{Z}}(x, \mathcal{Y}), \max_{y \in \mathcal{Y}} \text{dist}_{\mathcal{Z}}(y, \mathcal{X}) \right\}$$




The Hausdorff distance is defined between **subsets of a metric space**

Note that perturbing **one single point** can make  $d_{\mathcal{H}}^{\mathcal{Z}}$  arbitrarily large


## Example: Hausdorff distance, rigid case

$d_{\mathcal{H}}^{\mathbb{R}^3}(\mathcal{M}, \mathcal{N})$  is minimum when  $\mathcal{M}$  and  $\mathcal{N}$  are aligned



## Example: Hausdorff distance, non-rigid case

$d_{\mathcal{H}}^{\mathbb{R}^3}(\mathcal{M}, \mathcal{N})$  is minimum when  $\mathcal{M}$  and  $\mathcal{N}$  are aligned



## Example: Hausdorff distance in color space

$d_{\mathcal{H}}^{\mathbb{R}^3}(\mathcal{M}, \mathcal{N})$  is minimum when  $\mathcal{M}$  and  $\mathcal{N}$  have the same color histograms



## Example: Hausdorff distance in color space

$d_{\mathcal{H}}^{\mathbb{R}^3}(\mathcal{M}, \mathcal{N})$  is minimum when  $\mathcal{M}$  and  $\mathcal{N}$  have the same color histograms




## Example: evaluate quality of level-of-detail




"DIZZY" - GEARS OF WAR 2 - MODEL COURTESY OF EPIC GAMES ©

## Shortest paths



## Shortest paths




Euclidean




Geodesic

## Shortest paths on a graph




## Shortest paths on a graph




$$\sqrt{2} = d(v_2, v_3)$$

## Shortest paths on a graph



$$\sqrt{2} = d(v_2, v_3) \neq d(v_1, v_4) = 2$$


## Shortest paths on a graph



$$\sqrt{2} = d(v_2, v_3) \neq d(v_1, v_4) = 2$$

Shortest paths along edges provide upper bounds to exact geodesics

## Shortest paths on a graph




$$\sqrt{2} = d(v_2, v_3) \neq d(v_1, v_4) = 2$$

Shortest paths along edges provide **upper bounds** to exact geodesics

- Still useful with **high resolution** meshes or for **local** distances

## Shortest paths on a graph



$$\sqrt{2} = d(v_2, v_3) \neq d(v_1, v_4) = 2$$

Shortest paths along edges provide **upper bounds** to exact geodesics

- Still useful with **high resolution** meshes or for **local** distances
- Solved by **Dijkstra's algorithm** on the mesh graph

## Voronoi decomposition

For a given sampling  $\mathcal{S}$ , the associated **Voronoi regions** are defined as:


$$V_i(\mathcal{S}) = \{x \in \mathcal{X} : d_{\mathcal{X}}(x, x_i) < d_{\mathcal{X}}(x, x_j), x_j \neq i \in \mathcal{S}\}$$

## Voronoi decomposition

For a given sampling  $\mathcal{S}$ , the associated **Voronoi regions** are defined as:

$$V_i(\mathcal{S}) = \{x \in \mathcal{X} : d_{\mathcal{X}}(x, x_i) < d_{\mathcal{X}}(x, x_j), x_j \neq i \in \mathcal{S}\}$$

In the triangle example:




Three Voronoi regions, associated to the sampling  $\mathcal{S} = \{v_1, v_2, v_3\}$

## Examples: Voronoi decomposition



heat diffusion metric  
on the surface




Euclidean metric in color space  
(result of  $k$ -means)


## Examples: Voronoi decomposition



heat diffusion metric  
on the surface



Euclidean metric in color space  
(result of  $k$ -means)



## Exercise: Farthest point sampling

Implement FPS using this algorithm:

- Fix  $n$  and let  $\mathcal{S}^{(0)} = \{y\}$  for some  $y \in \mathcal{X}$
- Proceed recursively:
  - At step  $k$ , given  $\mathcal{S}^{(k-1)}$ , select  $x \in (\mathcal{X}, d_{\mathcal{X}})$  such that
$$x = \arg \max_{x \in \mathcal{X}} d_{\mathcal{X}}(x, \mathcal{S}^{(k-1)})$$
  - Set  $\mathcal{S}^{(k)} = \mathcal{S}^{(k-1)} \cup x$
  - Repeat until  $k = n$
- Test with different starting points  $y$
- Test with a fixed starting point and gradually increasing  $n$

Use FPS to define the initial seed for  $k$ -means in color space.