

# Computer Graphics

## Recap of linear algebra II

Arianna Rampini  
[rampini@di.uniroma1.it](mailto:rampini@di.uniroma1.it)



## Recap: Bases

A **basis** of  $V$  is a collection of vectors in  $V$  that is **linearly independent** and **spans**  $V$

- $\text{span}(v_1, \dots, v_n) = \{a_1v_1 + \dots + a_nv_n : a_1, \dots, a_n \in \mathbb{R}\}$
- $v_1, \dots, v_n \in V$  are **linearly independent** if and only if each  $v \in \text{span}(v_1, \dots, v_n)$  has only one representation as a linear combination of  $v_1, \dots, v_n$

You can think of a basis as the minimal set of vectors that generates the entire space

## Recap: Bases

Every vector  $v \in V$  can be expressed **uniquely** as a linear combination:

$$v = \sum_{i=1}^n c_i v_i$$

Considering the "vector matrix"  $n \times 1$  and the "basis matrix"  $n \times n$

$$\mathbf{c} = \begin{pmatrix} c_1 \\ \vdots \\ c_n \end{pmatrix} \quad \mathbf{V} = \begin{pmatrix} | & \cdots & | \\ v_1 & \cdots & v_n \\ | & \cdots & | \end{pmatrix}$$

the linear combination can be in **matrix notation** as:

$$v = \mathbf{V}\mathbf{c}$$

## Recap: Matrices

Consider a linear map  $T : V \rightarrow W$ , a basis  $v_1, \dots, v_n \in V$  and a basis  $w_1, \dots, w_m \in W$ .

The **matrix** of  $T$  in these bases is the  $m \times n$  array of values in  $\mathbb{R}$

$$\mathbf{T} = \begin{pmatrix} T_{1,1} & \cdots & T_{1,n} \\ \vdots & & \vdots \\ T_{m,1} & \cdots & T_{m,n} \end{pmatrix}$$

whose entries  $T_{i,j}$  are defined by

$$Tv_j = T_{1,j}w_1 + \cdots + T_{m,j}w_m$$

In other words, the matrix encodes **how basis vectors are mapped**, and this is enough to map all other vectors in their span, since:

$$Tv = T\left(\sum_j c_j v_j\right) = \sum_j T(c_j v_j) = \sum_j c_j Tv_j$$

## Recap: Product of “map matrix” and “vector matrix”

Using the matrix of  $T$  we have:

$$Tv = \mathbf{T}\mathbf{c}$$

because

$$\underbrace{\begin{pmatrix} T_{1,1} & \cdots & T_{1,n} \\ \vdots & & \vdots \\ T_{m,1} & \cdots & T_{m,n} \end{pmatrix}}_{\mathbf{T}} \underbrace{\begin{pmatrix} c_1 \\ \vdots \\ c_n \end{pmatrix}}_{\mathbf{c}} = \sum_{j=1}^n c_j \underbrace{\begin{pmatrix} T_{1,j} \\ \vdots \\ T_{m,j} \end{pmatrix}}_{Tv_j \text{ wrt } (w_1, \dots, w_m)}$$

(Recall that, for bases  $v_1, \dots, v_n \in V$  and  $w_1, \dots, w_m \in W$ :

$$Tv_j = T_{1,j}w_1 + \cdots + T_{m,j}w_m)$$

We see then that vector  $v = \sum_j c_j v_j$  is mapped to  $Tv = \sum_j c_j Tv_j$ .

In other words, matrix product is behaving as expected.

# Rank of a matrix

The **rank** of a matrix  $\mathbf{A} \in \mathbb{R}^{m \times n}$  is the dimension of the span of its columns

**Example:**

$$\mathbf{A} = \begin{pmatrix} 4 & 7 & 1 & 8 \\ 3 & 5 & 2 & 9 \end{pmatrix}$$

# Rank of a matrix

The **rank** of a matrix  $\mathbf{A} \in \mathbb{R}^{m \times n}$  is the dimension of the span of its columns

**Example:**

$$\mathbf{A} = \begin{pmatrix} 4 & 7 & 1 & 8 \\ 3 & 5 & 2 & 9 \end{pmatrix}$$

The rank is the dimension of

$$\text{span} \left( \begin{pmatrix} 4 \\ 3 \end{pmatrix}, \begin{pmatrix} 7 \\ 5 \end{pmatrix}, \begin{pmatrix} 1 \\ 2 \end{pmatrix}, \begin{pmatrix} 8 \\ 9 \end{pmatrix} \right) \in \mathbb{R}^2$$

# Rank of a matrix

The **rank** of a matrix  $\mathbf{A} \in \mathbb{R}^{m \times n}$  is the dimension of the span of its columns

**Example:**

$$\mathbf{A} = \begin{pmatrix} 4 & 7 & 1 & 8 \\ 3 & 5 & 2 & 9 \end{pmatrix}$$

The rank is the dimension of

$$\text{span} \left( \begin{pmatrix} 4 \\ 3 \end{pmatrix}, \begin{pmatrix} 7 \\ 5 \end{pmatrix}, \begin{pmatrix} 1 \\ 2 \end{pmatrix}, \begin{pmatrix} 8 \\ 9 \end{pmatrix} \right) \in \mathbb{R}^2$$

This span can not have dimension larger than 2, because  $\dim(\mathbb{R}^2) = 2$ .

In this example,  $\text{rank}(\mathbf{A}) = 2$



# Rank of a matrix

The **rank** of a matrix  $\mathbf{A} \in \mathbb{R}^{m \times n}$  is the dimension of the span of its columns

**Example:**

$$\mathbf{A} = \begin{pmatrix} 4 & 7 & 1 & 8 \\ 3 & 5 & 2 & 9 \end{pmatrix}$$

The rank is the dimension of

$$\text{span} \left( \begin{pmatrix} 4 \\ 3 \end{pmatrix}, \begin{pmatrix} 7 \\ 5 \end{pmatrix}, \begin{pmatrix} 1 \\ 2 \end{pmatrix}, \begin{pmatrix} 8 \\ 9 \end{pmatrix} \right) \in \mathbb{R}^2$$

This span can not have dimension larger than 2, because  $\dim(\mathbb{R}^2) = 2$ .

In this example,  $\text{rank}(\mathbf{A}) = 2$

Note that this result **does not depend** on a choice of basis, i.e., change of basis **preserves the rank**

## Example: Reduced bases

Consider the  $\mathbb{R}^{n \times k}$  matrix

$$\mathbf{V} = \begin{pmatrix} | & \cdots & \cdots & | \\ \mathbf{v}_1 & \cdots & \cdots & \mathbf{v}_k \\ | & \cdots & \cdots & | \end{pmatrix}$$

containing basis vectors as its columns, and the  $\mathbb{R}^{n \times k'}$  matrix

$$\mathbf{V}' = \begin{pmatrix} | & \cdots & | \\ \mathbf{v}_1 & \cdots & \mathbf{v}_{k'} \\ | & \cdots & | \end{pmatrix}$$

obtained by **truncating**  $\mathbf{V}$  to the first  $k' < k$  columns

## Example: Reduced bases

Consider the  $\mathbb{R}^{n \times k}$  matrix

$$\mathbf{V} = \begin{pmatrix} | & \cdots & \cdots & | \\ \mathbf{v}_1 & \cdots & \cdots & \mathbf{v}_k \\ | & \cdots & \cdots & | \end{pmatrix}$$

containing basis vectors as its columns, and the  $\mathbb{R}^{n \times k'}$  matrix

$$\mathbf{V}' = \begin{pmatrix} | & \cdots & | \\ \mathbf{v}_1 & \cdots & \mathbf{v}_{k'} \\ | & \cdots & | \end{pmatrix}$$

obtained by **truncating**  $\mathbf{V}$  to the first  $k' < k$  columns

Then,  $k = \text{rank}(\mathbf{V}) > \text{rank}(\mathbf{V}') = k'$

The rank reflects the expressive power of the **full** ( $\mathbf{V}$ ) and **reduced** ( $\mathbf{V}'$ ) bases

## Example: Reduced bases



full basis  
 $\text{rank}(\mathbf{V}) = k$



reduced basis  
 $\text{rank}(\mathbf{V}') = k' < k$

# Invariant subspaces

A subspace  $U$  of  $V$  is called **invariant** under  $T : V \rightarrow V$  if:

$$u \in U \quad \text{implies} \quad Tu \in U$$

# Invariant subspaces

A subspace  $U$  of  $V$  is called **invariant** under  $T : V \rightarrow V$  if:

$$u \in U \quad \text{implies} \quad Tu \in U$$

If  $T : V \rightarrow V$  also means  $T : U \rightarrow U$ , then  $U$  is an invariant subspace.

# Invariant subspaces

A subspace  $U$  of  $V$  is called **invariant** under  $T : V \rightarrow V$  if:

$$u \in U \quad \text{implies} \quad Tu \in U$$

If  $T : V \rightarrow V$  also means  $T : U \rightarrow U$ , then  $U$  is an invariant subspace.

Consider the 1-dimensional subspace:

$$U = \{\alpha v : \alpha \in \mathbb{R}\} = \text{span}(v)$$

# Invariant subspaces

A subspace  $U$  of  $V$  is called **invariant** under  $T : V \rightarrow V$  if:

$$u \in U \quad \text{implies} \quad Tu \in U$$

If  $T : V \rightarrow V$  also means  $T : U \rightarrow U$ , then  $U$  is an invariant subspace.

Consider the 1-dimensional subspace:

$$U = \{\alpha v : \alpha \in \mathbb{R}\} = \text{span}(v)$$

If  $U$  is invariant under  $T$ , then for some  $\lambda \in \mathbb{R}$ :

$$Tv = \lambda v$$



# Invariant subspaces

A subspace  $U$  of  $V$  is called **invariant** under  $T : V \rightarrow V$  if:

$$u \in U \quad \text{implies} \quad Tu \in U$$

If  $T : V \rightarrow V$  also means  $T : U \rightarrow U$ , then  $U$  is an invariant subspace.

Consider the 1-dimensional subspace:

$$U = \{\alpha v : \alpha \in \mathbb{R}\} = \text{span}(v)$$

If  $U$  is invariant under  $T$ , then for some  $\lambda \in \mathbb{R}$ :

$$Tv = \lambda v$$

Conversely, if  $Tv = \lambda v$  for some  $\lambda \in \mathbb{R}$ , then  $\text{span}(v)$  is a 1-dimensional subspace of  $V$  invariant under  $T$

# Eigenvalues and eigenvectors

$$Tv = \lambda v$$

Studying **invariant subspaces** is a key tool for studying linear maps

# Eigenvalues and eigenvectors

$$Tv = \lambda v$$

Studying **invariant subspaces** is a key tool for studying linear maps

We call  $\lambda$  an **eigenvalue** of  $T : V \rightarrow V$  if the above holds for some vector  $v \neq 0$ , called the associated **eigenvector**

# Eigenvalues and eigenvectors

$$Tv = \lambda v$$

Studying **invariant subspaces** is a key tool for studying linear maps

We call  $\lambda$  an **eigenvalue** of  $T : V \rightarrow V$  if the above holds for some vector  $v \neq 0$ , called the associated **eigenvector**

If the equation holds for  $m$  distinct eigenvalues and eigenvectors:

$$\begin{aligned}Tv_1 &= \lambda_1 v_1 \\ &\vdots \\ Tv_m &= \lambda_m v_m\end{aligned}$$

Then it can be proven that  $v_1, \dots, v_m$  are **linearly independent**

# Eigenvalues and eigenvectors

$$Tv = \lambda v$$

Studying **invariant subspaces** is a key tool for studying linear maps

We call  $\lambda$  an **eigenvalue** of  $T : V \rightarrow V$  if the above holds for some vector  $v \neq 0$ , called the associated **eigenvector**

If the equation holds for  $m$  distinct eigenvalues and eigenvectors:

$$\begin{aligned}Tv_1 &= \lambda_1 v_1 \\ &\vdots \\ Tv_m &= \lambda_m v_m\end{aligned}$$

Then it can be proven that  $v_1, \dots, v_m$  are **linearly independent**, and it must be:

$$m \leq \dim(V)$$

# Eigenvalues and eigenvectors

$$Tv_1 = \lambda_1 v_1$$

$$\vdots$$

$$Tv_m = \lambda_m v_m$$

In a sense, eigenvectors provide a **decomposition** of  $V$  into subspaces

# Eigenvalues and eigenvectors

$$Tv_1 = \lambda_1 v_1$$

$$\vdots$$

$$Tv_m = \lambda_m v_m$$

In a sense, eigenvectors provide a **decomposition** of  $V$  into subspaces

Additional notes:

- This decomposition only makes sense for  $T : V \rightarrow V$

# Eigenvalues and eigenvectors

$$Tv_1 = \lambda_1 v_1$$

$$\vdots$$

$$Tv_m = \lambda_m v_m$$

In a sense, eigenvectors provide a **decomposition** of  $V$  into subspaces

Additional notes:

- This decomposition only makes sense for  $T : V \rightarrow V$
- If  $V$  is a function space, eigenvectors are called **eigenfunctions**



# Eigenspaces

If distinct eigenvectors  $E = (v_1, \dots, v_m)$  correspond to the same eigenvalue  $\lambda$ , then  $E$  spans an eigenspace of  $T$

An eigenspace is a subspace of  $V$ , but has dimension greater than 1

# Eigenspaces

If distinct eigenvectors  $E = (v_1, \dots, v_m)$  correspond to the same eigenvalue  $\lambda$ , then  $E$  spans an eigenspace of  $T$

An eigenspace is a subspace of  $V$ , but has dimension greater than 1

To summarize:

- Certain linear maps  $T : V \rightarrow V$  induce invariant subspaces on  $V$

# Eigenspaces

If distinct eigenvectors  $E = (v_1, \dots, v_m)$  correspond to the same eigenvalue  $\lambda$ , then  $E$  spans an eigenspace of  $T$

An eigenspace is a subspace of  $V$ , but has dimension greater than 1

To summarize:

- Certain linear maps  $T : V \rightarrow V$  induce invariant subspaces on  $V$
- These subspaces are spanned by eigenvectors of  $T$

# Eigenspaces

If distinct eigenvectors  $E = (v_1, \dots, v_m)$  correspond to the same eigenvalue  $\lambda$ , then  $E$  spans an eigenspace of  $T$

An eigenspace is a subspace of  $V$ , but has dimension greater than 1

To summarize:

- Certain linear maps  $T : V \rightarrow V$  induce invariant subspaces on  $V$
- These subspaces are spanned by eigenvectors of  $T$
- Subspaces might be high-dimensional and are called eigenspaces

# Eigenspaces

If distinct eigenvectors  $E = (v_1, \dots, v_m)$  correspond to the same eigenvalue  $\lambda$ , then  $E$  spans an eigenspace of  $T$

An eigenspace is a subspace of  $V$ , but has dimension greater than 1

To summarize:

- Certain linear maps  $T : V \rightarrow V$  induce invariant subspaces on  $V$
- These subspaces are spanned by eigenvectors of  $T$
- Subspaces might be high-dimensional and are called eigenspaces
- Eigenspaces provide a form of decomposition of  $V$

# Inner product

We want to be able to measure **lengths** and **angles** among vectors

**Note:** We assume our vectors are **real-valued**

# Inner product

We want to be able to measure **lengths** and **angles** among vectors

To do so, we define the **inner product** as a function  $\langle u, v \rangle : V \times V \rightarrow \mathbb{R}$  with the properties:

- **non-negativity:**  $\langle v, v \rangle \geq 0$  for all  $v \in V$

**Note:** We assume our vectors are **real-valued**

# Inner product

We want to be able to measure **lengths** and **angles** among vectors

To do so, we define the **inner product** as a function  $\langle u, v \rangle : V \times V \rightarrow \mathbb{R}$  with the properties:

- **non-negativity:**  $\langle v, v \rangle \geq 0$  for all  $v \in V$
- **definiteness:**  $\langle v, v \rangle = 0$  iff  $v = 0$

**Note:** We assume our vectors are **real-valued**



# Inner product

We want to be able to measure **lengths** and **angles** among vectors

To do so, we define the **inner product** as a function  $\langle u, v \rangle : V \times V \rightarrow \mathbb{R}$  with the properties:

- **non-negativity:**  $\langle v, v \rangle \geq 0$  for all  $v \in V$
- **definiteness:**  $\langle v, v \rangle = 0$  iff  $v = 0$
- **additivity:**  $\langle u + v, w \rangle = \langle u, w \rangle + \langle v, w \rangle$  for all  $u, v, w \in V$

**Note:** We assume our vectors are **real-valued**

# Inner product

We want to be able to measure **lengths** and **angles** among vectors

To do so, we define the **inner product** as a function  $\langle u, v \rangle : V \times V \rightarrow \mathbb{R}$  with the properties:

- **non-negativity:**  $\langle v, v \rangle \geq 0$  for all  $v \in V$
- **definiteness:**  $\langle v, v \rangle = 0$  iff  $v = 0$
- **additivity:**  $\langle u + v, w \rangle = \langle u, w \rangle + \langle v, w \rangle$  for all  $u, v, w \in V$
- **homogeneity:**  $\langle \lambda u, v \rangle = \lambda \langle u, v \rangle$  for all  $\lambda \in \mathbb{R}$  and all  $u, v \in V$

**Note:** We assume our vectors are **real-valued**

# Inner product

We want to be able to measure **lengths** and **angles** among vectors

To do so, we define the **inner product** as a function  $\langle u, v \rangle : V \times V \rightarrow \mathbb{R}$  with the properties:

- **non-negativity:**  $\langle v, v \rangle \geq 0$  for all  $v \in V$
- **definiteness:**  $\langle v, v \rangle = 0$  iff  $v = 0$
- **additivity:**  $\langle u + v, w \rangle = \langle u, w \rangle + \langle v, w \rangle$  for all  $u, v, w \in V$
- **homogeneity:**  $\langle \lambda u, v \rangle = \lambda \langle u, v \rangle$  for all  $\lambda \in \mathbb{R}$  and all  $u, v \in V$
- **symmetry:**  $\langle u, v \rangle = \langle v, u \rangle$  for all  $u, v \in V$

**Note:** We assume our vectors are **real-valued**

# Examples: Inner products

- **Lists:**

The **Euclidean inner product** (or dot product) is defined by

$$\langle (u_1, \dots, u_n), (v_1, \dots, v_n) \rangle = u_1 v_1 + \dots + u_n v_n$$

This is the standard inner product for vectors in  $\mathbb{R}^n$

# Examples: Inner products

- **Lists:**

The **Euclidean inner product** (or dot product) is defined by

$$\langle (u_1, \dots, u_n), (v_1, \dots, v_n) \rangle = u_1 v_1 + \dots + u_n v_n$$

This is the standard inner product for vectors in  $\mathbb{R}^n$

- **Functions:**

On the vector space of continuous functions  $f : [-1, 1] \rightarrow \mathbb{R}$

$$\langle f, g \rangle = \int_{-1}^1 f(x)g(x)dx$$

# Norm

Each inner product determines a **norm**:

$$\|v\| = \sqrt{\langle v, v \rangle}$$

# Norm

Each inner product determines a **norm**:

$$\|v\| = \sqrt{\langle v, v \rangle}$$

## Examples:

- For lists with the Euclidean inner product:

$$\|(v_1, \dots, v_n)\| = \sqrt{v_1^2 + \dots + v_n^2}$$

# Norm

Each inner product determines a **norm**:

$$\|v\| = \sqrt{\langle v, v \rangle}$$

## Examples:

- For lists with the Euclidean inner product:

$$\|(v_1, \dots, v_n)\| = \sqrt{v_1^2 + \dots + v_n^2}$$

- For continuous functions  $f : [-1, 1] \rightarrow \mathbb{R}$ :

$$\|f\| = \sqrt{\int_{-1}^1 f(x)^2 dx}$$



# Orthogonality

Two vectors  $u, v \in V$  are **orthogonal** if  $\langle u, v \rangle = 0$

# Orthogonality

Two vectors  $u, v \in V$  are **orthogonal** if  $\langle u, v \rangle = 0$

For  $u, v \in \mathbb{R}^2$ , it can be shown that

$$\langle u, v \rangle = \|u\| \|v\| \cos \theta$$

where  $\theta \in \mathbb{R}$  is the angle between  $u, v$  if we think of them as **arrows** with initial point at the origin

# Orthogonality

Two vectors  $u, v \in V$  are **orthogonal** if  $\langle u, v \rangle = 0$

For  $u, v \in \mathbb{R}^2$ , it can be shown that

$$\langle u, v \rangle = \|u\| \|v\| \cos \theta$$

where  $\theta \in \mathbb{R}$  is the angle between  $u, v$  if we think of them as **arrows** with initial point at the origin

From this, we can think of the inner product as encoding a general notion of **angle** between two vectors:

$$\theta = \arccos \frac{\langle u, v \rangle}{\|u\| \|v\|}$$

For example, we can now think of “angle between two functions”

# Orthogonal bases

A basis  $(v_1, \dots, v_n)$  is orthogonal if all the vectors are orthogonal to each other

# Orthogonal bases

A basis  $(v_1, \dots, v_n)$  is orthogonal if all the vectors are orthogonal to each other; the basis is **orthonormal** if, in addition,  $\|v_i\| = 1$  for all  $v_i$

# Orthogonal bases

A basis  $(v_1, \dots, v_n)$  is orthogonal if all the vectors are orthogonal to each other; the basis is **orthonormal** if, in addition,  $\|v_i\| = 1$  for all  $v_i$

Note that every collection of orthogonal vectors is **linearly independent**

# Orthogonal bases

A basis  $(v_1, \dots, v_n)$  is orthogonal if all the vectors are orthogonal to each other; the basis is **orthonormal** if, in addition,  $\|v_i\| = 1$  for all  $v_i$

Note that every collection of orthogonal vectors is **linearly independent**

**Examples:**

# Orthogonal bases

A basis  $(v_1, \dots, v_n)$  is orthogonal if all the vectors are orthogonal to each other; the basis is **orthonormal** if, in addition,  $\|v_i\| = 1$  for all  $v_i$

Note that every collection of orthogonal vectors is **linearly independent**

## Examples:

- The standard basis in  $\mathbb{R}^n$  is orthonormal



# Orthogonal bases

A basis  $(v_1, \dots, v_n)$  is orthogonal if all the vectors are orthogonal to each other; the basis is **orthonormal** if, in addition,  $\|v_i\| = 1$  for all  $v_i$

Note that every collection of orthogonal vectors is **linearly independent**

## Examples:

- The standard basis in  $\mathbb{R}^n$  is orthonormal
- $(\frac{1}{2}, \frac{\sqrt{3}}{2}), (-\frac{\sqrt{3}}{2}, \frac{1}{2}) \in \mathbb{R}^2$  is orthonormal

# Orthogonal bases

A basis  $(v_1, \dots, v_n)$  is orthogonal if all the vectors are orthogonal to each other; the basis is **orthonormal** if, in addition,  $\|v_i\| = 1$  for all  $v_i$

Note that every collection of orthogonal vectors is **linearly independent**

## Examples:

- The standard basis in  $\mathbb{R}^n$  is orthonormal
- $(\frac{1}{2}, \frac{\sqrt{3}}{2}), (-\frac{\sqrt{3}}{2}, \frac{1}{2}) \in \mathbb{R}^2$  is orthonormal

# Orthogonal bases

A basis  $(v_1, \dots, v_n)$  is orthogonal if all the vectors are orthogonal to each other; the basis is **orthonormal** if, in addition,  $\|v_i\| = 1$  for all  $v_i$

Note that every collection of orthogonal vectors is **linearly independent**

## Examples:

- The standard basis in  $\mathbb{R}^n$  is orthonormal
- $(\frac{1}{2}, \frac{\sqrt{3}}{2}), (-\frac{\sqrt{3}}{2}, \frac{1}{2}) \in \mathbb{R}^2$  is orthonormal

Note that **any** orthogonal basis can be made **orthonormal** by rescaling each basis vector:

$$v_i \mapsto \frac{v_i}{\|v_i\|}$$

# Orthonormal bases

Given an orthonormal basis,  $v \in V$  can be written as a linear combination:

$$v = \langle v, v_1 \rangle v_1 + \cdots + \langle v, v_n \rangle v_n$$

So the combination coefficients are simply given by inner products

# Inner product in matrix notation

For vectors  $u, v \in V$  in the **standard basis**  $\{e_i\}$ , we can write:

$$\langle u, v \rangle = \left\langle \sum_i u_i e_i, \sum_j v_j e_j \right\rangle$$

# Inner product in matrix notation

For vectors  $u, v \in V$  in the **standard basis**  $\{e_i\}$ , we can write:

$$\begin{aligned}\langle u, v \rangle &= \left\langle \sum_i u_i e_i, \sum_j v_j e_j \right\rangle \\ &= \sum_{i,j} \langle u_i e_i, v_j e_j \rangle\end{aligned}$$

# Inner product in matrix notation

For vectors  $u, v \in V$  in the **standard basis**  $\{e_i\}$ , we can write:

$$\begin{aligned}\langle u, v \rangle &= \left\langle \sum_i u_i e_i, \sum_j v_j e_j \right\rangle \\ &= \sum_{i,j} \langle u_i e_i, v_j e_j \rangle \\ &= \sum_{i,j} u_i v_j \underbrace{\langle e_i, e_j \rangle}_{=0 \text{ if } i \neq j}\end{aligned}$$

# Inner product in matrix notation

For vectors  $u, v \in V$  in the **standard basis**  $\{e_i\}$ , we can write:

$$\begin{aligned}\langle u, v \rangle &= \left\langle \sum_i u_i e_i, \sum_j v_j e_j \right\rangle \\ &= \sum_{i,j} \langle u_i e_i, v_j e_j \rangle \\ &= \sum_{i,j} u_i v_j \underbrace{\langle e_i, e_j \rangle}_{=0 \text{ if } i \neq j} \\ &= \sum_i u_i v_i\end{aligned}$$

which corresponds to the standard Euclidean inner product



# Inner product in matrix notation

For vectors  $u, v \in V$  in the **standard basis**  $\{e_i\}$ , we can write:

$$\begin{aligned}\langle u, v \rangle &= \left\langle \sum_i u_i e_i, \sum_j v_j e_j \right\rangle \\ &= \sum_{i,j} \langle u_i e_i, v_j e_j \rangle \\ &= \sum_{i,j} u_i v_j \underbrace{\langle e_i, e_j \rangle}_{=0 \text{ if } i \neq j} \\ &= \sum_i u_i v_i\end{aligned}$$

which corresponds to the standard Euclidean inner product

In matrix notation, we can thus write

$$\langle u, v \rangle = \mathbf{u}^\top \mathbf{v}$$

# Inner product in matrix notation

For vectors  $u, v \in V$  in some other basis  $\{w_i\}$ , we can write:

$$\langle u, v \rangle = \sum_{i,j} u_i v_j \langle w_i, w_j \rangle$$

## Inner product in matrix notation

For vectors  $u, v \in V$  in some other basis  $\{w_i\}$ , we can write:

$$\langle u, v \rangle = \sum_{i,j} u_i v_j \underbrace{\langle w_i, w_j \rangle}_{\mathbf{w}_i^\top \mathbf{w}_j}$$

# Inner product in matrix notation

For vectors  $u, v \in V$  in some other basis  $\{w_i\}$ , we can write:

$$\begin{aligned}\langle u, v \rangle &= \sum_{i,j} u_i v_j \underbrace{\langle w_i, w_j \rangle}_{\mathbf{w}_i^\top \mathbf{w}_j} \\ &= \mathbf{u}^\top \mathbf{W}^\top \mathbf{W} \mathbf{v}\end{aligned}$$

where  $\mathbf{W}$  contains the basis vectors  $\mathbf{w}_i$  as its columns

# Inner product in matrix notation

For vectors  $u, v \in V$  in some other basis  $\{w_i\}$ , we can write:

$$\begin{aligned}\langle u, v \rangle &= \sum_{i,j} u_i v_j \underbrace{\langle w_i, w_j \rangle}_{\mathbf{w}_i^\top \mathbf{w}_j} \\ &= \mathbf{u}^\top \mathbf{W}^\top \mathbf{W} \mathbf{v}\end{aligned}$$

where  $\mathbf{W}$  contains the basis vectors  $\mathbf{w}_i$  as its columns

- If  $\{w_i\}$  is orthonormal, then  $\mathbf{W}^\top \mathbf{W} = \mathbf{I}$

# Inner product in matrix notation

For vectors  $u, v \in V$  in some other basis  $\{w_i\}$ , we can write:

$$\begin{aligned}\langle u, v \rangle &= \sum_{i,j} u_i v_j \underbrace{\langle w_i, w_j \rangle}_{\mathbf{w}_i^\top \mathbf{w}_j} \\ &= \mathbf{u}^\top \mathbf{W}^\top \mathbf{W} \mathbf{v}\end{aligned}$$

where  $\mathbf{W}$  contains the basis vectors  $\mathbf{w}_i$  as its columns

- If  $\{w_i\}$  is orthonormal, then  $\mathbf{W}^\top \mathbf{W} = \mathbf{I}$
- Another way to see the derivation above is as follows:

# Inner product in matrix notation

For vectors  $u, v \in V$  in some other basis  $\{w_i\}$ , we can write:

$$\begin{aligned}\langle u, v \rangle &= \sum_{i,j} u_i v_j \underbrace{\langle w_i, w_j \rangle}_{\mathbf{w}_i^\top \mathbf{w}_j} \\ &= \mathbf{u}^\top \mathbf{W}^\top \mathbf{W} \mathbf{v}\end{aligned}$$

where  $\mathbf{W}$  contains the basis vectors  $\mathbf{w}_i$  as its columns

- If  $\{w_i\}$  is orthonormal, then  $\mathbf{W}^\top \mathbf{W} = \mathbf{I}$
- Another way to see the derivation above is as follows:
  - $\mathbf{W}\mathbf{u}$  gives the coefficients of  $u$  in the **standard** basis (and similarly for  $v$ )

# Inner product in matrix notation

For vectors  $u, v \in V$  in some other basis  $\{w_i\}$ , we can write:

$$\begin{aligned}\langle u, v \rangle &= \sum_{i,j} u_i v_j \underbrace{\langle w_i, w_j \rangle}_{\mathbf{w}_i^\top \mathbf{w}_j} \\ &= \mathbf{u}^\top \mathbf{W}^\top \mathbf{W} \mathbf{v}\end{aligned}$$

where  $\mathbf{W}$  contains the basis vectors  $\mathbf{w}_i$  as its columns

- If  $\{w_i\}$  is orthonormal, then  $\mathbf{W}^\top \mathbf{W} = \mathbf{I}$
- Another way to see the derivation above is as follows:
  - $\mathbf{W}\mathbf{u}$  gives the coefficients of  $u$  in the **standard** basis (and similarly for  $v$ )
  - Once we are in the standard basis, we can write

$$\langle u, v \rangle = (\mathbf{W}\mathbf{u})^\top (\mathbf{W}\mathbf{v})$$



# Inner product in matrix notation

For vectors  $u, v \in V$  in some other basis  $\{w_i\}$ , we can write:

$$\begin{aligned}\langle u, v \rangle &= \sum_{i,j} u_i v_j \underbrace{\langle w_i, w_j \rangle}_{\mathbf{w}_i^\top \mathbf{w}_j} \\ &= \mathbf{u}^\top \mathbf{W}^\top \mathbf{W} \mathbf{v}\end{aligned}$$


where  $\mathbf{W}$  contains the basis vectors  $\mathbf{w}_i$  as its columns

- If  $\{w_i\}$  is orthonormal, then  $\mathbf{W}^\top \mathbf{W} = \mathbf{I}$
- Another way to see the derivation above is as follows:
  - $\mathbf{W}\mathbf{u}$  gives the coefficients of  $u$  in the **standard** basis (and similarly for  $v$ )
  - Once we are in the standard basis, we can write



$$\langle u, v \rangle = (\mathbf{W}\mathbf{u})^\top (\mathbf{W}\mathbf{v}) = \mathbf{u}^\top \mathbf{W}^\top \mathbf{W} \mathbf{v}$$

# Exercise


Find the two **linear maps**  $\mathbf{R}_1$  and  $\mathbf{R}_2$  that produce the following reflections:



$\mathbf{R}_1$



$\mathbf{R}_2$



## Suggested reading

See sections 3.F, 5.A – 6.B of:

S. Axler, “Linear algebra done right – 3rd edition”. Springer, 2015