

Computer Graphics


Local coordinates

Emanuele Rodolà
rodola@di.uniroma1.it



Exercises

- Farthest point sampling



Adjacency matrices as operators

We can see adjacency matrices as **operators** when applied to functions

For example, $\mathbf{g} = \mathbf{A}\mathbf{f}$ yields a **vertex-based** function g defined as:

$$g(v_i) = \sum_{e_{ij} \in E} f(v_j)$$

Adjacency matrices as operators

We can see adjacency matrices as **operators** when applied to functions

For example, $\mathbf{g} = \mathbf{A}\mathbf{f}$ yields a **vertex-based** function g defined as:

$$g(v_i) = \sum_{e_{ij} \in E} f(v_j)$$

On this observation, one can construct new operators such as $\mathbf{I} - \mathbf{A}$:

$$g(v_i) = f(v_i) - \sum_{e_{ij} \in E} f(v_j)$$

Adjacency matrices as operators

We can see adjacency matrices as **operators** when applied to functions

For example, $\mathbf{g} = \mathbf{A}\mathbf{f}$ yields a **vertex-based** function g defined as:


$$g(v_i) = \sum_{e_{ij} \in E} f(v_j)$$

On this observation, one can construct new operators such as $\mathbf{I} - \mathbf{A}$:

$$g(v_i) = f(v_i) - \sum_{e_{ij} \in E} f(v_j)$$

And similarly for more complex (but **linear**) operations

Example: Hole filling



original
 \mathbf{f}

noisy
 $\tilde{\mathbf{f}}$

denoised
 $\mathbf{A}\tilde{\mathbf{f}}$

Graph Laplacian

Given a mesh graph $G = (V, E)$, consider this condition on vertex v_i :

$$\mathbf{v}_i - \frac{1}{d_i} \sum_{j:(i,j) \in E} \mathbf{v}_j = 0$$

where d_i is the valence of v_i

Graph Laplacian

Given a **mesh graph** $G = (V, E)$, consider this condition on vertex v_i :

$$\mathbf{v}_i - \frac{1}{d_i} \sum_{j:(i,j) \in E} \mathbf{v}_j = 0$$

where d_i is the valence of v_i

The equation is satisfied if v_i lies in the **barycenter** of its neighbors

Graph Laplacian

Given a **mesh graph** $G = (V, E)$, consider this condition on vertex v_i :

$$\mathbf{v}_i - \frac{1}{d_i} \sum_{j:(i,j) \in E} \mathbf{v}_j = 0$$

where d_i is the valence of v_i

The equation is satisfied if v_i lies in the **barycenter** of its neighbors

In matrix notation, we define the $n \times n$ matrix \mathbf{L} as:

$$L_{ij} = \begin{cases} 1 & \text{if } i = j \\ -\frac{1}{d_i} & \text{if } e_{ij} \in E \\ 0 & \text{otherwise} \end{cases}$$

also known as the **graph Laplacian** of G

Graph Laplacian

Given a **mesh graph** $G = (V, E)$, consider this condition on vertex v_i :

$$\mathbf{v}_i - \frac{1}{d_i} \sum_{j:(i,j) \in E} \mathbf{v}_j = 0$$

where d_i is the valence of v_i

The equation is satisfied if v_i lies in the **barycenter** of its neighbors

In matrix notation, we define the $n \times n$ matrix \mathbf{L} as:

$$L_{ij} = \begin{cases} 1 & \text{if } i = j \\ -\frac{1}{d_i} & \text{if } e_{ij} \in E \\ 0 & \text{otherwise} \end{cases}$$

also known as the **graph Laplacian** of G ; we get to:

$$\mathbf{L}\mathbf{V} = \mathbf{0}$$

Least squares meshes

Is there an embedding satisfying $\mathbf{L}\mathbf{V} = \mathbf{0}$?

- Trivial embedding $\mathbf{V} = \mathbf{0}$

Least squares meshes

Is there an embedding satisfying $\mathbf{L}\mathbf{V} = \mathbf{0}$?

- Trivial embedding $\mathbf{V} = \mathbf{0}$
- One-dimensional subspace $\text{span}(\mathbf{1})$

Least squares meshes

Is there an **interesting** embedding satisfying $\mathbf{L}\mathbf{V} = \mathbf{0}$?

- Trivial embedding $\mathbf{V} = \mathbf{0}$
- One-dimensional subspace $\text{span}(\mathbf{1})$

Least squares meshes

Is there an **interesting** embedding satisfying $\mathbf{L}\mathbf{V} = \mathbf{0}$?

- Trivial embedding $\mathbf{V} = \mathbf{0}$
- One-dimensional subspace $\text{span}(\mathbf{1})$

Assume $m \geq 1$ **anchor vertices** $v_s \in \mathcal{A}$ with known 3D position

Least squares meshes

Is there an **interesting** embedding satisfying $\mathbf{L}\mathbf{V} = \mathbf{0}$?

- Trivial embedding $\mathbf{V} = \mathbf{0}$
- One-dimensional subspace $\text{span}(\mathbf{1})$

Assume $m \geq 1$ **anchor vertices** $v_s \in \mathcal{A}$ with known 3D position

Then, consider the linear system:

$$\begin{pmatrix} \mathbf{L} \\ \mathbf{A} \end{pmatrix} \mathbf{V} = \mathbf{b}$$

where

$$a_{ij} = \begin{cases} 1 & \text{if } v_j \in \mathcal{A} \\ 0 & \text{otherwise} \end{cases}, \quad b_k = \begin{cases} (0, 0, 0) & k \leq n \\ \mathbf{v}_{s_{k-n}} & n < k \leq n + m \end{cases}$$

Least squares meshes

$$\begin{pmatrix} \mathbf{L} \\ \mathbf{A} \end{pmatrix} \mathbf{v} \approx \mathbf{b}$$

Least squares meshes


$$\min_{\mathbf{V} \in \mathbb{R}^{n \times 3}} \left\| \begin{pmatrix} \mathbf{L} \\ \mathbf{A} \end{pmatrix} \mathbf{V} - \mathbf{b} \right\|_2^2$$

Least squares meshes

$$\min_{\mathbf{V} \in \mathbb{R}^{n \times 3}} \|\mathbf{L}\mathbf{V}\|_2^2 + \sum_{v_i \in \mathcal{A}} \|\mathbf{v}_i - \mathbf{b}_i\|_2^2$$


Least squares meshes

$$\min_{\mathbf{V} \in \mathbb{R}^{n \times 3}} \|\mathbf{L}\mathbf{V}\|_2^2 + \sum_{v_i \in \mathcal{A}} \|\mathbf{v}_i - \mathbf{b}_i\|_2^2$$




- Anchor constraints are not satisfied exactly
- At higher resolution, error distributes better among the constraints

Least squares meshes



Move the anchor positions to do **shape modeling**

Least squares meshes




Move the anchor positions to do **shape modeling**




Barycentric coordinates

How to express p in terms of v_1, v_2, v_3 ?



Barycentric coordinates

How to express p in terms of v_1, v_2, v_3 ?



In other words, we want a new [coordinate system](#) where

$$v_1 = (1, 0, 0)$$

$$v_2 = (0, 1, 0)$$


$$v_3 = (0, 0, 1)$$

and

$$p = (\lambda_1, \lambda_2, \lambda_3)$$

Barycentric coordinates

How to express p in terms of v_1, v_2, v_3 ?



In particular, we want a **convex combination**


$$p = \lambda_1 v_1 + \lambda_2 v_2 + \lambda_3 v_3$$

with

$$\lambda_1, \lambda_2, \lambda_3 \geq 0, \quad \lambda_1 + \lambda_2 + \lambda_3 = 1$$

Barycentric coordinates

How to express p in terms of v_1, v_2, v_3 ?




We come to the linear system

$$\begin{pmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \\ 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} \lambda_1 \\ \lambda_2 \\ \lambda_3 \end{pmatrix} = \begin{pmatrix} x \\ y \\ 1 \end{pmatrix}$$

which has a **unique** solution

Barycentric coordinates

How to express p in terms of v_1, v_2, v_3 ?



Solving the system, we get the closed form expressions:


$$\lambda_1 = \frac{(y_2 - y_3)(x - x_3) + (x_3 - x_2)(y - y_3)}{(y_2 - y_3)(x_1 - x_3) + (x_3 - x_2)(y_1 - y_3)}$$

$$\lambda_2 = \frac{(y_3 - y_1)(x - x_3) + (x_1 - x_3)(y - y_3)}{(y_2 - y_3)(x_1 - x_3) + (x_3 - x_2)(y_1 - y_3)}$$

$$\lambda_3 = 1 - \lambda_1 - \lambda_2$$

Barycentric coordinates

How to express p in terms of v_1, v_2, v_3 ?




Solving the system, we get the closed form expressions:

$$\lambda_1 = \frac{(y_2 - y_3)(x - x_3) + (x_3 - x_2)(y - y_3)}{(y_2 - y_3)(x_1 - x_3) + (x_3 - x_2)(y_1 - y_3)}$$
$$\lambda_2 = \frac{(y_3 - y_1)(x - x_3) + (x_1 - x_3)(y - y_3)}{(y_2 - y_3)(x_1 - x_3) + (x_3 - x_2)(y_1 - y_3)}$$
$$\lambda_3 = 1 - \lambda_1 - \lambda_2$$

If p is **outside** the triangle, we get at least one **negative** coordinate!


Barycentric coordinates: interpretation

The barycentric coordinates are proportional to the **triangle areas**



Barycentric coordinates: interpretation


The barycentric coordinates are proportional to the **triangle areas**



The triangle **centroid** (or **barycenter**) has coordinates $(0.33, 0.33, 0.33)$

Example: linear interpolation

Barycentric coordinates can be used to compute **linear interpolation** of function values:




Given function values $f(x_1)$, $f(x_2)$, $f(x_3)$ at the 3 vertices, the function values $f(x)$ inside the triangle are obtained by **barycentric coordinates**

See lesson 5 - "Visualization I" (Mar 13, 2019)

Mean value coordinates


This can also be seen as a triangle mesh with 4 vertices:



So we are asking: How to express any vertex v_0 as a combination of its neighbors?

Mean value coordinates

We consider the more general case with a **non-convex** polygon




Where we look for **coordinates** such that

$$v_0 = \sum_i \lambda_i v_i, \quad \lambda_i > 0, \quad \sum_i \lambda_i = 1$$

Mean value coordinates

We consider the more general case with a **non-convex** polygon




Where we look for **coordinates** such that

$$v_0 = \sum_i \lambda_i v_i, \quad \lambda_i > 0, \quad \sum_i \lambda_i = 1$$

There is **no unique solution!**

Mean value coordinates

We consider the more general case with a **non-convex** polygon




One possible solution is given by **Floater's mean value coordinates**:

$$w_i = \frac{\tan(\alpha/2) + \tan(\beta/2)}{\|v_i - v_0\|}, \quad \lambda_i = \frac{w_i}{\sum_j w_j}$$

Mean value coordinates

We consider the more general case with a **non-convex** polygon



One possible solution is given by **Floater's mean value coordinates**:

$$w_i = \frac{\tan(\alpha/2) + \tan(\beta/2)}{\|v_i - v_0\|}, \quad \lambda_i = \frac{w_i}{\sum_j w_j}$$



Floater, "Mean value coordinates". CAGD 20(1), 2003

Example

