

# Fundamentals of Computer Graphics

Mesh processing I

Emanuele Rodolà  
[rodola@di.uniroma1.it](mailto:rodola@di.uniroma1.it)



SAPIENZA  
UNIVERSITÀ DI ROMA

# Adjacency matrices

The mesh connectivity can be encoded in **adjacency matrices**

Let  $|V| = n$ ,  $|E| = e$ ,  $|F| = m$  for a mesh  $M = (V, E, F)$

## Adjacency matrices: Vertex-to-vertex

The mesh connectivity can be encoded in **adjacency matrices**

Let  $|V| = n$ ,  $|E| = e$ ,  $|F| = m$  for a mesh  $M = (V, E, F)$

The **vertex-to-vertex** adjacency is defined as the  $n \times n$  binary matrix:

$$\mathbf{A} = \begin{pmatrix} 0 & 1 & 0 & \cdots & 1 \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ 1 & 0 & 1 & \cdots & 0 \end{pmatrix}$$

where  $a_{ij} = 1$  if vertex  $v_i$  is connected to  $v_j$  (that is,  $e_{ij} \in E$ )

# Adjacency matrices: Vertex-to-vertex

The mesh connectivity can be encoded in **adjacency matrices**

Let  $|V| = n$ ,  $|E| = e$ ,  $|F| = m$  for a mesh  $M = (V, E, F)$

The **vertex-to-vertex** adjacency is defined as the  $n \times n$  binary matrix:

$$\mathbf{A} = \begin{pmatrix} 0 & 1 & 0 & \cdots & 1 \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ 1 & 0 & 1 & \cdots & 0 \end{pmatrix}$$

where  $a_{ij} = 1$  if vertex  $v_i$  is connected to  $v_j$  (that is,  $e_{ij} \in E$ )

- The **diagonal** is always 0

# Adjacency matrices: Vertex-to-vertex

The mesh connectivity can be encoded in **adjacency matrices**

Let  $|V| = n$ ,  $|E| = e$ ,  $|F| = m$  for a mesh  $M = (V, E, F)$

The **vertex-to-vertex** adjacency is defined as the  $n \times n$  binary matrix:

$$\mathbf{A} = \begin{pmatrix} 0 & 1 & 0 & \cdots & 1 \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ 1 & 0 & 1 & \cdots & 0 \end{pmatrix}$$

where  $a_{ij} = 1$  if vertex  $v_i$  is connected to  $v_j$  (that is,  $e_{ij} \in E$ )

- The **diagonal** is always 0
- $\mathbf{A}$  is **symmetric**

# Adjacency matrices: Vertex-to-vertex

The mesh connectivity can be encoded in **adjacency matrices**

Let  $|V| = n$ ,  $|E| = e$ ,  $|F| = m$  for a mesh  $M = (V, E, F)$

The **vertex-to-vertex** adjacency is defined as the  $n \times n$  binary matrix:

$$\mathbf{A} = \begin{pmatrix} 0 & 1 & 0 & \cdots & 1 \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ 1 & 0 & 1 & \cdots & 0 \end{pmatrix}$$

where  $a_{ij} = 1$  if vertex  $v_i$  is connected to  $v_j$  (that is,  $e_{ij} \in E$ )

- The **diagonal** is always 0
- $\mathbf{A}$  is **symmetric**
- Each row and column has at least one 1 (that is,  $\sum_{ij} a_{ij} = e$ )

# Adjacency matrices: Vertex-to-triangle

The mesh connectivity can be encoded in [adjacency matrices](#)

Let  $|V| = n$ ,  $|E| = e$ ,  $|F| = m$  for a mesh  $M = (V, E, F)$

The [vertex-to-triangle](#) adjacency is defined as the  $n \times m$  binary matrix:

$$\mathbf{P} = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 & 1 & 1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 1 & 0 & \cdots & 1 & 0 & 1 \end{pmatrix}$$

where  $p_{ij} = 1$  if vertex  $v_i$  belongs to triangle  $t_j$

# Adjacency matrices: Vertex-to-triangle

The mesh connectivity can be encoded in [adjacency matrices](#)

Let  $|V| = n$ ,  $|E| = e$ ,  $|F| = m$  for a mesh  $M = (V, E, F)$

The [vertex-to-triangle](#) adjacency is defined as the  $n \times m$  binary matrix:

$$\mathbf{P} = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 & 1 & 1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 1 & 0 & \cdots & 1 & 0 & 1 \end{pmatrix}$$

where  $p_{ij} = 1$  if vertex  $v_i$  belongs to triangle  $t_j$

- Each [row](#) has at least one 1 (each vertex belongs to some triangle)



# Adjacency matrices: Vertex-to-triangle

The mesh connectivity can be encoded in **adjacency matrices**

Let  $|V| = n$ ,  $|E| = e$ ,  $|F| = m$  for a mesh  $M = (V, E, F)$

The **vertex-to-triangle** adjacency is defined as the  $n \times m$  binary matrix:

$$\mathbf{P} = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 & 1 & 1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 1 & 0 & \cdots & 1 & 0 & 1 \end{pmatrix}$$

where  $p_{ij} = 1$  if vertex  $v_i$  belongs to triangle  $t_j$

- Each **row** has at least one 1 (each vertex belongs to some triangle)
- Each **column** sums up to 3 (each triangle has exactly 3 vertices)

# Adjacency matrices: Vertex-vertex co-occurrence

Consider the product:

$$\mathbf{P}\mathbf{P}^{\top} = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 & 1 & 1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 1 & 0 & \cdots & 1 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & \cdots & \cdots & 0 \\ 0 & \cdots & \cdots & 1 \\ 0 & \cdots & \cdots & 0 \\ \vdots & \cdots & \cdots & \vdots \\ 0 & \cdots & \cdots & 1 \\ 1 & \cdots & \cdots & 0 \\ 1 & \cdots & \cdots & 1 \end{pmatrix}$$

# Adjacency matrices: Vertex-vertex co-occurrence

Consider the product:

$$\mathbf{PP}^\top = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 & 1 & 1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 1 & 0 & \cdots & 1 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & \cdots & \cdots & 0 \\ 0 & \cdots & \cdots & 1 \\ 0 & \cdots & \cdots & 0 \\ \vdots & \cdots & \cdots & \vdots \\ 0 & \cdots & \cdots & 1 \\ 1 & \cdots & \cdots & 0 \\ 1 & \cdots & \cdots & 1 \end{pmatrix}$$

$\mathbf{PP}^\top$  is a  $n \times n$  matrix with the same **zero pattern** as  $\mathbf{A}$

# Adjacency matrices: Vertex-vertex co-occurrence

Consider the product:

$$\mathbf{PP}^\top = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 & 1 & 1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots & \vdots \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots & \vdots \\ 0 & 1 & 0 & \cdots & 1 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & \cdots & \cdots & 0 \\ 0 & \cdots & \cdots & 1 \\ 0 & \cdots & \cdots & 0 \\ \vdots & \cdots & \cdots & \vdots \\ 0 & \cdots & \cdots & 1 \\ 1 & \cdots & \cdots & 0 \\ 1 & \cdots & \cdots & 1 \end{pmatrix}$$

$\mathbf{PP}^\top$  is a  $n \times n$  matrix with the same **zero pattern** as  $\mathbf{A}$

Cell  $(i, j)$  counts the number of **triangles** where  $v_i$  and  $v_j$  **co-occur**

# Adjacency matrices: Vertex-vertex co-occurrence

Consider the product:

$$\mathbf{PP}^\top = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 & 1 & 1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots & \vdots \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots & \vdots \\ 0 & 1 & 0 & \cdots & 1 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & \cdots & \cdots & 0 \\ 0 & \cdots & \cdots & 1 \\ 0 & \cdots & \cdots & 0 \\ \vdots & \cdots & \cdots & \vdots \\ 0 & \cdots & \cdots & 1 \\ 1 & \cdots & \cdots & 0 \\ 1 & \cdots & \cdots & 1 \end{pmatrix}$$

$\mathbf{PP}^\top$  is a  $n \times n$  matrix with the same **zero pattern** as  $\mathbf{A}$

Cell  $(i, j)$  counts the number of **triangles** where  $v_i$  and  $v_j$  **co-occur**

- $\text{diag}(\mathbf{PP}^\top)$  contains the number of triangles for each vertex

# Adjacency matrices: Vertex-vertex co-occurrence

Consider the product:

$$\mathbf{PP}^\top = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 & 1 & 1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots & \vdots \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots & \vdots \\ 0 & 1 & 0 & \cdots & 1 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & \cdots & \cdots & 0 \\ 0 & \cdots & \cdots & 1 \\ 0 & \cdots & \cdots & 0 \\ \vdots & \cdots & \cdots & \vdots \\ 0 & \cdots & \cdots & 1 \\ 1 & \cdots & \cdots & 0 \\ 1 & \cdots & \cdots & 1 \end{pmatrix}$$

$\mathbf{PP}^\top$  is a  $n \times n$  matrix with the same **zero pattern** as  $\mathbf{A}$

Cell  $(i, j)$  counts the number of **triangles** where  $v_i$  and  $v_j$  **co-occur**

- $\text{diag}(\mathbf{PP}^\top)$  contains the number of triangles for each vertex
- All other values are 0 ( $e_{ij} \notin E$ )

# Adjacency matrices: Vertex-vertex co-occurrence

Consider the product:

$$\mathbf{PP}^\top = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 & 1 & 1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots & \vdots \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots & \vdots \\ 0 & 1 & 0 & \cdots & 1 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & \cdots & \cdots & 0 \\ 0 & \cdots & \cdots & 1 \\ 0 & \cdots & \cdots & 0 \\ \vdots & \cdots & \cdots & \vdots \\ 0 & \cdots & \cdots & 1 \\ 1 & \cdots & \cdots & 0 \\ 1 & \cdots & \cdots & 1 \end{pmatrix}$$

$\mathbf{PP}^\top$  is a  $n \times n$  matrix with the same **zero pattern** as  $\mathbf{A}$

Cell  $(i, j)$  counts the number of **triangles** where  $v_i$  and  $v_j$  **co-occur**

- $\text{diag}(\mathbf{PP}^\top)$  contains the number of triangles for each vertex
- All other values are 0 ( $e_{ij} \notin E$ ), 1 (boundary)

# Adjacency matrices: Vertex-vertex co-occurrence

Consider the product:

$$\mathbf{PP}^\top = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 & 1 & 1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots & \vdots \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots & \vdots \\ 0 & 1 & 0 & \cdots & 1 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & \cdots & \cdots & 0 \\ 0 & \cdots & \cdots & 1 \\ 0 & \cdots & \cdots & 0 \\ \vdots & \cdots & \cdots & \vdots \\ 0 & \cdots & \cdots & 1 \\ 1 & \cdots & \cdots & 0 \\ 1 & \cdots & \cdots & 1 \end{pmatrix}$$

$\mathbf{PP}^\top$  is a  $n \times n$  matrix with the same **zero pattern** as  $\mathbf{A}$

Cell  $(i, j)$  counts the number of **triangles** where  $v_i$  and  $v_j$  **co-occur**

- $\text{diag}(\mathbf{PP}^\top)$  contains the number of triangles for each vertex
- All other values are 0 ( $e_{ij} \notin E$ ), 1 (boundary), or 2 (interior)



## Adjacency matrices: Triangle-to-triangle

Consider the product:

$$\mathbf{P}^\top \mathbf{P} = \begin{pmatrix} 1 & \dots & \dots & 0 \\ 0 & \dots & \dots & 1 \\ 0 & \dots & \dots & 0 \\ \vdots & \dots & \dots & \vdots \\ 0 & \dots & \dots & 1 \\ 1 & \dots & \dots & 0 \\ 1 & \dots & \dots & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & \dots & 0 & 1 & 1 \\ \vdots & \dots & \dots & \dots & \dots & \dots & \vdots \\ \vdots & \dots & \dots & \dots & \dots & \dots & \vdots \\ 0 & 1 & 0 & \dots & 1 & 0 & 1 \end{pmatrix}$$

which is a  $m \times m$  matrix

# Adjacency matrices: Triangle-to-triangle

Consider the product:

$$\mathbf{P}^\top \mathbf{P} = \begin{pmatrix} 1 & \dots & \dots & 0 \\ 0 & \dots & \dots & 1 \\ 0 & \dots & \dots & 0 \\ \vdots & \dots & \dots & \vdots \\ 0 & \dots & \dots & 1 \\ 1 & \dots & \dots & 0 \\ 1 & \dots & \dots & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & \dots & 0 & 1 & 1 \\ \vdots & \dots & \dots & \dots & \dots & \dots & \vdots \\ \vdots & \dots & \dots & \dots & \dots & \dots & \vdots \\ 0 & 1 & 0 & \dots & 1 & 0 & 1 \end{pmatrix}$$

which is a  $m \times m$  matrix

Cell  $(i, j)$  counts the number of **vertices** that  $t_i$  and  $t_j$  **share**

# Adjacency matrices: Triangle-to-triangle

Consider the product:

$$\mathbf{P}^\top \mathbf{P} = \begin{pmatrix} 1 & \dots & \dots & 0 \\ 0 & \dots & \dots & 1 \\ 0 & \dots & \dots & 0 \\ \vdots & \dots & \dots & \vdots \\ 0 & \dots & \dots & 1 \\ 1 & \dots & \dots & 0 \\ 1 & \dots & \dots & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & \dots & 0 & 1 & 1 \\ \vdots & \dots & \dots & \dots & \dots & \dots & \vdots \\ \vdots & \dots & \dots & \dots & \dots & \dots & \vdots \\ 0 & 1 & 0 & \dots & 1 & 0 & 1 \end{pmatrix}$$

which is a  $m \times m$  matrix

Cell  $(i, j)$  counts the number of **vertices** that  $t_i$  and  $t_j$  **share**

- $\text{diag}(\mathbf{P}\mathbf{P}^\top)$  is always 3 (=number of vertices per triangle)

# Adjacency matrices: Triangle-to-triangle

Consider the product:

$$\mathbf{P}^\top \mathbf{P} = \begin{pmatrix} 1 & \dots & \dots & 0 \\ 0 & \dots & \dots & 1 \\ 0 & \dots & \dots & 0 \\ \vdots & \dots & \dots & \vdots \\ 0 & \dots & \dots & 1 \\ 1 & \dots & \dots & 0 \\ 1 & \dots & \dots & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & \dots & 0 & 1 & 1 \\ \vdots & \dots & \dots & \dots & \dots & \vdots & \vdots \\ \vdots & \dots & \dots & \dots & \dots & \vdots & \vdots \\ 0 & 1 & 0 & \dots & 1 & 0 & 1 \end{pmatrix}$$

which is a  $m \times m$  matrix

Cell  $(i, j)$  counts the number of **vertices** that  $t_i$  and  $t_j$  **share**

- $\text{diag}(\mathbf{P}\mathbf{P}^\top)$  is always 3 (=number of vertices per triangle)
- All other values are 0 (non-adjacent triangles)

# Adjacency matrices: Triangle-to-triangle

Consider the product:

$$\mathbf{P}^\top \mathbf{P} = \begin{pmatrix} 1 & \dots & \dots & 0 \\ 0 & \dots & \dots & 1 \\ 0 & \dots & \dots & 0 \\ \vdots & \dots & \dots & \vdots \\ 0 & \dots & \dots & 1 \\ 1 & \dots & \dots & 0 \\ 1 & \dots & \dots & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & \dots & 0 & 1 & 1 \\ \vdots & \dots & \dots & \dots & \dots & \vdots \\ \vdots & \dots & \dots & \dots & \dots & \vdots \\ 0 & 1 & 0 & \dots & 1 & 0 & 1 \end{pmatrix}$$

which is a  $m \times m$  matrix

Cell  $(i, j)$  counts the number of **vertices** that  $t_i$  and  $t_j$  **share**

- $\text{diag}(\mathbf{P}\mathbf{P}^\top)$  is always 3 (=number of vertices per triangle)
- All other values are 0 (non-adjacent triangles), 1 (triangles share one vertex)

# Adjacency matrices: Triangle-to-triangle

Consider the product:


$$\mathbf{P}^\top \mathbf{P} = \begin{pmatrix} 1 & \dots & \dots & 0 \\ 0 & \dots & \dots & 1 \\ 0 & \dots & \dots & 0 \\ \vdots & \dots & \dots & \vdots \\ 0 & \dots & \dots & 1 \\ 1 & \dots & \dots & 0 \\ 1 & \dots & \dots & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & \dots & 0 & 1 & 1 \\ \vdots & \dots & \dots & \dots & \dots & \dots & \vdots \\ \vdots & \dots & \dots & \dots & \dots & \dots & \vdots \\ 0 & 1 & 0 & \dots & 1 & 0 & 1 \end{pmatrix}$$

which is a  $m \times m$  matrix


Cell  $(i, j)$  counts the number of **vertices** that  $t_i$  and  $t_j$  **share**

- $\text{diag}(\mathbf{P}\mathbf{P}^\top)$  is always 3 (=number of vertices per triangle)
- All other values are 0 (non-adjacent triangles), 1 (triangles share one vertex), or 2 (triangles share an edge)


# Examples: Adjacency



vertex-to-vertex




vertex-to-triangle




triangle-to-triangle


## Examples: Adjacency



vertex-to-vertex



vertex-to-triangle



triangle-to-triangle

In general we have  $m \approx 2n$ , and with  $n$  in the order of several thousands these adjacency matrices can be **very large** (quadratic in  $n$ )

It is advisable to use **sparse** data structures to store them



## Adjacency matrices: Powers

The  $k$ -th power of  $\mathbf{A}$  corresponds to composing  $\mathbf{A}$  with itself  $k \geq 1$  times

For example, for  $k = 2$ :

$$\mathbf{A}^2 = \mathbf{A}\mathbf{A} = \begin{pmatrix} 0 & 1 & 0 & \cdots & 1 \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ 1 & 0 & 1 & \cdots & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 & 0 & \cdots & 1 \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ 1 & 0 & 1 & \cdots & 0 \end{pmatrix}$$

## Adjacency matrices: Powers

The  $k$ -th power of  $\mathbf{A}$  corresponds to composing  $\mathbf{A}$  with itself  $k \geq 1$  times

For example, for  $k = 2$ :

$$\mathbf{A}^2 = \mathbf{A}\mathbf{A} = \begin{pmatrix} 0 & 1 & 0 & \cdots & 1 \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ 1 & 0 & 1 & \cdots & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 & 0 & \cdots & 1 \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ 1 & 0 & 1 & \cdots & 0 \end{pmatrix}$$

The result is a  $n \times n$  matrix encoding **2nd order adjacency**

## Adjacency matrices: Powers

The  $k$ -th power of  $\mathbf{A}$  corresponds to composing  $\mathbf{A}$  with itself  $k \geq 1$  times

For  $k > 1$ :

$$\mathbf{A}^k = \mathbf{A} \cdots \mathbf{A} = \begin{pmatrix} 0 & 1 & 0 & \cdots & 1 \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ 1 & 0 & 1 & \cdots & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & 1 & 0 & \cdots & 1 \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ 1 & 0 & 1 & \cdots & 0 \end{pmatrix}$$

The result is a  $n \times n$  matrix encoding  $k$ -th order adjacency

## Adjacency matrices: Powers

The  $k$ -th power of  $\mathbf{A}$  corresponds to composing  $\mathbf{A}$  with itself  $k \geq 1$  times

For  $k > 1$ :

$$\mathbf{A}^k = \mathbf{A} \cdots \mathbf{A} = \begin{pmatrix} 0 & 1 & 0 & \cdots & 1 \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ 1 & 0 & 1 & \cdots & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & 1 & 0 & \cdots & 1 \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ 1 & 0 & 1 & \cdots & 0 \end{pmatrix}$$

The result is a  $n \times n$  matrix encoding  $k$ -th order adjacency

- The same applies to triangle-to-triangle adjacency
- The  $k$ -th power of vertex-to-triangle is given by  $\mathbf{A}^k \mathbf{P}$

# Adjacency matrices: Powers

The  $k$ -th power of  $\mathbf{A}$  corresponds to composing  $\mathbf{A}$  with itself  $k \geq 1$  times

For  $k > 1$ :


$$\mathbf{A}^k = \mathbf{A} \cdots \mathbf{A} = \begin{pmatrix} 0 & 1 & 0 & \cdots & 1 \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ 1 & 0 & 1 & \cdots & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & 1 & 0 & \cdots & 1 \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ 1 & 0 & 1 & \cdots & 0 \end{pmatrix}$$

The result is a  $n \times n$  matrix encoding  $k$ -th order adjacency


- The same applies to triangle-to-triangle adjacency
- The  $k$ -th power of vertex-to-triangle is given by  $\mathbf{A}^k \mathbf{P}$

Manipulating adjacency is useful in many tasks relying upon local context


# Examples: Powers



vertex-to-vertex  
 $k = 1$




vertex-to-triangle  
 $k = 1$




triangle-to-triangle  
 $k = 1$


# Examples: Powers



vertex-to-vertex  
 $k = 2$




vertex-to-triangle  
 $k = 2$




triangle-to-triangle  
 $k = 2$


# Examples: Powers



vertex-to-vertex  
 $k = 3$



vertex-to-triangle  
 $k = 3$



triangle-to-triangle  
 $k = 3$



# Adjacency matrices as operators

We can see adjacency matrices as **operators** when applied to functions

For example,  $\mathbf{g} = \mathbf{A}\mathbf{f}$  yields a **vertex-based** function  $g$  defined as:

$$g(v_i) = \sum_{e_{ij} \in E} f(v_j)$$

# Adjacency matrices as operators

We can see adjacency matrices as **operators** when applied to functions

For example,  $\mathbf{g} = \mathbf{A}\mathbf{f}$  yields a **vertex-based** function  $g$  defined as:

$$g(v_i) = \sum_{e_{ij} \in E} f(v_j)$$

On this observation, one can construct new operators such as  $\mathbf{I} - \mathbf{A}$ :

$$g(v_i) = f(v_i) - \sum_{e_{ij} \in E} f(v_j)$$

# Adjacency matrices as operators

We can see adjacency matrices as **operators** when applied to functions

For example,  $\mathbf{g} = \mathbf{A}\mathbf{f}$  yields a **vertex-based** function  $g$  defined as:

$$g(v_i) = \sum_{e_{ij} \in E} f(v_j)$$

On this observation, one can construct new operators such as  $\mathbf{I} - \mathbf{A}$ :

$$g(v_i) = f(v_i) - \sum_{e_{ij} \in E} f(v_j)$$

And similarly for **triangle-based** functions

# Adjacency matrices: Point clouds

Adjacency is a general notion that can be extended to **point clouds**

(Such notions of adjacency can of course be used on meshes as well)

# Adjacency matrices: Point clouds

Adjacency is a general notion that can be extended to **point clouds**

For example, use **Euclidean distance** within a threshold  $\tau$ :

$$a_{ij} = \begin{cases} 1 & \text{if } \|\mathbf{v}_i - \mathbf{v}_j\|_2 \leq \tau \\ 0 & \text{otherwise} \end{cases}$$




(Such notions of adjacency can of course be used on meshes as well)

# Adjacency matrices: Point clouds


Adjacency is a general notion that can be extended to **point clouds**

For example, use **Euclidean distance** within a threshold  $\tau$ :

$$a_{ij} = \begin{cases} 1 & \text{if } \|\mathbf{v}_i - \mathbf{v}_j\|_2 \leq \tau \\ 0 & \text{otherwise} \end{cases}$$



$k = 1$




$k = 2$

Similarly to before,  $\mathbf{A}^k$  encodes  $k$ -th order adjacency

(Such notions of adjacency can of course be used on meshes as well)

## Example: Hole filling




original  
 $\mathbf{f}$

noisy  
 $\tilde{\mathbf{f}}$


denoised  
 $\mathbf{A}\tilde{\mathbf{f}}$

## Shortest paths





## Shortest paths




Euclidean




Geodesic

# Shortest paths on a graph




# Shortest paths on a graph




$$\sqrt{2} = d(v_2, v_3)$$

# Shortest paths on a graph



$$\sqrt{2} = d(v_2, v_3) \neq d(v_1, v_4) = 2$$


# Shortest paths on a graph



$$\sqrt{2} = d(v_2, v_3) \neq d(v_1, v_4) = 2$$

Shortest paths along edges provide **upper bounds** to exact geodesics

# Shortest paths on a graph




$$\sqrt{2} = d(v_2, v_3) \neq d(v_1, v_4) = 2$$

Shortest paths along edges provide **upper bounds** to exact geodesics

- Still useful with **high resolution** meshes or for **local** distances

# Shortest paths on a graph



$$\sqrt{2} = d(v_2, v_3) \neq d(v_1, v_4) = 2$$

Shortest paths along edges provide **upper bounds** to exact geodesics

- Still useful with **high resolution** meshes or for **local** distances
- Solved by **Dijkstra's algorithm** on the mesh graph

# Graph Laplacian

Given a **mesh graph**  $G = (V, E)$ , consider this condition on vertex  $v_i$ :

$$\mathbf{v}_i - \frac{1}{d_i} \sum_{j:(i,j) \in E} \mathbf{v}_j = 0$$

where  $d_i$  is the valence of  $v_i$



# Graph Laplacian

Given a **mesh graph**  $G = (V, E)$ , consider this condition on vertex  $v_i$ :

$$\mathbf{v}_i - \frac{1}{d_i} \sum_{j:(i,j) \in E} \mathbf{v}_j = 0$$

where  $d_i$  is the valence of  $v_i$

The equation is satisfied if  $v_i$  lies in the **barycenter** of its neighbors

# Graph Laplacian

Given a **mesh graph**  $G = (V, E)$ , consider this condition on vertex  $v_i$ :

$$\mathbf{v}_i - \frac{1}{d_i} \sum_{j:(i,j) \in E} \mathbf{v}_j = 0$$

where  $d_i$  is the valence of  $v_i$

The equation is satisfied if  $v_i$  lies in the **barycenter** of its neighbors

In matrix notation, we define the  $n \times n$  matrix  $\mathbf{L}$  as:

$$L_{ij} = \begin{cases} 1 & \text{if } i = j \\ -\frac{1}{d_i} & \text{if } e_{ij} \in E \\ 0 & \text{otherwise} \end{cases}$$

also known as the **graph Laplacian** of  $G$

# Graph Laplacian

Given a **mesh graph**  $G = (V, E)$ , consider this condition on vertex  $v_i$ :

$$\mathbf{v}_i - \frac{1}{d_i} \sum_{j:(i,j) \in E} \mathbf{v}_j = 0$$

where  $d_i$  is the valence of  $v_i$

The equation is satisfied if  $v_i$  lies in the **barycenter** of its neighbors

In matrix notation, we define the  $n \times n$  matrix  $\mathbf{L}$  as:

$$L_{ij} = \begin{cases} 1 & \text{if } i = j \\ -\frac{1}{d_i} & \text{if } e_{ij} \in E \\ 0 & \text{otherwise} \end{cases}$$

also known as the **graph Laplacian** of  $G$ ; we get to:

$$\mathbf{L}\mathbf{V} = \mathbf{0}$$

# Least squares meshes

Is there an embedding satisfying  $\mathbf{L}\mathbf{V} = \mathbf{0}$ ?

- Trivial embedding  $\mathbf{V} = \mathbf{0}$

# Least squares meshes

Is there an embedding satisfying  $\mathbf{L}\mathbf{V} = \mathbf{0}$ ?

- Trivial embedding  $\mathbf{V} = \mathbf{0}$
- One-dimensional subspace  $\text{span}(\mathbf{1})$

# Least squares meshes

Is there an **interesting** embedding satisfying  $\mathbf{L}\mathbf{V} = \mathbf{0}$ ?

- Trivial embedding  $\mathbf{V} = \mathbf{0}$
- One-dimensional subspace  $\text{span}(\mathbf{1})$

# Least squares meshes

Is there an **interesting** embedding satisfying  $\mathbf{L}\mathbf{V} = \mathbf{0}$ ?

- Trivial embedding  $\mathbf{V} = \mathbf{0}$
- One-dimensional subspace  $\text{span}(\mathbf{1})$

Assume  $m \geq 1$  **anchor vertices**  $v_s \in \mathcal{A}$  with known 3D position

# Least squares meshes

Is there an **interesting** embedding satisfying  $\mathbf{L}\mathbf{V} = \mathbf{0}$ ?

- Trivial embedding  $\mathbf{V} = \mathbf{0}$
- One-dimensional subspace  $\text{span}(\mathbf{1})$

Assume  $m \geq 1$  **anchor vertices**  $v_s \in \mathcal{A}$  with known 3D position

Then, consider the linear system:

$$\begin{pmatrix} \mathbf{L} \\ \mathbf{A} \end{pmatrix} \mathbf{V} = \mathbf{b}$$

where

$$a_{ij} = \begin{cases} 1 & \text{if } v_j \in \mathcal{A} \\ 0 & \text{otherwise} \end{cases}, \quad b_k = \begin{cases} (0, 0, 0) & k \leq n \\ \mathbf{v}_{s_{k-n}} & n < k \leq n + m \end{cases}$$



# Least squares meshes

$$\begin{pmatrix} \mathbf{L} \\ \mathbf{A} \end{pmatrix} \mathbf{v} \approx \mathbf{b}$$

# Least squares meshes

$$\min_{\mathbf{v} \in \mathbb{R}^{n \times 3}} \left\| \begin{pmatrix} \mathbf{L} \\ \mathbf{A} \end{pmatrix} \mathbf{v} - \mathbf{b} \right\|_2^2$$

# Least squares meshes


$$\min_{\mathbf{V} \in \mathbb{R}^{n \times 3}} \|\mathbf{L}\mathbf{V}\|_2^2 + \sum_{i=1}^{n+m} \|\mathbf{v}_i - \mathbf{b}_i\|_2^2$$

# Least squares meshes

$$\min_{\mathbf{V} \in \mathbb{R}^{n \times 3}} \|\mathbf{L}\mathbf{V}\|_2^2 + \sum_{v_i \notin \mathcal{A}} \|\mathbf{v}_i\|_2^2 + \sum_{v_i \in \mathcal{A}} \|\mathbf{v}_i - \mathbf{b}_i\|_2^2$$


# Least squares meshes

$$\min_{\mathbf{V} \in \mathbb{R}^{n \times 3}} \|\mathbf{L}\mathbf{V}\|_2^2 + \sum_{v_i \notin \mathcal{A}} \|\mathbf{v}_i\|_2^2 + \sum_{v_i \in \mathcal{A}} \|\mathbf{v}_i - \mathbf{b}_i\|_2^2$$



- Anchor constraints are not satisfied exactly
- At higher resolution, error distributes better among the constraints

# Least squares meshes



Move the anchor positions to do **shape modeling**

Sorkine and Cohen-Or, "Least-squares meshes". Proc. SMI, 2004

## Exercise: Least squares meshes

Implement the example in Figure 4 from:

Sorkine and Cohen-Or, “Least-squares meshes”. Proc. SMI, 2004

# Midterm

Monday 5 November, same room and time as always



# Midterm

Monday 5 November, same room and time as always

## **Evaluation:**

- Take it as a **self-evaluation** – will **not** be used in the final grade!
- May be used as a basis for the **oral** examination

# Midterm

Monday 5 November, same room and time as always

## Evaluation:

- Take it as a **self-evaluation** – will **not** be used in the final grade!
- May be used as a basis for the **oral** examination

## What:

- **Multiple choice** as well as **open** questions
- Covers **everything** we did, today included
- **No coding** questions, expect mathematics