

Metodi Numerici dell'Informatica

Orthogonality and QR decomposition

Emanuele Rodolà
rodola@di.uniroma1.it



SAPIENZA
UNIVERSITÀ DI ROMA

Motivation

Column space

For any $\mathbf{A} \in \mathbb{R}^{n \times m}$, the linear problem:

$$\mathbf{Ax} = \mathbf{b}$$

can be seen as:

“write \mathbf{b} as a linear combination of the columns of \mathbf{A} with coefficients stored in \mathbf{x} ”.

Column space

For any $\mathbf{A} \in \mathbb{R}^{n \times m}$, the linear problem:

$$\mathbf{Ax} = \mathbf{b}$$

can be seen as:

“write \mathbf{b} as a linear combination of the columns of \mathbf{A} with coefficients stored in \mathbf{x} ”.

$$\mathbf{Ax} = \begin{pmatrix} | & | & \cdots \\ \mathbf{a}_1 & \mathbf{a}_2 & \cdots \\ | & | & \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \end{pmatrix} = x_1 \begin{pmatrix} | \\ \mathbf{a}_1 \\ | \end{pmatrix} + x_2 \begin{pmatrix} | \\ \mathbf{a}_2 \\ | \end{pmatrix} + \cdots$$

Numerical instability

Consider the following situation:


$$\mathbf{A} = \begin{pmatrix} | & | & \cdots \\ \mathbf{a}_1 & \mathbf{a}_2 & \cdots \\ | & | & \cdots \end{pmatrix} = \begin{pmatrix} 0 & 0.0001 & \cdots \\ \vdots & \vdots & \cdots \\ 1 & 1.0001 & \cdots \end{pmatrix}$$

Numerical instability

Consider the following situation:

$$\mathbf{A} = \begin{pmatrix} | & | & \cdots \\ \mathbf{a}_1 & \mathbf{a}_2 & \cdots \\ | & | & \cdots \end{pmatrix} = \begin{pmatrix} 0 & 0.0001 & \cdots \\ \vdots & \vdots & \cdots \\ 1 & 1.0001 & \cdots \end{pmatrix}$$

The columns are almost **linearly dependent**:



This problem is **poorly conditioned**, since we can write either:

$$\mathbf{b} \approx x_1 \mathbf{a}_1 + x_2 \mathbf{a}_2$$

$$\mathbf{b} \approx x_2 \mathbf{a}_1 + x_1 \mathbf{a}_2$$

Column space

$$\mathbf{A}\mathbf{x} = \begin{pmatrix} | & | & \cdots \\ \mathbf{a}_1 & \mathbf{a}_2 & \cdots \\ | & | & \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \end{pmatrix} = x_1 \begin{pmatrix} | \\ \mathbf{a}_1 \\ | \end{pmatrix} + x_2 \begin{pmatrix} | \\ \mathbf{a}_2 \\ | \end{pmatrix} + \cdots$$

- A solution might not exist
- The solution might be unique

Column space

$$\mathbf{A}\mathbf{x} = \begin{pmatrix} | & | & \cdots \\ \mathbf{a}_1 & \mathbf{a}_2 & \cdots \\ | & | & \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \end{pmatrix} = x_1 \begin{pmatrix} | \\ \mathbf{a}_1 \\ | \end{pmatrix} + x_2 \begin{pmatrix} | \\ \mathbf{a}_2 \\ | \end{pmatrix} + \cdots$$

- A solution might not exist
- The solution might be unique

It ultimately depends on the structure of the **column space** of \mathbf{A} , which we define as $\text{col } \mathbf{A} = \text{span}(\mathbf{a}_1, \mathbf{a}_2, \dots)$.

Column space

$$\mathbf{A}\mathbf{x} = \begin{pmatrix} | & | & \cdots \\ \mathbf{a}_1 & \mathbf{a}_2 & \cdots \\ | & | & \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \end{pmatrix} = x_1 \begin{pmatrix} | \\ \mathbf{a}_1 \\ | \end{pmatrix} + x_2 \begin{pmatrix} | \\ \mathbf{a}_2 \\ | \end{pmatrix} + \cdots$$

- A solution might not exist
- The solution might be unique

It ultimately depends on the structure of the **column space** of \mathbf{A} , which we define as $\text{col } \mathbf{A} = \text{span}(\mathbf{a}_1, \mathbf{a}_2, \dots)$.

Can we find a simpler **representation** for the column space?

Column space

$$\mathbf{A}\mathbf{x} = \begin{pmatrix} | & | & \cdots \\ \mathbf{a}_1 & \mathbf{a}_2 & \cdots \\ | & | & \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \end{pmatrix} = x_1 \begin{pmatrix} | \\ \mathbf{a}_1 \\ | \end{pmatrix} + x_2 \begin{pmatrix} | \\ \mathbf{a}_2 \\ | \end{pmatrix} + \cdots$$

- A solution might not exist
- The solution might be unique

It ultimately depends on the structure of the **column space** of \mathbf{A} , which we define as $\text{col } \mathbf{A} = \text{span}(\mathbf{a}_1, \mathbf{a}_2, \dots)$.


Can we find a simpler **representation** for the column space?

Can we deal with **numerical instability** explicitly?

Orthogonality


Projections

Consider two vectors \vec{a} and \vec{b} . Which multiple of \vec{a} is closest to \vec{b} ?



Projections

Consider two vectors \mathbf{a} and \mathbf{b} . Which multiple of \mathbf{a} is closest to \mathbf{b} ?




This can be phrased as a **projection** problem:

$$\min_c \|c \mathbf{a} - \mathbf{b}\|_2^2$$

Projections

Consider two vectors \mathbf{a} and \mathbf{b} . Which multiple of \mathbf{a} is closest to \mathbf{b} ?




This can be phrased as a **projection** problem:

$$\min_c \|c \mathbf{a} - \mathbf{b}\|_2^2$$

The optimal c^* gives us the projection $\text{proj}_{\mathbf{a}} \mathbf{b} \equiv c^* \mathbf{a}$.

Projections

Consider two vectors \mathbf{a} and \mathbf{b} . Which multiple of \mathbf{a} is closest to \mathbf{b} ?




This can be phrased as a **projection** problem:

$$\min_c \|c \mathbf{a} - \mathbf{b}\|_2^2$$

The optimal c^* gives us the projection $\text{proj}_{\mathbf{a}} \mathbf{b} \equiv c^* \mathbf{a}$.


Further, the **complement** $\mathbf{b} - \text{proj}_{\mathbf{a}} \mathbf{b}$ is orthogonal to $\text{proj}_{\mathbf{a}} \mathbf{b}$.

Projections



The projection problem is a least-squares problem where $\mathbf{a} \mathbf{c} \approx \mathbf{b}$.

Projections




The projection problem is a least-squares problem where $\mathbf{a}c \approx \mathbf{b}$.

We have the normal equations $\mathbf{a}^\top \mathbf{a} c = \mathbf{a}^\top \mathbf{b}$, hence

$$c = \frac{\mathbf{a}^\top \mathbf{b}}{\mathbf{a}^\top \mathbf{a}}$$

Projections




The projection problem is a least-squares problem where $\mathbf{a}c \approx \mathbf{b}$.

We have the normal equations $\mathbf{a}^\top \mathbf{a}c = \mathbf{a}^\top \mathbf{b}$, hence

$$c = \frac{\mathbf{a}^\top \mathbf{b}}{\|\mathbf{a}\|_2^2}$$

Projections



The projection problem is a least-squares problem where $\mathbf{a}c \approx \mathbf{b}$.

We have the normal equations $\mathbf{a}^\top \mathbf{a}c = \mathbf{a}^\top \mathbf{b}$, hence

$$c = \frac{\mathbf{a}^\top \mathbf{b}}{\|\mathbf{a}\|_2^2}$$

Therefore:

$$\text{proj}_{\mathbf{a}} \mathbf{b} = \frac{\mathbf{a}^\top \mathbf{b}}{\|\mathbf{a}\|_2^2} \mathbf{a}$$

Projections

Similarly, we can project \mathbf{b} onto an entire space.

Projections

Similarly, we can project \mathbf{b} onto an entire space.

Assume we are given an **orthonormal** basis $\{\mathbf{a}_1, \dots, \mathbf{a}_k\}$; then, we can project onto its span by solving the problem:

$$\min_{c_1, \dots, c_k} \|c_1 \mathbf{a}_1 + \dots + c_k \mathbf{a}_k - \mathbf{b}\|_2^2$$


Projections

Similarly, we can project \mathbf{b} onto an entire space.

Assume we are given an **orthonormal** basis $\{\mathbf{a}_1, \dots, \mathbf{a}_k\}$; then, we can project onto its span by solving the problem:

$$\min_{c_1, \dots, c_k} \|c_1 \mathbf{a}_1 + \dots + c_k \mathbf{a}_k - \mathbf{b}\|_2^2$$

Example in \mathbb{R}^3 and $k = 2$:




Projections

Similarly, we can project \mathbf{b} onto an entire space.

Assume we are given an **orthonormal** basis $\{\mathbf{a}_1, \dots, \mathbf{a}_k\}$; then, we can project onto its span by solving the problem:

$$\min_{c_1, \dots, c_k} \|c_1 \mathbf{a}_1 + \dots + c_k \mathbf{a}_k - \mathbf{b}\|_2^2$$

Example in \mathbb{R}^3 and $k = 2$:



The projection is: $\text{proj}_{\text{span}(\mathbf{a}_1, \dots, \mathbf{a}_k)} \mathbf{b} = (\mathbf{a}_1^\top \mathbf{b}) \mathbf{a}_1 + \dots + (\mathbf{a}_k^\top \mathbf{b}) \mathbf{a}_k$

Orthogonal matrices

Consider the trivial linear problem:

$$\mathbf{I}\mathbf{x} = \mathbf{b}$$

where \mathbf{I} is the identity matrix. The obvious solution is $\mathbf{x} = \mathbf{b}$.

Orthogonal matrices

Consider the trivial linear problem:

$$\mathbf{I}\mathbf{x} = \mathbf{b}$$

where \mathbf{I} is the identity matrix. The obvious solution is $\mathbf{x} = \mathbf{b}$.

From the past lectures, we also know that the least-squares problem

$$\mathbf{A}\mathbf{x} \approx \mathbf{b}$$

corresponds to a linear problem

$$\mathbf{A}^\top \mathbf{A}\mathbf{x} = \mathbf{A}^\top \mathbf{b}$$

Orthogonal matrices

Consider the trivial linear problem:

$$\mathbf{I}\mathbf{x} = \mathbf{b}$$

where \mathbf{I} is the identity matrix. The obvious solution is $\mathbf{x} = \mathbf{b}$.

From the past lectures, we also know that the least-squares problem

$$\mathbf{A}\mathbf{x} \approx \mathbf{b}$$

corresponds to a linear problem

$$\mathbf{A}^\top \mathbf{A}\mathbf{x} = \mathbf{A}^\top \mathbf{b}$$

In the lucky case where $\mathbf{A}^\top \mathbf{A} = \mathbf{I}$ (i.e. matrix \mathbf{A} is **orthogonal**), we boil down to the trivial case. Further, this implies that $\mathbf{A}^{-1} = \mathbf{A}^\top$.

Let's analyze this more carefully.

Orthogonal matrices

For clarity, let us call \mathbf{Q} the orthogonal matrices.

The product $\mathbf{Q}^\top \mathbf{Q}$ has the structure:

$$\mathbf{Q}^\top \mathbf{Q} = \begin{pmatrix} - & \vec{q}_1^\top & - \\ - & \vec{q}_2^\top & - \\ & \vdots & \\ - & \vec{q}_n^\top & \end{pmatrix} \begin{pmatrix} | & | & & | \\ \vec{q}_1 & \vec{q}_2 & \cdots & \vec{q}_n \\ | & | & & | \end{pmatrix} = \begin{pmatrix} \vec{q}_1 \cdot \vec{q}_1 & \vec{q}_1 \cdot \vec{q}_2 & \cdots & \vec{q}_1 \cdot \vec{q}_n \\ \vec{q}_2 \cdot \vec{q}_1 & \vec{q}_2 \cdot \vec{q}_2 & \cdots & \vec{q}_2 \cdot \vec{q}_n \\ \vdots & \vdots & \cdots & \vdots \\ \vec{q}_n \cdot \vec{q}_1 & \vec{q}_n \cdot \vec{q}_2 & \cdots & \vec{q}_n \cdot \vec{q}_n \end{pmatrix}$$

Orthogonal matrices

For clarity, let us call \mathbf{Q} the orthogonal matrices.

The product $\mathbf{Q}^\top \mathbf{Q}$ has the structure:

$$\mathbf{Q}^\top \mathbf{Q} = \begin{pmatrix} - & \vec{q}_1^\top & - \\ - & \vec{q}_2^\top & - \\ & \vdots & \\ - & \vec{q}_n^\top & \end{pmatrix} \begin{pmatrix} | & | & \cdots & | \\ \vec{q}_1 & \vec{q}_2 & \cdots & \vec{q}_n \\ | & | & & | \end{pmatrix} = \begin{pmatrix} \vec{q}_1 \cdot \vec{q}_1 & \vec{q}_1 \cdot \vec{q}_2 & \cdots & \vec{q}_1 \cdot \vec{q}_n \\ \vec{q}_2 \cdot \vec{q}_1 & \vec{q}_2 \cdot \vec{q}_2 & \cdots & \vec{q}_2 \cdot \vec{q}_n \\ \vdots & \vdots & \cdots & \vdots \\ \vec{q}_n \cdot \vec{q}_1 & \vec{q}_n \cdot \vec{q}_2 & \cdots & \vec{q}_n \cdot \vec{q}_n \end{pmatrix}$$

Then, $\mathbf{Q}^\top \mathbf{Q} = \mathbf{I}$ means that the columns of \mathbf{Q} have **unit norm** and they are **orthogonal** to one another.

Orthogonal matrices

For clarity, let us call \mathbf{Q} the orthogonal matrices.

The product $\mathbf{Q}^\top \mathbf{Q}$ has the structure:

$$\mathbf{Q}^\top \mathbf{Q} = \begin{pmatrix} - & \vec{q}_1^\top & - \\ - & \vec{q}_2^\top & - \\ & \vdots & \\ - & \vec{q}_n^\top & \end{pmatrix} \begin{pmatrix} | & | & \cdots & | \\ \vec{q}_1 & \vec{q}_2 & & \vec{q}_n \\ | & | & & | \end{pmatrix} = \begin{pmatrix} \vec{q}_1 \cdot \vec{q}_1 & \vec{q}_1 \cdot \vec{q}_2 & \cdots & \vec{q}_1 \cdot \vec{q}_n \\ \vec{q}_2 \cdot \vec{q}_1 & \vec{q}_2 \cdot \vec{q}_2 & \cdots & \vec{q}_2 \cdot \vec{q}_n \\ \vdots & \vdots & \cdots & \vdots \\ \vec{q}_n \cdot \vec{q}_1 & \vec{q}_n \cdot \vec{q}_2 & \cdots & \vec{q}_n \cdot \vec{q}_n \end{pmatrix}$$

Then, $\mathbf{Q}^\top \mathbf{Q} = \mathbf{I}$ means that the columns of \mathbf{Q} have **unit norm** and they are **orthogonal** to one another.

The columns of \mathbf{Q} form an **orthonormal basis** for the column space.

Orthogonal matrices

For clarity, let us call \mathbf{Q} the orthogonal matrices.

The product $\mathbf{Q}^\top \mathbf{Q}$ has the structure:

$$\mathbf{Q}^\top \mathbf{Q} = \begin{pmatrix} - & \vec{q}_1^\top & - \\ - & \vec{q}_2^\top & - \\ & \vdots & \\ - & \vec{q}_n^\top & \end{pmatrix} \begin{pmatrix} | & | & \cdots & | \\ \vec{q}_1 & \vec{q}_2 & \cdots & \vec{q}_n \\ | & | & & | \end{pmatrix} = \begin{pmatrix} \vec{q}_1 \cdot \vec{q}_1 & \vec{q}_1 \cdot \vec{q}_2 & \cdots & \vec{q}_1 \cdot \vec{q}_n \\ \vec{q}_2 \cdot \vec{q}_1 & \vec{q}_2 \cdot \vec{q}_2 & \cdots & \vec{q}_2 \cdot \vec{q}_n \\ \vdots & \vdots & \cdots & \vdots \\ \vec{q}_n \cdot \vec{q}_1 & \vec{q}_n \cdot \vec{q}_2 & \cdots & \vec{q}_n \cdot \vec{q}_n \end{pmatrix}$$

Then, $\mathbf{Q}^\top \mathbf{Q} = \mathbf{I}$ means that the columns of \mathbf{Q} have **unit norm** and they are **orthogonal** to one another.

The columns of \mathbf{Q} form an **orthonormal basis** for the column space.

Examples:

- The **standard basis** of indicator vectors is an orthonormal basis

Orthogonal matrices

For clarity, let us call \mathbf{Q} the orthogonal matrices.

The product $\mathbf{Q}^\top \mathbf{Q}$ has the structure:

$$\mathbf{Q}^\top \mathbf{Q} = \begin{pmatrix} - & \vec{q}_1^\top & - \\ - & \vec{q}_2^\top & - \\ & \vdots & \\ - & \vec{q}_n^\top & \end{pmatrix} \begin{pmatrix} | & | & & | \\ \vec{q}_1 & \vec{q}_2 & \cdots & \vec{q}_n \\ | & | & & | \end{pmatrix} = \begin{pmatrix} \vec{q}_1 \cdot \vec{q}_1 & \vec{q}_1 \cdot \vec{q}_2 & \cdots & \vec{q}_1 \cdot \vec{q}_n \\ \vec{q}_2 \cdot \vec{q}_1 & \vec{q}_2 \cdot \vec{q}_2 & \cdots & \vec{q}_2 \cdot \vec{q}_n \\ \vdots & \vdots & \cdots & \vdots \\ \vec{q}_n \cdot \vec{q}_1 & \vec{q}_n \cdot \vec{q}_2 & \cdots & \vec{q}_n \cdot \vec{q}_n \end{pmatrix}$$

Then, $\mathbf{Q}^\top \mathbf{Q} = \mathbf{I}$ means that the columns of \mathbf{Q} have **unit norm** and they are **orthogonal** to one another.

The columns of \mathbf{Q} form an **orthonormal basis** for the column space.

Examples:

- The **standard basis** of indicator vectors is an orthonormal basis
- The identity matrix \mathbf{I} is orthogonal

Orthogonal matrices

For clarity, let us call \mathbf{Q} the orthogonal matrices.

The product $\mathbf{Q}^\top \mathbf{Q}$ has the structure:

$$\mathbf{Q}^\top \mathbf{Q} = \begin{pmatrix} - & \vec{q}_1^\top & - \\ - & \vec{q}_2^\top & - \\ & \vdots & \\ - & \vec{q}_n^\top & \end{pmatrix} \begin{pmatrix} | & | & \cdots & | \\ \vec{q}_1 & \vec{q}_2 & \cdots & \vec{q}_n \\ | & | & & | \end{pmatrix} = \begin{pmatrix} \vec{q}_1 \cdot \vec{q}_1 & \vec{q}_1 \cdot \vec{q}_2 & \cdots & \vec{q}_1 \cdot \vec{q}_n \\ \vec{q}_2 \cdot \vec{q}_1 & \vec{q}_2 \cdot \vec{q}_2 & \cdots & \vec{q}_2 \cdot \vec{q}_n \\ \vdots & \vdots & \cdots & \vdots \\ \vec{q}_n \cdot \vec{q}_1 & \vec{q}_n \cdot \vec{q}_2 & \cdots & \vec{q}_n \cdot \vec{q}_n \end{pmatrix}$$

Then, $\mathbf{Q}^\top \mathbf{Q} = \mathbf{I}$ means that the columns of \mathbf{Q} have **unit norm** and they are **orthogonal** to one another.

The columns of \mathbf{Q} form an **orthonormal basis** for the column space.

Examples:

- The **standard basis** of indicator vectors is an orthonormal basis
- The identity matrix \mathbf{I} is orthogonal
- Any **permutation matrix** is orthogonal

Isometries

Orthogonal matrices **preserve lengths**:

$$\|Q\mathbf{x}\|_2^2$$

Isometries

Orthogonal matrices **preserve lengths**:

$$\|\mathbf{Q}\mathbf{x}\|_2^2 = \mathbf{x}^\top \mathbf{Q}^\top \mathbf{Q} \mathbf{x}$$

Isometries

Orthogonal matrices **preserve lengths**:

$$\|\mathbf{Q}\mathbf{x}\|_2^2 = \mathbf{x}^\top \mathbf{Q}^\top \mathbf{Q} \mathbf{x} = \mathbf{x}^\top \mathbf{I} \mathbf{x}$$

Isometries

Orthogonal matrices **preserve lengths**:

$$\|\mathbf{Q}\mathbf{x}\|_2^2 = \mathbf{x}^\top \mathbf{Q}^\top \mathbf{Q} \mathbf{x} = \mathbf{x}^\top \mathbf{I} \mathbf{x} = \mathbf{x}^\top \mathbf{x}$$

Isometries

Orthogonal matrices **preserve lengths**:

$$\|\mathbf{Q}\mathbf{x}\|_2^2 = \mathbf{x}^\top \mathbf{Q}^\top \mathbf{Q} \mathbf{x} = \mathbf{x}^\top \mathbf{I} \mathbf{x} = \mathbf{x}^\top \mathbf{x} = \|\mathbf{x}\|_2^2$$

Isometries

Orthogonal matrices **preserve lengths**:

$$\|\mathbf{Q}\mathbf{x}\|_2^2 = \mathbf{x}^\top \mathbf{Q}^\top \mathbf{Q} \mathbf{x} = \mathbf{x}^\top \mathbf{I} \mathbf{x} = \mathbf{x}^\top \mathbf{x} = \|\mathbf{x}\|_2^2$$

Orthogonal matrices also **preserve angles** (i.e. inner products):

$$\langle \mathbf{Q}\mathbf{x}, \mathbf{Q}\mathbf{y} \rangle = \mathbf{x}^\top \mathbf{Q}^\top \mathbf{Q} \mathbf{y} = \mathbf{x}^\top \mathbf{I} \mathbf{y} = \mathbf{x}^\top \mathbf{y}$$

Isometries

Orthogonal matrices **preserve lengths**:

$$\|\mathbf{Q}\mathbf{x}\|_2^2 = \mathbf{x}^\top \mathbf{Q}^\top \mathbf{Q} \mathbf{x} = \mathbf{x}^\top \mathbf{I} \mathbf{x} = \mathbf{x}^\top \mathbf{x} = \|\mathbf{x}\|_2^2$$

Orthogonal matrices also **preserve angles** (i.e. inner products):

$$\langle \mathbf{Q}\mathbf{x}, \mathbf{Q}\mathbf{y} \rangle = \mathbf{x}^\top \mathbf{Q}^\top \mathbf{Q} \mathbf{y} = \mathbf{x}^\top \mathbf{I} \mathbf{y} = \mathbf{x}^\top \mathbf{y}$$

By these properties, the map $\mathbf{x} \mapsto \mathbf{Q}\mathbf{x}$ is an **isometry** of \mathbb{R}^n .

Isometries


Orthogonal matrices **preserve lengths**:

$$\|Q\mathbf{x}\|_2^2 = \mathbf{x}^\top Q^\top Q \mathbf{x} = \mathbf{x}^\top \mathbf{I} \mathbf{x} = \mathbf{x}^\top \mathbf{x} = \|\mathbf{x}\|_2^2$$

Orthogonal matrices also **preserve angles** (i.e. inner products):

$$\langle Q\mathbf{x}, Q\mathbf{y} \rangle = \mathbf{x}^\top Q^\top Q \mathbf{y} = \mathbf{x}^\top \mathbf{I} \mathbf{y} = \mathbf{x}^\top \mathbf{y}$$

By these properties, the map $\mathbf{x} \mapsto Q\mathbf{x}$ is an **isometry** of \mathbb{R}^n .



Non-orthogonal matrices

When solving $\mathbf{Ax} = \mathbf{b}$, most of the times \mathbf{A} is **not** orthogonal.

However, orthogonal matrices can still help us to simplify the problem.

Non-orthogonal matrices

When solving $\mathbf{Ax} = \mathbf{b}$, most of the times \mathbf{A} is **not** orthogonal.

However, orthogonal matrices can still help us to simplify the problem.

General idea Rewrite \mathbf{A} as a product of two matrices, so that $\mathbf{A}^\top \mathbf{Ax} = \mathbf{A}^\top \mathbf{b}$ becomes easier to solve.

Non-orthogonal matrices

When solving $\mathbf{Ax} = \mathbf{b}$, most of the times \mathbf{A} is **not** orthogonal.

However, orthogonal matrices can still help us to simplify the problem.

General idea Rewrite \mathbf{A} as a product of two matrices, so that $\mathbf{A}^\top \mathbf{Ax} = \mathbf{A}^\top \mathbf{b}$ becomes easier to solve.

In particular, suppose we can write:

$$\mathbf{A} = \mathbf{QR}$$

where \mathbf{Q} is **orthogonal** and \mathbf{R} is **invertible**.

Non-orthogonal matrices

When solving $\mathbf{Ax} = \mathbf{b}$, most of the times \mathbf{A} is **not** orthogonal.

However, orthogonal matrices can still help us to simplify the problem.

General idea Rewrite \mathbf{A} as a product of two matrices, so that $\mathbf{A}^\top \mathbf{Ax} = \mathbf{A}^\top \mathbf{b}$ becomes easier to solve.

In particular, suppose we can write:

$$\mathbf{A} = \mathbf{QR}$$

where \mathbf{Q} is **orthogonal** and \mathbf{R} is **invertible**.

Then, the operation

$$\mathbf{AR}^{-1} = \mathbf{Q}$$

can be seen as transforming the columns of \mathbf{A} to make them orthogonal.

QR decomposition

QR decomposition

It turns out that **any** matrix \mathbf{A} admits the factorization:

$$\mathbf{A} = \mathbf{Q}\mathbf{R}$$

which means that we can always “orthogonalize” the columns of any \mathbf{A} .

QR decomposition

It turns out that **any** matrix \mathbf{A} admits the factorization:

$$\mathbf{A} = \mathbf{Q}\mathbf{R}$$

which means that we can always “orthogonalize” the columns of any \mathbf{A} .

Further, it can be shown that

$$\text{col } \mathbf{A}\mathbf{R}^{-1} = \text{col } \mathbf{A}$$

QR decomposition

It turns out that **any** matrix **A** admits the factorization:

$$\mathbf{A} = \mathbf{Q}\mathbf{R}$$

which means that we can always “orthogonalize” the columns of any **A**.

Further, it can be shown that

$$\text{col } \mathbf{A}\mathbf{R}^{-1} = \text{col } \mathbf{A}$$

i.e., orthogonalization does not change the column space of **A**.

QR decomposition

It turns out that **any** matrix \mathbf{A} admits the factorization:

$$\mathbf{A} = \mathbf{Q}\mathbf{R}$$

which means that we can always “orthogonalize” the columns of any \mathbf{A} .

Further, it can be shown that

$$\text{col } \mathbf{A}\mathbf{R}^{-1} = \text{col } \mathbf{A}$$

i.e., orthogonalization does not change the column space of \mathbf{A} .

Since $\mathbf{Q} = \mathbf{A}\mathbf{R}^{-1}$, then

$$\text{col } \mathbf{Q} = \text{col } \mathbf{A}$$

means that the columns of \mathbf{Q} are an orthonormal basis for $\text{col } \mathbf{A}$.

Better conditioning

How does this help us?

Recall the “almost parallel” numerical issue we encountered before:

$$\mathbf{A} = \begin{pmatrix} | & | & \cdots \\ \mathbf{a}_1 & \mathbf{a}_2 & \cdots \\ | & | & \cdots \end{pmatrix} = \begin{pmatrix} 0 & 0.0001 & \cdots \\ \vdots & \vdots & \cdots \\ 1 & 1.0001 & \cdots \end{pmatrix}$$


Better conditioning

How does this help us?

Recall the “almost parallel” numerical issue we encountered before:

$$\mathbf{A} = \left(\begin{array}{c|c|c} | & | & \\ \mathbf{a}_1 & \mathbf{a}_2 & \cdots \\ | & | & \end{array} \right) = \begin{pmatrix} 0 & 0.0001 & \\ \vdots & \vdots & \cdots \\ 1 & 1.0001 & \end{pmatrix}$$

With orthogonalization, we make the problem **better conditioned** by simply applying invertible transformations to the columns of \mathbf{A} .



QR for least squares

Using QR factorization, the normal equations:

$$\mathbf{A}^\top \mathbf{A} \mathbf{x} = \mathbf{A}^\top \mathbf{b}$$

become:

$$(\mathbf{QR})^\top \mathbf{QR} \mathbf{x} = (\mathbf{QR})^\top \mathbf{b}$$

QR for least squares

Using QR factorization, the normal equations:

$$\mathbf{A}^\top \mathbf{A} \mathbf{x} = \mathbf{A}^\top \mathbf{b}$$

become:

$$\mathbf{R}^\top \mathbf{Q}^\top \mathbf{Q} \mathbf{R} \mathbf{x} = \mathbf{R}^\top \mathbf{Q}^\top \mathbf{b}$$

QR for least squares

Using QR factorization, the normal equations:

$$\mathbf{A}^\top \mathbf{A} \mathbf{x} = \mathbf{A}^\top \mathbf{b}$$

become:

$$\mathbf{R}^\top \mathbf{I} \mathbf{R} \mathbf{x} = \mathbf{R}^\top \mathbf{Q}^\top \mathbf{b}$$

QR for least squares

Using QR factorization, the normal equations:

$$\mathbf{A}^\top \mathbf{A} \mathbf{x} = \mathbf{A}^\top \mathbf{b}$$

become:

$$\mathbf{R}^\top \mathbf{R} \mathbf{x} = \mathbf{R}^\top \mathbf{Q}^\top \mathbf{b}$$

QR for least squares

Using QR factorization, the normal equations:

$$\mathbf{A}^\top \mathbf{A} \mathbf{x} = \mathbf{A}^\top \mathbf{b}$$

become:

$$\mathbf{R}^\top \mathbf{R} \mathbf{x} = \mathbf{R}^\top \mathbf{Q}^\top \mathbf{b}$$

And since \mathbf{R} is invertible:

$$\mathbf{R} \mathbf{x} = \mathbf{Q}^\top \mathbf{b}$$

QR for least squares

Using QR factorization, the normal equations:

$$\mathbf{A}^\top \mathbf{A} \mathbf{x} = \mathbf{A}^\top \mathbf{b}$$

become:

$$\mathbf{R}^\top \mathbf{R} \mathbf{x} = \mathbf{R}^\top \mathbf{Q}^\top \mathbf{b}$$

And since \mathbf{R} is invertible:


$$\mathbf{R} \mathbf{x} = \mathbf{Q}^\top \mathbf{b}$$

One additional property of matrix \mathbf{R} in QR decomposition, is that it is **upper triangular**.


$$U = \begin{bmatrix} u_{1,1} & u_{1,2} & u_{1,3} & \dots & u_{1,n} \\ & u_{2,2} & u_{2,3} & \dots & u_{2,n} \\ & & \ddots & \ddots & \vdots \\ & & & \ddots & u_{n-1,n} \\ 0 & & & & u_{n,n} \end{bmatrix}$$

This makes the system immediate to solve by **back-substitution**.


Gram-Schmidt algorithm



Gram-Schmidt algorithm




input




rescale to
unit norm


Gram-Schmidt algorithm



input




rescale to
unit norm




project

Gram-Schmidt algorithm




Gram-Schmidt algorithm



Observe the following:

- The orthogonalized vectors span the same space of $\mathbf{v}_1, \mathbf{v}_2$


Gram-Schmidt algorithm



Observe the following:

- The orthogonalized vectors span the same space of $\mathbf{v}_1, \mathbf{v}_2$
- We can introduce more input vectors one at a time, and orthogonalize them with respect to the previous ones


Gram-Schmidt algorithm



Observe the following:

- The orthogonalized vectors span the same space of $\mathbf{v}_1, \mathbf{v}_2$
- We can introduce more input vectors one at a time, and orthogonalize them with respect to the previous ones
- If the input vectors are columns of \mathbf{A} , the orthogonalized vectors are the columns of \mathbf{Q} .


Gram-Schmidt algorithm



Observe the following:

- The orthogonalized vectors span the same space of $\mathbf{v}_1, \mathbf{v}_2$
- We can introduce more input vectors one at a time, and orthogonalize them with respect to the previous ones
- If the input vectors are columns of \mathbf{A} , the orthogonalized vectors are the columns of \mathbf{Q} . Further, we can compute $\mathbf{R} = \mathbf{Q}^\top \mathbf{A}$.

Gram-Schmidt algorithm



function GRAM-SCHMIDT($\vec{v}_1, \vec{v}_2, \dots, \vec{v}_k$)

▷ Computes an orthonormal basis $\hat{a}_1, \dots, \hat{a}_k$ for span $\{\vec{v}_1, \dots, \vec{v}_k\}$

▷ Assumes $\vec{v}_1, \dots, \vec{v}_k$ are linearly independent.

$\hat{a}_1 \leftarrow \vec{v}_1 / \|\vec{v}_1\|_2$

▷ Nothing to project out of the first vector

for $i \leftarrow 2, 3, \dots, k$

$\vec{p} \leftarrow \vec{0}$

▷ Projection of \vec{v}_i onto span $\{\hat{a}_1, \dots, \hat{a}_{i-1}\}$

for $j \leftarrow 1, 2, \dots, i-1$

$\vec{p} \leftarrow \vec{p} + (\vec{v}_i \cdot \hat{a}_j) \hat{a}_j$

▷ Projecting onto orthonormal basis

$\vec{r} \leftarrow \vec{v}_i - \vec{p}$

▷ Residual is orthogonal to current basis

$\hat{a}_i \leftarrow \vec{r} / \|\vec{r}\|_2$

▷ Normalize this residual and add it to the basis

return $\{\hat{a}_1, \dots, \hat{a}_k\}$

Reduced QR

Let us take a closer look at the dimensions:

$$\mathbf{Ax} = \mathbf{b}$$

In general, \mathbf{A} will be tall with size $m \times n$.

Reduced QR

Let us take a closer look at the dimensions:

$$\mathbf{Ax} = \mathbf{b}$$

In general, \mathbf{A} will be tall with size $m \times n$. Then it can be:

$$\underbrace{\mathbf{A}}_{m \times n} = \underbrace{\mathbf{Q}}_{m \times n} \underbrace{\mathbf{R}}_{n \times n}$$

or

$$\underbrace{\mathbf{A}}_{m \times n} = \underbrace{\mathbf{Q}}_{m \times m} \underbrace{\mathbf{R}}_{m \times n}$$

Reduced QR

Let us take a closer look at the dimensions:

$$\mathbf{Ax} = \mathbf{b}$$

In general, \mathbf{A} will be tall with size $m \times n$. Then it can be:

$$\underbrace{\mathbf{A}}_{m \times n} = \underbrace{\mathbf{Q}}_{m \times n} \underbrace{\mathbf{R}}_{n \times n}$$

or

$$\underbrace{\mathbf{A}}_{m \times n} = \underbrace{\mathbf{Q}}_{m \times m} \underbrace{\mathbf{R}}_{m \times n}$$

- Gram-Schmidt can be modified to output either case

Reduced QR

Let us take a closer look at the dimensions:

$$\mathbf{Ax} = \mathbf{b}$$

In general, \mathbf{A} will be tall with size $m \times n$. Then it can be:

$$\underbrace{\mathbf{A}}_{m \times n} = \underbrace{\mathbf{Q}}_{m \times n} \underbrace{\mathbf{R}}_{n \times n}$$

or

$$\underbrace{\mathbf{A}}_{m \times n} = \underbrace{\mathbf{Q}}_{m \times m} \underbrace{\mathbf{R}}_{m \times n}$$

- Gram-Schmidt can be modified to output either case
- \mathbf{A} has $k \leq n$ linearly independent columns...

Reduced QR

Let us take a closer look at the dimensions:

$$\mathbf{A}\mathbf{x} = \mathbf{b}$$

In general, \mathbf{A} will be tall with size $m \times n$. Then it can be:

$$\underbrace{\mathbf{A}}_{m \times n} = \underbrace{\mathbf{Q}}_{m \times n} \underbrace{\mathbf{R}}_{n \times n}$$

or

$$\underbrace{\mathbf{A}}_{m \times n} = \underbrace{\mathbf{Q}}_{m \times m} \underbrace{\mathbf{R}}_{m \times n}$$

- Gram-Schmidt can be modified to output either case
- \mathbf{A} has $k \leq n$ linearly independent columns...
- ...therefore, we can discard the last $n - k$ columns of \mathbf{Q}

Suggested reading

Read Sections 5.1 – 5.4 of the book:

J. Solomon, “Numerical Algorithms”