

Metodi Numerici dell'Informatica

Spectral decomposition

Emanuele Rodolà
rodola@di.uniroma1.it




SAPIENZA
UNIVERSITÀ DI ROMA

Motivation

Principal axis


Consider the two-dimensional data in this plot:




(a) Input data

Principal axis

Consider the two-dimensional data in this plot:



(a) Input data



(b) Principal axis


Q: Find the vector \mathbf{v} such that each data point \mathbf{x}_i can be written as

$$\mathbf{x}_i = c_i \mathbf{v}$$


where each \mathbf{x}_i has its own c_i

Principal axis


Consider the two-dimensional data in this plot:



(a) Input data



(b) Principal axis



(c) Projection error


Q: Find the vector \mathbf{v} such that each data point \mathbf{x}_i can be written as

$$\mathbf{x}_i = c_i \mathbf{v}$$


where each \mathbf{x}_i has its own c_i

Principal axis


Consider the two-dimensional data in this plot:



(a) Input data



(b) Principal axis



(c) Projection error

$$\begin{aligned} \min_{\mathbf{v}} \quad & \sum_i \|\mathbf{x}_i - \text{proj}_{\mathbf{v}} \mathbf{x}_i\|_2^2 \\ \text{s.t.} \quad & \|\mathbf{v}\|_2 = 1 \end{aligned}$$

Principal axis

$$\begin{aligned} \min_{\mathbf{v}} \quad & \sum_i \|\mathbf{x}_i - \text{proj}_{\mathbf{v}} \mathbf{x}_i\|_2^2 \\ \text{s.t.} \quad & \|\mathbf{v}\|_2 = 1 \end{aligned}$$

Principal axis

$$\begin{aligned} \min_{\mathbf{v}} \quad & \sum_i \left\| \mathbf{x}_i - \frac{\mathbf{x}_i^\top \mathbf{v}}{\mathbf{v}^\top \mathbf{v}} \mathbf{v} \right\|_2^2 \\ \text{s.t.} \quad & \|\mathbf{v}\|_2 = 1 \end{aligned}$$

Principal axis

$$\begin{aligned} \min_{\mathbf{v}} \quad & \sum_i \|\mathbf{x}_i - (\mathbf{x}_i^\top \mathbf{v}) \mathbf{v}\|_2^2 \\ \text{s.t.} \quad & \|\mathbf{v}\|_2 = 1 \end{aligned}$$

Principal axis

$$\begin{aligned} \min_{\mathbf{v}} \quad & \sum_i (\mathbf{x}_i - (\mathbf{x}_i^\top \mathbf{v}) \mathbf{v})^\top (\mathbf{x}_i - (\mathbf{x}_i^\top \mathbf{v}) \mathbf{v}) \\ \text{s.t.} \quad & \|\mathbf{v}\|_2 = 1 \end{aligned}$$

Principal axis

$$\begin{aligned} \min_{\mathbf{v}} \quad & \sum_i (\|\mathbf{x}_i\|_2^2 - 2(\mathbf{x}_i^\top \mathbf{v})(\mathbf{x}_i^\top \mathbf{v}) + (\mathbf{x}_i^\top \mathbf{v})^2 \|\mathbf{v}\|_2^2) \\ \text{s.t.} \quad & \|\mathbf{v}\|_2 = 1 \end{aligned}$$

Principal axis

$$\begin{aligned} \min_{\mathbf{v}} \quad & \sum_i (\|\mathbf{x}_i\|_2^2 - 2(\mathbf{x}_i^\top \mathbf{v})^2 + (\mathbf{x}_i^\top \mathbf{v})^2 \|\mathbf{v}\|_2^2) \\ \text{s.t.} \quad & \|\mathbf{v}\|_2 = 1 \end{aligned}$$

Principal axis

$$\begin{aligned} \min_{\mathbf{v}} \quad & \sum_i (\|\mathbf{x}_i\|_2^2 - (\mathbf{x}_i^\top \mathbf{v})^2) \\ \text{s.t.} \quad & \|\mathbf{v}\|_2 = 1 \end{aligned}$$

Principal axis

$$\begin{aligned} \min_{\mathbf{v}} \quad & - \sum_i (\mathbf{x}_i^\top \mathbf{v})^2 \\ \text{s.t.} \quad & \|\mathbf{v}\|_2 = 1 \end{aligned}$$

Principal axis

$$\begin{aligned} \min_{\mathbf{v}} \quad & -\|\mathbf{X}^\top \mathbf{v}\|_2^2 \\ \text{s.t.} \quad & \|\mathbf{v}\|_2 = 1 \end{aligned}$$

where matrix \mathbf{X} contains the vectors \mathbf{x}_i as its columns.

Principal axis

$$\begin{aligned} \max_{\mathbf{v}} \quad & \|\mathbf{X}^\top \mathbf{v}\|_2^2 \\ \text{s.t.} \quad & \|\mathbf{v}\|_2 = 1 \end{aligned}$$

where matrix \mathbf{X} contains the vectors \mathbf{x}_i as its columns.

This can also be written as

$$\begin{aligned} \max_{\mathbf{v}} \quad & \mathbf{v}^\top \underbrace{\mathbf{X}\mathbf{X}^\top}_{\text{symmetric}} \mathbf{v} \\ \text{s.t.} \quad & \|\mathbf{v}\|_2 = 1 \end{aligned}$$

Principal axis

$$\begin{aligned} \max_{\mathbf{v}} \quad & \|\mathbf{X}^\top \mathbf{v}\|_2^2 \\ \text{s.t.} \quad & \|\mathbf{v}\|_2 = 1 \end{aligned}$$

where matrix \mathbf{X} contains the vectors \mathbf{x}_i as its columns.

This can also be written as

$$\begin{aligned} \max_{\mathbf{v}} \quad & \mathbf{v}^\top \underbrace{\mathbf{X}\mathbf{X}^\top}_{\text{symmetric}} \mathbf{v} \\ \text{s.t.} \quad & \|\mathbf{v}\|_2 = 1 \end{aligned}$$

The global maximizer \mathbf{v}^* of this problem is the **principal component** of the data contained in the matrix \mathbf{X} .

Eigenvectors and eigenvalues

Eigenvalue equation

An **eigenvector** \mathbf{x} of a square matrix \mathbf{A} is any vector satisfying

$$\mathbf{Ax} = \lambda\mathbf{x}$$

for some (possibly complex) number λ that we call **eigenvalue**.

Eigenvalue equation

An **eigenvector** \mathbf{x} of a square matrix \mathbf{A} is any vector satisfying

$$\mathbf{A}\mathbf{x} = \lambda\mathbf{x}$$

for some (possibly complex) number λ that we call **eigenvalue**.

A few basic facts:

The **scale** of an eigenvector is not important. In particular:

$$\mathbf{A}\mathbf{c}\mathbf{x}$$

Eigenvalue equation

An **eigenvector** \mathbf{x} of a square matrix \mathbf{A} is any vector satisfying

$$\mathbf{A}\mathbf{x} = \lambda\mathbf{x}$$

for some (possibly complex) number λ that we call **eigenvalue**.

A few basic facts:

The **scale** of an eigenvector is not important. In particular:

$$\mathbf{A}c\mathbf{x} = c\mathbf{A}\mathbf{x}$$

Eigenvalue equation

An **eigenvector** \mathbf{x} of a square matrix \mathbf{A} is any vector satisfying

$$\mathbf{A}\mathbf{x} = \lambda\mathbf{x}$$

for some (possibly complex) number λ that we call **eigenvalue**.

A few basic facts:

The **scale** of an eigenvector is not important. In particular:

$$\mathbf{A}c\mathbf{x} = c\mathbf{A}\mathbf{x} = c\lambda\mathbf{x}$$

Eigenvalue equation

An **eigenvector** \mathbf{x} of a square matrix \mathbf{A} is any vector satisfying

$$\mathbf{A}\mathbf{x} = \lambda\mathbf{x}$$

for some (possibly complex) number λ that we call **eigenvalue**.

A few basic facts:

The **scale** of an eigenvector is not important. In particular:

$$\mathbf{A}c\mathbf{x} = c\mathbf{A}\mathbf{x} = c\lambda\mathbf{x} = \lambda c\mathbf{x}$$

Eigenvalue equation

An **eigenvector** \mathbf{x} of a square matrix \mathbf{A} is any vector satisfying

$$\mathbf{A}\mathbf{x} = \lambda\mathbf{x}$$

for some (possibly complex) number λ that we call **eigenvalue**.

A few basic facts:

The **scale** of an eigenvector is not important. In particular:

$$\mathbf{A}c\mathbf{x} = c\mathbf{A}\mathbf{x} = c\lambda\mathbf{x} = \lambda c\mathbf{x}$$

For this reason, we can restrict our search to eigenvectors with $\|\mathbf{x}\|_2 = 1$.

Eigenvalue equation

An **eigenvector** \mathbf{x} of a square matrix \mathbf{A} is any vector satisfying

$$\mathbf{A}\mathbf{x} = \lambda\mathbf{x}$$

for some (possibly complex) number λ that we call **eigenvalue**.

A few basic facts:

The **scale** of an eigenvector is not important. In particular:

$$\mathbf{A}c\mathbf{x} = c\mathbf{A}\mathbf{x} = c\lambda\mathbf{x} = \lambda c\mathbf{x}$$

For this reason, we can restrict our search to eigenvectors with $\|\mathbf{x}\|_2 = 1$.

Clearly, \mathbf{x} and $-\mathbf{x}$ are both eigenvectors with the same eigenvalue.

Similarity

Eigenvectors can be scaled without changing their eigenvalues:

$$\mathbf{A}c\mathbf{x} = \lambda c\mathbf{x}$$

Similarity

Eigenvectors can be scaled without changing their eigenvalues:

$$\mathbf{A}c\mathbf{x} = \lambda c\mathbf{x}$$

Instead of just scaling, consider an invertible linear transformation:

$$\mathbf{A}\mathbf{T}\mathbf{x} = \lambda\mathbf{T}\mathbf{x}$$

which is saying that $\mathbf{T}\mathbf{x}$ is an eigenvector of \mathbf{A} with eigenvalue λ .

Similarity

Eigenvectors can be scaled without changing their eigenvalues:

$$\mathbf{A}c\mathbf{x} = \lambda c\mathbf{x}$$

Instead of just scaling, consider an invertible linear transformation:

$$\mathbf{A}\mathbf{T}\mathbf{x} = \lambda\mathbf{T}\mathbf{x}$$

which is saying that $\mathbf{T}\mathbf{x}$ is an eigenvector of \mathbf{A} with eigenvalue λ .

Is it true that also \mathbf{x} is an eigenvector of \mathbf{A} ? Let's check:

$$\mathbf{A}\mathbf{T}\mathbf{x} = \lambda\mathbf{T}\mathbf{x}$$

Similarity

Eigenvectors can be scaled without changing their eigenvalues:

$$\mathbf{A}c\mathbf{x} = \lambda c\mathbf{x}$$

Instead of just scaling, consider an invertible linear transformation:

$$\mathbf{A}\mathbf{T}\mathbf{x} = \lambda\mathbf{T}\mathbf{x}$$

which is saying that $\mathbf{T}\mathbf{x}$ is an eigenvector of \mathbf{A} with eigenvalue λ .

Is it true that also \mathbf{x} is an eigenvector of \mathbf{A} ? Let's check:

$$\mathbf{T}^{-1}\mathbf{A}\mathbf{T}\mathbf{x} = \lambda\mathbf{x}$$

Similarity

Eigenvectors can be scaled without changing their eigenvalues:

$$\mathbf{A}c\mathbf{x} = \lambda c\mathbf{x}$$

Instead of just scaling, consider an invertible linear transformation:

$$\mathbf{A}\mathbf{T}\mathbf{x} = \lambda\mathbf{T}\mathbf{x}$$

which is saying that $\mathbf{T}\mathbf{x}$ is an eigenvector of \mathbf{A} with eigenvalue λ .

Is it true that also \mathbf{x} is an eigenvector of \mathbf{A} ? Let's check:

$$\mathbf{T}^{-1}\mathbf{A}\mathbf{T}\mathbf{x} = \lambda\mathbf{x}$$

Similarity

Eigenvectors can be scaled without changing their eigenvalues:

$$\mathbf{A}c\mathbf{x} = \lambda c\mathbf{x}$$

Instead of just scaling, consider an invertible **linear transformation**:

$$\mathbf{A}\mathbf{T}\mathbf{x} = \lambda\mathbf{T}\mathbf{x}$$

which is saying that $\mathbf{T}\mathbf{x}$ is an eigenvector of \mathbf{A} with eigenvalue λ .

Is it true that also \mathbf{x} is an eigenvector of \mathbf{A} ? Let's check:

$$\mathbf{T}^{-1}\mathbf{A}\mathbf{T}\mathbf{x} = \lambda\mathbf{x}$$

In other words, \mathbf{x} is an eigenvector of $\mathbf{T}^{-1}\mathbf{A}\mathbf{T}$ with eigenvalue λ .

Similarity

Eigenvectors can be scaled without changing their eigenvalues:

$$\mathbf{A}c\mathbf{x} = \lambda c\mathbf{x}$$

Instead of just scaling, consider an invertible **linear transformation**:

$$\mathbf{A}\mathbf{T}\mathbf{x} = \lambda\mathbf{T}\mathbf{x}$$

which is saying that $\mathbf{T}\mathbf{x}$ is an eigenvector of \mathbf{A} with eigenvalue λ .

Is it true that also \mathbf{x} is an eigenvector of \mathbf{A} ? Let's check:

$$\mathbf{T}^{-1}\mathbf{A}\mathbf{T}\mathbf{x} = \lambda\mathbf{x}$$

In other words, \mathbf{x} is an eigenvector of $\mathbf{T}^{-1}\mathbf{A}\mathbf{T}$ with eigenvalue λ .

We say that $\mathbf{B} = \mathbf{T}^{-1}\mathbf{A}\mathbf{T}$ is a **similarity transformation**. \mathbf{A} and \mathbf{B} have the same eigenvalues.

More basic facts

A few more basic facts that will become useful later:

- Orthogonal matrices

Observe:

$$Q\mathbf{x} = \lambda\mathbf{x}$$

More basic facts

A few more basic facts that will become useful later:

- Orthogonal matrices

Observe:

$$\|Q\mathbf{x}\|_2^2 = \|\lambda\mathbf{x}\|_2^2$$

More basic facts

A few more basic facts that will become useful later:

- Orthogonal matrices

Observe:

$$(\mathbf{Q}\mathbf{x})^\top \mathbf{Q}\mathbf{x} = |\lambda|^2 \|\mathbf{x}\|_2^2$$

More basic facts

A few more basic facts that will become useful later:

- Orthogonal matrices

Observe:

$$\mathbf{x}^\top \mathbf{Q}^\top \mathbf{Q} \mathbf{x} = |\lambda|^2 \|\mathbf{x}\|_2^2$$

More basic facts

A few more basic facts that will become useful later:

- Orthogonal matrices

Observe:

$$\mathbf{x}^\top \mathbf{x} = |\lambda|^2 \|\mathbf{x}\|_2^2$$

More basic facts

A few more basic facts that will become useful later:

- Orthogonal matrices

Observe:

$$\|\mathbf{x}\|_2^2 = |\lambda|^2 \|\mathbf{x}\|_2^2$$

More basic facts

A few more basic facts that will become useful later:

- Orthogonal matrices

Observe:

$$1 = |\lambda|^2$$

More basic facts

A few more basic facts that will become useful later:

- Orthogonal matrices

Observe:

$$1 = |\lambda|$$

More basic facts

A few more basic facts that will become useful later:

- Orthogonal matrices

We obtained:

$$\lambda = \pm 1$$

- Diagonal and upper-triangular matrices

The eigenvalues are the entries along the main diagonal.

More basic facts

A few more basic facts that will become useful later:

- Orthogonal matrices

We obtained:

$$\lambda = \pm 1$$

- Diagonal and upper-triangular matrices

The eigenvalues are the entries along the main diagonal.

- Commuting matrices

Consider two matrices **A** and **B**. One can prove:

$$\mathbf{AB} = \mathbf{BA} \quad \Leftrightarrow \quad \mathbf{A} \text{ and } \mathbf{B} \text{ have the same eigenvectors}$$

Big questions

- How to compute eigenvalues and eigenvectors?

Big questions

- How to compute eigenvalues and eigenvectors?
- What to do with them?

A tentative answer

Why do we care about eigenvectors and eigenvalues?

$$\mathbf{A}\mathbf{x} = \lambda\mathbf{x}$$

A tentative answer

Why do we care about eigenvectors and eigenvalues?

$$\mathbf{A}\mathbf{x} = \lambda\mathbf{x}$$

A first trivial (but quite valid!) observation:

- If \mathbf{x} is an eigenvector, multiplying by \mathbf{A} equals simply scaling \mathbf{x} .

A tentative answer

Why do we care about eigenvectors and eigenvalues?

$$\mathbf{A}\mathbf{x} = \lambda\mathbf{x}$$

A first trivial (but quite valid!) observation:

- If \mathbf{x} is an eigenvector, multiplying by \mathbf{A} equals simply scaling \mathbf{x} .

But what if \mathbf{y} is some arbitrary vector?

Suppose the eigenvectors $\{\mathbf{x}_i\}$ form a basis. Then:

A tentative answer

Why do we care about eigenvectors and eigenvalues?

$$\mathbf{A}\mathbf{x} = \lambda\mathbf{x}$$

A first trivial (but quite valid!) observation:

- If \mathbf{x} is an eigenvector, multiplying by \mathbf{A} equals simply scaling \mathbf{x} .

But what if \mathbf{y} is some arbitrary vector?

Suppose the eigenvectors $\{\mathbf{x}_i\}$ form a basis. Then:

$$\mathbf{y} = \sum_i \alpha_i \mathbf{x}_i$$

A tentative answer

Why do we care about eigenvectors and eigenvalues?

$$\mathbf{A}\mathbf{x} = \lambda\mathbf{x}$$

A first trivial (but quite valid!) observation:

- If \mathbf{x} is an eigenvector, multiplying by \mathbf{A} equals simply scaling \mathbf{x} .

But what if \mathbf{y} is some arbitrary vector?

Suppose the eigenvectors $\{\mathbf{x}_i\}$ form a basis. Then:

$$\mathbf{y} = \sum_i \alpha_i \mathbf{x}_i$$

- If \mathbf{y} is an arbitrary vector, multiplying by \mathbf{A} equals

$$\mathbf{A}\mathbf{y} = \mathbf{A} \sum_i \alpha_i \mathbf{x}_i$$

A tentative answer

Why do we care about eigenvectors and eigenvalues?

$$\mathbf{A}\mathbf{x} = \lambda\mathbf{x}$$

A first trivial (but quite valid!) observation:

- If \mathbf{x} is an eigenvector, multiplying by \mathbf{A} equals simply scaling \mathbf{x} .

But what if \mathbf{y} is some arbitrary vector?

Suppose the eigenvectors $\{\mathbf{x}_i\}$ form a basis. Then:

$$\mathbf{y} = \sum_i \alpha_i \mathbf{x}_i$$

- If \mathbf{y} is an arbitrary vector, multiplying by \mathbf{A} equals

$$\mathbf{A}\mathbf{y} = \sum_i \alpha_i \mathbf{A}\mathbf{x}_i$$

A tentative answer

Why do we care about eigenvectors and eigenvalues?

$$\mathbf{A}\mathbf{x} = \lambda\mathbf{x}$$

A first trivial (but quite valid!) observation:

- If \mathbf{x} is an eigenvector, multiplying by \mathbf{A} equals simply scaling \mathbf{x} .

But what if \mathbf{y} is some arbitrary vector?

Suppose the eigenvectors $\{\mathbf{x}_i\}$ form a basis. Then:

$$\mathbf{y} = \sum_i \alpha_i \mathbf{x}_i$$


- If \mathbf{y} is an arbitrary vector, multiplying by \mathbf{A} equals

$$\mathbf{A}\mathbf{y} = \sum_i \alpha_i \lambda_i \mathbf{x}_i$$


Min-max theorem

Back to our motivation:


$$\begin{aligned} \max_{\mathbf{v}} \mathbf{v}^\top \underbrace{\mathbf{X}\mathbf{X}^\top}_{\text{symmetric}} \mathbf{v} \\ \text{s.t. } \|\mathbf{v}\|_2 = 1 \end{aligned}$$



(a) Input data



(b) Principal axis



(c) Projection error

Min-max theorem

Back to our motivation:

$$\begin{aligned} \max_{\mathbf{v}} \mathbf{v}^\top \mathbf{A} \mathbf{v} \\ \text{s.t. } \|\mathbf{v}\|_2 = 1 \end{aligned}$$

where \mathbf{A} is *symmetric*.

Min-max theorem

Back to our motivation:

$$\begin{aligned} \max_{\mathbf{v}} \mathbf{v}^\top \mathbf{A} \mathbf{v} \\ \text{s.t. } \|\mathbf{v}\|_2 = 1 \end{aligned}$$

where \mathbf{A} is *symmetric*.

Theorem If \mathbf{A} is symmetric, then its maximum eigenvalue is given by $\max_{\mathbf{v}} \frac{\mathbf{v}^\top \mathbf{A} \mathbf{v}}{\|\mathbf{v}\|_2^2}$, and \mathbf{v} is the corresponding eigenvector.

Min-max theorem

Back to our motivation:

$$\begin{aligned} \max_{\mathbf{v}} \quad & \mathbf{v}^\top \mathbf{A} \mathbf{v} \\ \text{s.t.} \quad & \|\mathbf{v}\|_2 = 1 \end{aligned}$$

where \mathbf{A} is [symmetric](#).

Theorem If \mathbf{A} is symmetric, then its maximum eigenvalue is given by $\max_{\mathbf{v}} \frac{\mathbf{v}^\top \mathbf{A} \mathbf{v}}{\|\mathbf{v}\|_2^2}$, and \mathbf{v} is the corresponding eigenvector. More in general:

$$\lambda_{\min} \leq \frac{\mathbf{v}^\top \mathbf{A} \mathbf{v}}{\|\mathbf{v}\|_2^2} \leq \lambda_{\max}$$

Min-max theorem

Back to our motivation:

$$\begin{aligned} \max_{\mathbf{v}} \quad & \mathbf{v}^\top \mathbf{A} \mathbf{v} \\ \text{s.t.} \quad & \|\mathbf{v}\|_2 = 1 \end{aligned}$$

where \mathbf{A} is **symmetric**.

Theorem If \mathbf{A} is symmetric, then its maximum eigenvalue is given by $\max_{\mathbf{v}} \frac{\mathbf{v}^\top \mathbf{A} \mathbf{v}}{\|\mathbf{v}\|_2^2}$, and \mathbf{v} is the corresponding eigenvector. More in general:

$$\lambda_{\min} \leq \frac{\mathbf{v}^\top \mathbf{A} \mathbf{v}}{\|\mathbf{v}\|_2^2} \leq \lambda_{\max}$$

The ratio $\frac{\mathbf{v}^\top \mathbf{A} \mathbf{v}}{\|\mathbf{v}\|_2^2}$ is called **Rayleigh quotient**.

Symmetric matrices

A matrix \mathbf{A} is **symmetric** if $\mathbf{A} = \mathbf{A}^\top$.

Symmetric matrices

A matrix \mathbf{A} is **symmetric** if $\mathbf{A} = \mathbf{A}^\top$.

Consider an eigenvalue-eigenvector pair (λ, \mathbf{x}) , where $\|\mathbf{x}\|_2^2 = 1$.

All eigenvalues of symmetric matrices are **real**:

$$\lambda \mathbf{x}^\top \mathbf{x} = (\lambda \mathbf{x})^\top \mathbf{x}$$

Symmetric matrices

A matrix \mathbf{A} is **symmetric** if $\mathbf{A} = \mathbf{A}^\top$.

Consider an eigenvalue-eigenvector pair (λ, \mathbf{x}) , where $\|\mathbf{x}\|_2^2 = 1$.

All eigenvalues of symmetric matrices are **real**:

$$\begin{aligned}\lambda \mathbf{x}^\top \mathbf{x} &= (\lambda \mathbf{x})^\top \mathbf{x} \\ &= (\mathbf{A} \mathbf{x})^\top \mathbf{x}\end{aligned}$$

Symmetric matrices

A matrix \mathbf{A} is **symmetric** if $\mathbf{A} = \mathbf{A}^\top$.

Consider an eigenvalue-eigenvector pair (λ, \mathbf{x}) , where $\|\mathbf{x}\|_2^2 = 1$.

All eigenvalues of symmetric matrices are **real**:

$$\begin{aligned}\lambda \mathbf{x}^\top \mathbf{x} &= (\lambda \mathbf{x})^\top \mathbf{x} \\ &= (\mathbf{A} \mathbf{x})^\top \mathbf{x} \\ &= \mathbf{x}^\top \mathbf{A}^\top \mathbf{x}\end{aligned}$$

Symmetric matrices

A matrix \mathbf{A} is **symmetric** if $\mathbf{A} = \mathbf{A}^\top$.

Consider an eigenvalue-eigenvector pair (λ, \mathbf{x}) , where $\|\mathbf{x}\|_2^2 = 1$.

All eigenvalues of symmetric matrices are **real**:

$$\begin{aligned}\lambda \mathbf{x}^\top \mathbf{x} &= (\lambda \mathbf{x})^\top \mathbf{x} \\ &= (\mathbf{A} \mathbf{x})^\top \mathbf{x} \\ &= \mathbf{x}^\top \mathbf{A}^\top \mathbf{x} \\ &= \mathbf{x}^\top (\mathbf{A} \mathbf{x})\end{aligned}$$

Symmetric matrices

A matrix \mathbf{A} is **symmetric** if $\mathbf{A} = \mathbf{A}^\top$.

Consider an eigenvalue-eigenvector pair (λ, \mathbf{x}) , where $\|\mathbf{x}\|_2^2 = 1$.

All eigenvalues of symmetric matrices are **real**:

$$\begin{aligned}\lambda \mathbf{x}^\top \mathbf{x} &= (\lambda \mathbf{x})^\top \mathbf{x} \\ &= (\mathbf{A} \mathbf{x})^\top \mathbf{x} \\ &= \mathbf{x}^\top \mathbf{A}^\top \mathbf{x} \\ &= \mathbf{x}^\top (\mathbf{A} \mathbf{x}) \\ &= \overline{(\mathbf{A} \mathbf{x})^\top \mathbf{x}}\end{aligned}$$

Symmetric matrices

A matrix \mathbf{A} is **symmetric** if $\mathbf{A} = \mathbf{A}^\top$.

Consider an eigenvalue-eigenvector pair (λ, \mathbf{x}) , where $\|\mathbf{x}\|_2^2 = 1$.

All eigenvalues of symmetric matrices are **real**:

$$\begin{aligned}\lambda \mathbf{x}^\top \mathbf{x} &= (\lambda \mathbf{x})^\top \mathbf{x} \\ &= (\mathbf{A} \mathbf{x})^\top \mathbf{x} \\ &= \mathbf{x}^\top \mathbf{A}^\top \mathbf{x} \\ &= \mathbf{x}^\top (\mathbf{A} \mathbf{x}) \\ &= \overline{(\mathbf{A} \mathbf{x})^\top \mathbf{x}} \\ &= \overline{(\lambda \mathbf{x})^\top \mathbf{x}}\end{aligned}$$

Symmetric matrices

A matrix \mathbf{A} is **symmetric** if $\mathbf{A} = \mathbf{A}^\top$.

Consider an eigenvalue-eigenvector pair (λ, \mathbf{x}) , where $\|\mathbf{x}\|_2^2 = 1$.

All eigenvalues of symmetric matrices are **real**:

$$\begin{aligned}\lambda \mathbf{x}^\top \mathbf{x} &= (\lambda \mathbf{x})^\top \mathbf{x} \\ &= (\mathbf{A} \mathbf{x})^\top \mathbf{x} \\ &= \mathbf{x}^\top \mathbf{A}^\top \mathbf{x} \\ &= \mathbf{x}^\top (\mathbf{A} \mathbf{x}) \\ &= \overline{(\mathbf{A} \mathbf{x})^\top \mathbf{x}} \\ &= \overline{(\lambda \mathbf{x})^\top \mathbf{x}} \\ &= \bar{\lambda} \mathbf{x}^\top \mathbf{x}\end{aligned}$$

Symmetric matrices

A matrix \mathbf{A} is **symmetric** if $\mathbf{A} = \mathbf{A}^\top$.

Consider an eigenvalue-eigenvector pair (λ, \mathbf{x}) , where $\|\mathbf{x}\|_2^2 = 1$.

All eigenvalues of symmetric matrices are **real**:

$$\begin{aligned}\lambda \mathbf{x}^\top \mathbf{x} &= (\lambda \mathbf{x})^\top \mathbf{x} \\ &= (\mathbf{A} \mathbf{x})^\top \mathbf{x} \\ &= \mathbf{x}^\top \mathbf{A}^\top \mathbf{x} \\ &= \mathbf{x}^\top (\mathbf{A} \mathbf{x}) \\ &= \overline{(\mathbf{A} \mathbf{x})^\top \mathbf{x}} \\ &= \overline{(\lambda \mathbf{x})^\top \mathbf{x}} \\ &= \bar{\lambda} \mathbf{x}^\top \mathbf{x}\end{aligned}$$

Consider distinct eigenvectors $\mathbf{x}_i, \mathbf{x}_j$ with $\lambda_i \neq \lambda_j$:

Symmetric matrices

A matrix \mathbf{A} is **symmetric** if $\mathbf{A} = \mathbf{A}^\top$.

Consider an eigenvalue-eigenvector pair (λ, \mathbf{x}) , where $\|\mathbf{x}\|_2^2 = 1$.

All eigenvalues of symmetric matrices are **real**:

$$\begin{aligned}\lambda \mathbf{x}^\top \mathbf{x} &= (\lambda \mathbf{x})^\top \mathbf{x} \\ &= (\mathbf{A} \mathbf{x})^\top \mathbf{x} \\ &= \mathbf{x}^\top \mathbf{A}^\top \mathbf{x} \\ &= \mathbf{x}^\top (\mathbf{A} \mathbf{x}) \\ &= \overline{(\mathbf{A} \mathbf{x})^\top \mathbf{x}} \\ &= \overline{(\lambda \mathbf{x})^\top \mathbf{x}} \\ &= \bar{\lambda} \mathbf{x}^\top \mathbf{x}\end{aligned}$$

Consider distinct eigenvectors $\mathbf{x}_i, \mathbf{x}_j$ with $\lambda_i \neq \lambda_j$:

$$(\mathbf{A} \mathbf{x}_i)^\top \mathbf{x}_j = \mathbf{x}_i^\top \mathbf{A} \mathbf{x}_j$$

Symmetric matrices

A matrix \mathbf{A} is **symmetric** if $\mathbf{A} = \mathbf{A}^\top$.

Consider an eigenvalue-eigenvector pair (λ, \mathbf{x}) , where $\|\mathbf{x}\|_2^2 = 1$.

All eigenvalues of symmetric matrices are **real**:

$$\begin{aligned}\lambda \mathbf{x}^\top \mathbf{x} &= (\lambda \mathbf{x})^\top \mathbf{x} \\ &= (\mathbf{A} \mathbf{x})^\top \mathbf{x} \\ &= \mathbf{x}^\top \mathbf{A}^\top \mathbf{x} \\ &= \mathbf{x}^\top (\mathbf{A} \mathbf{x}) \\ &= \overline{(\mathbf{A} \mathbf{x})^\top \mathbf{x}} \\ &= \overline{(\lambda \mathbf{x})^\top \mathbf{x}} \\ &= \bar{\lambda} \mathbf{x}^\top \mathbf{x}\end{aligned}$$

Consider distinct eigenvectors $\mathbf{x}_i, \mathbf{x}_j$ with $\lambda_i \neq \lambda_j$:

$$(\lambda_i \mathbf{x}_i)^\top \mathbf{x}_j = \mathbf{x}_i^\top \lambda_j \mathbf{x}_j$$

Symmetric matrices

A matrix \mathbf{A} is **symmetric** if $\mathbf{A} = \mathbf{A}^\top$.

Consider an eigenvalue-eigenvector pair (λ, \mathbf{x}) , where $\|\mathbf{x}\|_2^2 = 1$.

All eigenvalues of symmetric matrices are **real**:

$$\begin{aligned}\lambda \mathbf{x}^\top \mathbf{x} &= (\lambda \mathbf{x})^\top \mathbf{x} \\ &= (\mathbf{A} \mathbf{x})^\top \mathbf{x} \\ &= \mathbf{x}^\top \mathbf{A}^\top \mathbf{x} \\ &= \mathbf{x}^\top (\mathbf{A} \mathbf{x}) \\ &= \overline{(\mathbf{A} \mathbf{x})^\top \mathbf{x}} \\ &= \overline{(\lambda \mathbf{x})^\top \mathbf{x}} \\ &= \bar{\lambda} \mathbf{x}^\top \mathbf{x}\end{aligned}$$

Consider distinct eigenvectors $\mathbf{x}_i, \mathbf{x}_j$ with $\lambda_i \neq \lambda_j$:

$$\lambda_i \mathbf{x}_i^\top \mathbf{x}_j = \lambda_j \mathbf{x}_i^\top \mathbf{x}_j$$

For the equality to hold, it must be $\mathbf{x}_i^\top \mathbf{x}_j = 0$, i.e. \mathbf{x}_i and \mathbf{x}_j are **orthogonal**.

Spectral theorem

The set of eigenvalues $\{\lambda_i\}$ of a matrix \mathbf{A} is called the **spectrum**.

Spectral theorem

The set of eigenvalues $\{\lambda_i\}$ of a matrix \mathbf{A} is called the **spectrum**.

We can write the eigenvalue equation as:

$$\mathbf{A}\mathbf{X} = \mathbf{X}\mathbf{\Lambda}$$

If \mathbf{A} is symmetric, then \mathbf{X} is an **orthogonal** matrix of eigenvectors, and $\mathbf{\Lambda}$ is a **diagonal** matrix of real eigenvalues.

Spectral theorem

The set of eigenvalues $\{\lambda_i\}$ of a matrix \mathbf{A} is called the **spectrum**.

We can write the eigenvalue equation as:

$$\mathbf{X}^\top \mathbf{A} \mathbf{X} = \mathbf{\Lambda}$$

If \mathbf{A} is symmetric, then \mathbf{X} is an **orthogonal** matrix of eigenvectors, and $\mathbf{\Lambda}$ is a **diagonal** matrix of real eigenvalues.

Spectral theorem

The set of eigenvalues $\{\lambda_i\}$ of a matrix \mathbf{A} is called the **spectrum**.

We can write the eigenvalue equation as:

$$\mathbf{X}^\top \mathbf{A} \mathbf{X} = \mathbf{\Lambda}$$

If \mathbf{A} is symmetric, then \mathbf{X} is an **orthogonal** matrix of eigenvectors, and $\mathbf{\Lambda}$ is a **diagonal** matrix of real eigenvalues.

We call it the **spectral decomposition** of \mathbf{A} .

Spectral theorem

The set of eigenvalues $\{\lambda_i\}$ of a matrix \mathbf{A} is called the **spectrum**.

We can write the eigenvalue equation as:

$$\mathbf{X}^\top \mathbf{A} \mathbf{X} = \mathbf{\Lambda}$$

If \mathbf{A} is symmetric, then \mathbf{X} is an **orthogonal** matrix of eigenvectors, and $\mathbf{\Lambda}$ is a **diagonal** matrix of real eigenvalues.

We call it the **spectral decomposition** of \mathbf{A} .

Observe the similarity with our motivational problem.

$$\begin{aligned} \max_{\mathbf{x}} \quad & \mathbf{x}^\top \mathbf{A} \mathbf{x} \\ \text{s.t.} \quad & \|\mathbf{x}\|_2 = 1 \end{aligned}$$

Spectral theorem

The set of eigenvalues $\{\lambda_i\}$ of a matrix \mathbf{A} is called the **spectrum**.

We can write the eigenvalue equation as:

$$\mathbf{X}^\top \mathbf{A} \mathbf{X} = \mathbf{\Lambda}$$

If \mathbf{A} is symmetric, then \mathbf{X} is an **orthogonal** matrix of eigenvectors, and $\mathbf{\Lambda}$ is a **diagonal** matrix of real eigenvalues.

We call it the **spectral decomposition** of \mathbf{A} .

Observe the similarity with our motivational problem. We can modify it to solve for **all** eigenvectors and eigenvalues:

$$\begin{aligned} \min_{\mathbf{X}} \quad & \text{tr}(\mathbf{X}^\top \mathbf{A} \mathbf{X}) \\ \text{s.t.} \quad & \mathbf{X}^\top \mathbf{X} = \mathbf{I} \end{aligned}$$

Finding eigenvalues

Power iteration

Very simple algorithm to find the **largest** eigenvalue/eigenvector:

```
function NORMALIZED-ITERATION( $A$ )  
   $\vec{v} \leftarrow \text{ARBITRARY}(n)$   
  for  $k \leftarrow 1, 2, 3, \dots$   
     $\vec{w} \leftarrow A\vec{v}$   
     $\vec{v} \leftarrow \vec{w} / \|\vec{w}\|$   
  return  $\vec{v}$ 
```

Power iteration

Very simple algorithm to find the **largest** eigenvalue/eigenvector:

```
function NORMALIZED-ITERATION( $A$ )  
   $\vec{v} \leftarrow \text{ARBITRARY}(n)$   
  for  $k \leftarrow 1, 2, 3, \dots$   
     $\vec{w} \leftarrow A\vec{v}$   
     $\vec{v} \leftarrow \vec{w} / \|\vec{w}\|$   
  return  $\vec{v}$ 
```

The normalization is needed to reduce the numerical error.

Without normalization, it will still converge to the principal eigenvector (but with a very large scale).

Inverse iteration

To find the **smallest** eigenvalue/eigenvector, we first observe that:

$$\mathbf{Ax} = \lambda\mathbf{x} \implies \mathbf{A}^{-1}\mathbf{x} = \frac{1}{\lambda}\mathbf{x}$$

Inverse iteration

To find the **smallest** eigenvalue/eigenvector, we first observe that:

$$\mathbf{A}\mathbf{x} = \lambda\mathbf{x} \implies \mathbf{A}^{-1}\mathbf{x} = \frac{1}{\lambda}\mathbf{x}$$

Then, we can just apply the power method to \mathbf{A}^{-1} :

```
function INVERSE-ITERATION( $A$ )  
   $\vec{v} \leftarrow \text{ARBITRARY}(n)$   
  for  $k \leftarrow 1, 2, 3, \dots$   
     $\vec{w} \leftarrow A^{-1}\vec{v}$   
     $\vec{v} \leftarrow \vec{w}/\|\vec{w}\|$   
  return  $\vec{v}$ 
```

Inverse iteration

To find the **smallest** eigenvalue/eigenvector, we first observe that:

$$\mathbf{A}\mathbf{x} = \lambda\mathbf{x} \implies \mathbf{A}^{-1}\mathbf{x} = \frac{1}{\lambda}\mathbf{x}$$

Then, we can just apply the power method to \mathbf{A}^{-1} :

```
function INVERSE-ITERATION( $A$ )  
   $\vec{v} \leftarrow \text{ARBITRARY}(n)$   
  for  $k \leftarrow 1, 2, 3, \dots$   
     $\vec{w} \leftarrow A^{-1}\vec{v}$   
     $\vec{v} \leftarrow \vec{w}/\|\vec{w}\|$   
  return  $\vec{v}$ 
```

In practice, you don't invert \mathbf{A} but apply LU decomposition.

Shifting

For a matrix \mathbf{A} , we have eigenvalues $\{\lambda_i\}$ and eigenvectors $\{\mathbf{x}_i\}$.

Then:

$$(\mathbf{A} - \sigma \mathbf{I})\mathbf{x}_i =$$

Shifting

For a matrix \mathbf{A} , we have eigenvalues $\{\lambda_i\}$ and eigenvectors $\{\mathbf{x}_i\}$.

Then:

$$(\mathbf{A} - \sigma \mathbf{I})\mathbf{x}_i = \mathbf{A}\mathbf{x}_i - \sigma \mathbf{x}_i =$$

Shifting

For a matrix \mathbf{A} , we have eigenvalues $\{\lambda_i\}$ and eigenvectors $\{\mathbf{x}_i\}$.

Then:

$$(\mathbf{A} - \sigma \mathbf{I})\mathbf{x}_i = \mathbf{A}\mathbf{x}_i - \sigma \mathbf{x}_i = \lambda_i \mathbf{x}_i - \sigma \mathbf{x}_i =$$

Shifting

For a matrix \mathbf{A} , we have eigenvalues $\{\lambda_i\}$ and eigenvectors $\{\mathbf{x}_i\}$.

Then:

$$(\mathbf{A} - \sigma \mathbf{I})\mathbf{x}_i = \mathbf{A}\mathbf{x}_i - \sigma \mathbf{x}_i = \lambda_i \mathbf{x}_i - \sigma \mathbf{x}_i = (\lambda_i - \sigma)\mathbf{x}_i$$

Shifting

For a matrix \mathbf{A} , we have eigenvalues $\{\lambda_i\}$ and eigenvectors $\{\mathbf{x}_i\}$.

Then:

$$(\mathbf{A} - \sigma \mathbf{I})\mathbf{x}_i = \mathbf{A}\mathbf{x}_i - \sigma \mathbf{x}_i = \lambda_i \mathbf{x}_i - \sigma \mathbf{x}_i = (\lambda_i - \sigma)\mathbf{x}_i$$

which means that the eigenvalues of $\mathbf{A} - \sigma \mathbf{I}$ are $\lambda_i - \sigma$, i.e., by shifting the matrix, we get a corresponding **shift in the eigenvalues**.

Shifting

For a matrix \mathbf{A} , we have eigenvalues $\{\lambda_i\}$ and eigenvectors $\{\mathbf{x}_i\}$.

Then:

$$(\mathbf{A} - \sigma \mathbf{I})\mathbf{x}_i = \mathbf{A}\mathbf{x}_i - \sigma \mathbf{x}_i = \lambda_i \mathbf{x}_i - \sigma \mathbf{x}_i = (\lambda_i - \sigma)\mathbf{x}_i$$

which means that the eigenvalues of $\mathbf{A} - \sigma \mathbf{I}$ are $\lambda_i - \sigma$, i.e., by shifting the matrix, we get a corresponding [shift in the eigenvalues](#).

If we think that σ is near an eigenvalue of \mathbf{A} , then $\mathbf{A} - \sigma \mathbf{I}$ has an eigenvalue close to 0.

Shifting

For a matrix \mathbf{A} , we have eigenvalues $\{\lambda_i\}$ and eigenvectors $\{\mathbf{x}_i\}$.

Then:

$$(\mathbf{A} - \sigma \mathbf{I})\mathbf{x}_i = \mathbf{A}\mathbf{x}_i - \sigma \mathbf{x}_i = \lambda_i \mathbf{x}_i - \sigma \mathbf{x}_i = (\lambda_i - \sigma)\mathbf{x}_i$$

which means that the eigenvalues of $\mathbf{A} - \sigma \mathbf{I}$ are $\lambda_i - \sigma$, i.e., by shifting the matrix, we get a corresponding **shift in the eigenvalues**.

If we think that σ is near an eigenvalue of \mathbf{A} , then $\mathbf{A} - \sigma \mathbf{I}$ has an eigenvalue close to 0.

We can use this fact to estimate **portions of the spectrum**:

- Provide a **guess** σ for an eigenvalue

Shifting

For a matrix \mathbf{A} , we have eigenvalues $\{\lambda_i\}$ and eigenvectors $\{\mathbf{x}_i\}$.

Then:

$$(\mathbf{A} - \sigma \mathbf{I})\mathbf{x}_i = \mathbf{A}\mathbf{x}_i - \sigma \mathbf{x}_i = \lambda_i \mathbf{x}_i - \sigma \mathbf{x}_i = (\lambda_i - \sigma)\mathbf{x}_i$$

which means that the eigenvalues of $\mathbf{A} - \sigma \mathbf{I}$ are $\lambda_i - \sigma$, i.e., by shifting the matrix, we get a corresponding **shift in the eigenvalues**.

If we think that σ is near an eigenvalue of \mathbf{A} , then $\mathbf{A} - \sigma \mathbf{I}$ has an eigenvalue close to 0.

We can use this fact to estimate **portions of the spectrum**:

- Provide a **guess** σ for an eigenvalue
- Compute the shifted matrix $\mathbf{B} = \mathbf{A} - \sigma \mathbf{I}$

Shifting

For a matrix \mathbf{A} , we have eigenvalues $\{\lambda_i\}$ and eigenvectors $\{\mathbf{x}_i\}$.

Then:

$$(\mathbf{A} - \sigma \mathbf{I})\mathbf{x}_i = \mathbf{A}\mathbf{x}_i - \sigma \mathbf{x}_i = \lambda_i \mathbf{x}_i - \sigma \mathbf{x}_i = (\lambda_i - \sigma)\mathbf{x}_i$$

which means that the eigenvalues of $\mathbf{A} - \sigma \mathbf{I}$ are $\lambda_i - \sigma$, i.e., by shifting the matrix, we get a corresponding **shift in the eigenvalues**.

If we think that σ is near an eigenvalue of \mathbf{A} , then $\mathbf{A} - \sigma \mathbf{I}$ has an eigenvalue close to 0.

We can use this fact to estimate **portions of the spectrum**:

- Provide a **guess** σ for an eigenvalue
- Compute the shifted matrix $\mathbf{B} = \mathbf{A} - \sigma \mathbf{I}$
- Apply the **inverse iteration** on \mathbf{B}

Finding multiple eigenvalues at once

Recall that if $\mathbf{B} = \mathbf{T}^{-1}\mathbf{A}\mathbf{T}$ then \mathbf{A} and \mathbf{B} have the same spectra.

Finding multiple eigenvalues at once

Recall that if $\mathbf{B} = \mathbf{T}^{-1}\mathbf{A}\mathbf{T}$ then \mathbf{A} and \mathbf{B} have the same spectra.

For example, we could first compute

$$\mathbf{A} = \mathbf{Q}\mathbf{R}$$

and then overwrite:

$$\mathbf{A} \leftarrow \mathbf{Q}^{\top}\mathbf{A}\mathbf{Q}$$

and still have the guarantee that the eigenvalues stay the same.

Finding multiple eigenvalues at once

Recall that if $\mathbf{B} = \mathbf{T}^{-1}\mathbf{A}\mathbf{T}$ then \mathbf{A} and \mathbf{B} have the same spectra.

For example, we could first compute

$$\mathbf{A} = \mathbf{Q}\mathbf{R}$$

and then overwrite:

$$\mathbf{A} \leftarrow \mathbf{Q}^{\top}\mathbf{A}\mathbf{Q}$$

and still have the guarantee that the eigenvalues stay the same.

Now observe:

$$\mathbf{Q}^{\top}\mathbf{A}\mathbf{Q} =$$

Finding multiple eigenvalues at once

Recall that if $\mathbf{B} = \mathbf{T}^{-1}\mathbf{A}\mathbf{T}$ then \mathbf{A} and \mathbf{B} have the same spectra.

For example, we could first compute

$$\mathbf{A} = \mathbf{Q}\mathbf{R}$$

and then overwrite:

$$\mathbf{A} \leftarrow \mathbf{Q}^{\top}\mathbf{A}\mathbf{Q}$$

and still have the guarantee that the eigenvalues stay the same.

Now observe:

$$\mathbf{Q}^{\top}\mathbf{A}\mathbf{Q} = \mathbf{Q}^{\top}(\mathbf{Q}\mathbf{R})\mathbf{Q} =$$

Finding multiple eigenvalues at once

Recall that if $\mathbf{B} = \mathbf{T}^{-1}\mathbf{A}\mathbf{T}$ then \mathbf{A} and \mathbf{B} have the same spectra.

For example, we could first compute

$$\mathbf{A} = \mathbf{Q}\mathbf{R}$$

and then overwrite:

$$\mathbf{A} \leftarrow \mathbf{Q}^\top \mathbf{A} \mathbf{Q}$$

and still have the guarantee that the eigenvalues stay the same.

Now observe:

$$\mathbf{Q}^\top \mathbf{A} \mathbf{Q} = \mathbf{Q}^\top (\mathbf{Q}\mathbf{R}) \mathbf{Q} = (\mathbf{Q}^\top \mathbf{Q}) \mathbf{R} \mathbf{Q} =$$

Finding multiple eigenvalues at once

Recall that if $\mathbf{B} = \mathbf{T}^{-1}\mathbf{A}\mathbf{T}$ then \mathbf{A} and \mathbf{B} have the same spectra.

For example, we could first compute

$$\mathbf{A} = \mathbf{Q}\mathbf{R}$$

and then overwrite:

$$\mathbf{A} \leftarrow \mathbf{Q}^\top \mathbf{A} \mathbf{Q}$$

and still have the guarantee that the eigenvalues stay the same.

Now observe:

$$\mathbf{Q}^\top \mathbf{A} \mathbf{Q} = \mathbf{Q}^\top (\mathbf{Q}\mathbf{R}) \mathbf{Q} = (\mathbf{Q}^\top \mathbf{Q}) \mathbf{R} \mathbf{Q} = \mathbf{R} \mathbf{Q}$$

Finding multiple eigenvalues at once

Recall that if $\mathbf{B} = \mathbf{T}^{-1}\mathbf{A}\mathbf{T}$ then \mathbf{A} and \mathbf{B} have the same spectra.

For example, we could first compute

$$\mathbf{A} = \mathbf{Q}\mathbf{R}$$

and then overwrite:

$$\mathbf{A} \leftarrow \mathbf{Q}^\top \mathbf{A} \mathbf{Q}$$

and still have the guarantee that the eigenvalues stay the same.

Now observe:

$$\mathbf{Q}^\top \mathbf{A} \mathbf{Q} = \mathbf{Q}^\top (\mathbf{Q}\mathbf{R}) \mathbf{Q} = (\mathbf{Q}^\top \mathbf{Q}) \mathbf{R} \mathbf{Q} = \mathbf{R} \mathbf{Q}$$

In other words, this operation **preserves the eigenvalues**:

$$\mathbf{A} \leftarrow \mathbf{R} \mathbf{Q}$$

QR iteration

If we keep doing this iteratively, we get the algorithm:

```
function QR-ITERATION( $A \in \mathbb{R}^{n \times n}$ )  
  for  $k \leftarrow 1, 2, 3, \dots$   
     $Q, R \leftarrow \text{QR-FACTORIZE}(A)$   
     $A \leftarrow RQ$   
  return  $\text{diag}(R)$ 
```

QR iteration

If we keep doing this iteratively, we get the algorithm:

```
function QR-ITERATION( $A \in \mathbb{R}^{n \times n}$ )  
  for  $k \leftarrow 1, 2, 3, \dots$   
     $Q, R \leftarrow \text{QR-FACTORIZE}(A)$   
     $A \leftarrow RQ$   
  return  $\text{diag}(R)$ 
```

- The generated matrices $\mathbf{A}_1, \mathbf{A}_2, \dots$ all have the same eigenvalues

QR iteration

If we keep doing this iteratively, we get the algorithm:

```
function QR-ITERATION( $A \in \mathbb{R}^{n \times n}$ )  
  for  $k \leftarrow 1, 2, 3, \dots$   
     $Q, R \leftarrow \text{QR-FACTORIZE}(A)$   
     $A \leftarrow RQ$   
  return  $\text{diag}(R)$ 
```

- The generated matrices $\mathbf{A}_1, \mathbf{A}_2, \dots$ all have the same eigenvalues
- The iterations **converge**, i.e. at some point $\mathbf{A}_k \approx \mathbf{A}_{k-1}$

QR iteration

If we keep doing this iteratively, we get the algorithm:

```
function QR-ITERATION( $A \in \mathbb{R}^{n \times n}$ )  
  for  $k \leftarrow 1, 2, 3, \dots$   
     $Q, R \leftarrow \text{QR-FACTORIZE}(A)$   
     $A \leftarrow RQ$   
  return  $\text{diag}(R)$ 
```

- The generated matrices $\mathbf{A}_1, \mathbf{A}_2, \dots$ all have the same eigenvalues
- The iterations **converge**, i.e. at some point $\mathbf{A}_k \approx \mathbf{A}_{k-1}$
- At convergence, we get $\mathbf{Q}_\infty^\top \mathbf{A}_\infty \mathbf{Q}_\infty = \mathbf{A}_\infty$, which implies $\mathbf{Q}_\infty \mathbf{R}_\infty = \mathbf{R}_\infty \mathbf{Q}_\infty$ and $\mathbf{A}_\infty \mathbf{Q}_\infty = \mathbf{Q}_\infty \mathbf{A}_\infty$

QR iteration

If we keep doing this iteratively, we get the algorithm:

```
function QR-ITERATION( $A \in \mathbb{R}^{n \times n}$ )  
  for  $k \leftarrow 1, 2, 3, \dots$   
     $Q, R \leftarrow \text{QR-FACTORIZE}(A)$   
     $A \leftarrow RQ$   
  return  $\text{diag}(R)$ 
```

- The generated matrices $\mathbf{A}_1, \mathbf{A}_2, \dots$ all have the same eigenvalues
- The iterations **converge**, i.e. at some point $\mathbf{A}_k \approx \mathbf{A}_{k-1}$
- At convergence, we get $\mathbf{Q}_\infty^\top \mathbf{A}_\infty \mathbf{Q}_\infty = \mathbf{A}_\infty$, which implies $\mathbf{Q}_\infty \mathbf{R}_\infty = \mathbf{R}_\infty \mathbf{Q}_\infty$ and $\mathbf{A}_\infty \mathbf{Q}_\infty = \mathbf{Q}_\infty \mathbf{A}_\infty$
- If $\mathbf{A}_\infty \mathbf{x} = \lambda \mathbf{x}$, then we can write

$$\lambda \mathbf{x} = \mathbf{A}_\infty \mathbf{x} =$$

QR iteration

If we keep doing this iteratively, we get the algorithm:

```
function QR-ITERATION( $A \in \mathbb{R}^{n \times n}$ )  
  for  $k \leftarrow 1, 2, 3, \dots$   
     $Q, R \leftarrow \text{QR-FACTORIZE}(A)$   
     $A \leftarrow RQ$   
  return  $\text{diag}(R)$ 
```

- The generated matrices $\mathbf{A}_1, \mathbf{A}_2, \dots$ all have the same eigenvalues
- The iterations **converge**, i.e. at some point $\mathbf{A}_k \approx \mathbf{A}_{k-1}$
- At convergence, we get $\mathbf{Q}_\infty^\top \mathbf{A}_\infty \mathbf{Q}_\infty = \mathbf{A}_\infty$, which implies $\mathbf{Q}_\infty \mathbf{R}_\infty = \mathbf{R}_\infty \mathbf{Q}_\infty$ and $\mathbf{A}_\infty \mathbf{Q}_\infty = \mathbf{Q}_\infty \mathbf{A}_\infty$
- If $\mathbf{A}_\infty \mathbf{x} = \lambda \mathbf{x}$, then we can write

$$\lambda \mathbf{x} = \mathbf{A}_\infty \mathbf{x} = \mathbf{Q}_\infty \mathbf{R}_\infty \mathbf{x} =$$

QR iteration

If we keep doing this iteratively, we get the algorithm:

```
function QR-ITERATION( $A \in \mathbb{R}^{n \times n}$ )  
  for  $k \leftarrow 1, 2, 3, \dots$   
     $Q, R \leftarrow \text{QR-FACTORIZE}(A)$   
     $A \leftarrow RQ$   
  return  $\text{diag}(R)$ 
```

- The generated matrices A_1, A_2, \dots all have the same eigenvalues
- The iterations **converge**, i.e. at some point $A_k \approx A_{k-1}$
- At convergence, we get $Q_\infty^\top A_\infty Q_\infty = A_\infty$, which implies $Q_\infty R_\infty = R_\infty Q_\infty$ and $A_\infty Q_\infty = Q_\infty A_\infty$
- If $A_\infty x = \lambda x$, then we can write

$$\lambda x = A_\infty x = Q_\infty R_\infty x = R_\infty Q_\infty x =$$

QR iteration

If we keep doing this iteratively, we get the algorithm:

```
function QR-ITERATION( $A \in \mathbb{R}^{n \times n}$ )  
  for  $k \leftarrow 1, 2, 3, \dots$   
     $Q, R \leftarrow \text{QR-FACTORIZE}(A)$   
     $A \leftarrow RQ$   
  return  $\text{diag}(R)$ 
```

- The generated matrices A_1, A_2, \dots all have the same eigenvalues
- The iterations **converge**, i.e. at some point $A_k \approx A_{k-1}$
- At convergence, we get $Q_\infty^\top A_\infty Q_\infty = A_\infty$, which implies $Q_\infty R_\infty = R_\infty Q_\infty$ and $A_\infty Q_\infty = Q_\infty A_\infty$
- If $A_\infty x = \lambda x$, then we can write

$$\lambda x = A_\infty x = Q_\infty R_\infty x = R_\infty Q_\infty x = \pm R_\infty x$$

QR iteration

If we keep doing this iteratively, we get the algorithm:

```
function QR-ITERATION( $A \in \mathbb{R}^{n \times n}$ )  
  for  $k \leftarrow 1, 2, 3, \dots$   
     $Q, R \leftarrow \text{QR-FACTORIZE}(A)$   
     $A \leftarrow RQ$   
  return  $\text{diag}(R)$ 
```

- The generated matrices $\mathbf{A}_1, \mathbf{A}_2, \dots$ all have the same eigenvalues
- The iterations **converge**, i.e. at some point $\mathbf{A}_k \approx \mathbf{A}_{k-1}$
- At convergence, we get $\mathbf{Q}_\infty^\top \mathbf{A}_\infty \mathbf{Q}_\infty = \mathbf{A}_\infty$, which implies $\mathbf{Q}_\infty \mathbf{R}_\infty = \mathbf{R}_\infty \mathbf{Q}_\infty$ and $\mathbf{A}_\infty \mathbf{Q}_\infty = \mathbf{Q}_\infty \mathbf{A}_\infty$
- If $\mathbf{A}_\infty \mathbf{x} = \lambda \mathbf{x}$, then we can write

$$\lambda \mathbf{x} = \mathbf{A}_\infty \mathbf{x} = \mathbf{Q}_\infty \mathbf{R}_\infty \mathbf{x} = \mathbf{R}_\infty \mathbf{Q}_\infty \mathbf{x} = \pm \mathbf{R}_\infty \mathbf{x}$$

- In other words, the eigenvalues of \mathbf{A}_∞ (and hence of \mathbf{A}) equal the diagonal elements of \mathbf{R}_∞ up to sign.

Suggested reading

Read Sections 6.1.1, 6.2, 6.2.1, 6.3.1, 6.3.2, 6.3.3, 6.4.2 of the book:

J. Solomon, “Numerical Algorithms”