

Project1: Interim Data Analysis Plan

Eduardo Romo

March 01, 2026

Introduction

Highly Active Antiretroviral Therapy (HAART) is the standard treatment regimen for HIV infection and substantially reduces disease progression by suppressing viral replication and improving immune function, as reflected by increases in CD4+ T-cell counts. The Multi-center AIDS Cohort Study is a prospective cohort study of HIV infected homosexual and bisexual men that includes longitudinal laboratory, clinical, and quality of life measurements collected over multiple years following HAART initiation.

Investigators are interested in whether baseline hard drug use is associated with poorer treatment response after initiating HAART. The primary scientific hypothesis is that participants reporting hard drug use at baseline experience worse outcomes two years after treatment initiation compared with those who do not report hard drug use. Treatment response was evaluated using four complementary outcomes: change in viral load, CD4+ T cell count, and physical and mental quality of life scores.

The corresponding statistical hypotheses evaluate whether mean treatment response at two years differs by baseline hard drug use status for each outcome, adjusting for age, BMI, smoking status, education, and race/ethnicity. For each outcome, the null hypothesis is that there is no difference in mean treatment response between subjects who report hard drug use at baseline and those who do not, while the alternative hypothesis is that a difference

exists. These hypotheses will be evaluated using both bayesian and frequentist methods.

Methods

Data Cleaning and Management

The initial dataset included 715 unique patients with longitudinal clinical and quality of life measurements following HAART initiation. Data were recorded in long format with repeated observations by study year. This analysis focused on baseline (year 0) and two year follow up (year 2) measurements to evaluate short term treatment response.

Several categorical variables were recoded to improve interpretability and reduce sparse cell counts. Race was collapsed to White vs Other, smoking status to Current smoker vs Not current smoker, adherence to 95% vs <95%, and education to College degree vs No college degree. Baseline hard drug use was represented as a binary exposure (Yes vs No). BMI values outside plausible physiologic ranges (0 or 250) were treated as implausible and coded as missing.

Treatment response outcomes were defined using baseline and year 2 measurements: change in mental quality of life, change in physical quality of life, change in CD4+ T-cell count, and the log10 ratio of viral load (year 2 vs baseline). Viral load was log-transformed to reduce skewness and improve distributional symmetry; baseline viral load was similarly transformed for model adjustment. Histograms were used to assess outcome distributions.

The analysis dataset was created by restricting data to year 0 and 2, reshaping to wide format (one row per patient), and computing derived outcomes. Missingness was examined overall and stratified by baseline hard drug use. Because outcome missingness differed by exposure status, missing completely at random was considered unlikely. Analyses were therefore conducted using a complete-case approach, assuming outcomes were missing at random conditional on observed covariates included in the models. The final analytic sample included 476 participants.

Statistical Analysis

The primary objective was to estimate the association between baseline hard drug and treatment response at two years. For each outcome, two linear regression models were fit: (1) a primary model adjusting for baseline outcome value, age, BMI, smoking status, education, and race/ethnicity; and (2) a secondary model additionally adjusting for year 2 adherence to assess how the hard drug use coefficient changed after accounting for adherence (a naive mediation analysis). Frequentist linear models were fit primarily as a validation check against the bayesian estimates.

Frequentist results were summarized using point estimates, standard errors, two-sided p-values, and 95% confidence interval. Bayesian models were fit using brms package with the cmdstanr backend. Non-informative priors were specified to reflect minimal prior knowledge: Normal(0, 100²) for intercept and coefficients and Half-Normal(0, 100²) for residual standard deviation. Posterior inference for the hard drug use coefficient was summarized using posterior means, posterior standard deviations, and 95% highest density posterior intervals (HPDIs). Clinical relevance was assessed using posterior probabilities that the absolute effect exceeded pre-specified clinically meaningful thresholds: 2 points for quality-of-life outcomes, 50 cells for CD4+ count, and 0.5 for log10 viral load ratio. Models were fit using four chains with 20,000 iterations per chain, including 5,000 warmup iterations.

Predictive performance of Bayesian models was evaluated using leave-one-out cross-validation (LOO). Differences in LOO information criterion (ΔLOOIC) were used to compare full and reduced models, with lower LOOIC indicating improved expected predictive performance. Model diagnostics included trace plots, autocorrelation checks, split-chain $\hat{R}$ values near 1, and effective sample sizes exceeding 1000 for all parameters.

Results

Descriptive statistics

Baseline characteristics stratified by baseline hard drug use are shown in Table 1. 39 (8.2%) subjects of the 476 in the study reported hard drug use at baseline. Overall demographic and clinical characteristics were generally similar across groups;

Table 1: Baseline characteristics stratified by baseline hard drug use.

Variable	Overall	No hard drugs	Yes hard drugs
Sample size			
N	476	437	39
BMI			
Mean (SD)	25.2 (4.3)	25.3 (4.4)	23.6 (3.4)
Missing, n (%)	13 (2.7%)	10 (2.3%)	3 (7.7%)
Age			
Mean (SD)	43.3 (8.7)	43.1 (8.7)	44.6 (9.5)
CD4+ Count			
Mean (SD)	373.5 (200.5)	375.4 (201.1)	352.2 (194.7)
Log Viral Load			
Mean (SD)	4.5 (0.9)	4.5 (0.9)	4.5 (0.9)
Mental Quality of Life			
Mean (SD)	44.9 (13.5)	45.1 (13.7)	42.3 (11.2)
Physical Quality of Life			
Mean (SD)	51.0 (9.1)	51.3 (9.1)	47.7 (8.5)
Smoking status			
Not current smoker	291 (61.1%)	282 (64.5%)	9 (23.1%)
Current smoker	185 (38.9%)	155 (35.5%)	30 (76.9%)
Adherence category (year 2)			
>= 95% Adherence	426 (89.5%)	388 (88.8%)	38 (97.4%)
< 95% Adherence	50 (10.5%)	49 (11.2%)	1 (2.6%)
Race/ethnicity			
White	298 (62.6%)	279 (63.8%)	19 (48.7%)
Other	178 (37.4%)	158 (36.2%)	20 (51.3%)
Education			
No College Degree	271 (56.9%)	242 (55.4%)	29 (74.4%)
College Degree	205 (43.1%)	195 (44.6%)	10 (25.6%)

however, participants reporting hard drug use were more likely to be current smokers (76.9% vs 36.5%) and less likely to have a college degree (25.6% vs 44.6%). Baseline values for CD4+ count, viral load, and quality of life measures were generally comparable between groups, with slightly lower baseline mental and physical quality of life scores among hard drug users.

Figure 1 displays boxplots of the four treatment response outcomes. Descriptively, hard drug users appeared to show similar improvements in physical quality of life and CD4+ cell count, whereas differences in mental quality of life and viral load were less substantial.




Figure 1: Treatment Response (Year 2 - Year 0) by Hard Drug Use

Main analysis: association between hard drug use and treatment response

Table 2 summarizes regression results evaluating the association between baseline hard drug use and treatment response outcome, adjusting for baseline outcome value, age, BMI, smok-

ing status, education, and race/ethnicity. Bayesian estimates are presented as the primary results, with frequentist estimates included for comparison.

For change in mental quality of life, the posterior mean coefficient for hard drug use was -0.29 (95% HPDI: -3.78, 3.16), indicating little evidence of an association. The posterior probability (0.261) of a clinically meaningful effect was low. For physical quality of life, hard drug use was associated with poorer treatment response (posterior mean -3.34 ; 95% HPDI: -6.05 , -0.60). The posterior probability (0.834) that the effect exceeded the predefined clinically meaningful threshold suggested moderate evidence of a clinically relevant reduction in improvement among hard drug users. The Bayesian model estimated a mean difference of -148.85 cells/ μ L (95% HPDI: -209.06 , -90.29), indicating substantially poorer immunologic recovery among participants reporting hard drug use. Posterior probability (~ 1.00) results supported strong evidence of a clinically meaningful effect. For viral load change, the posterior mean coefficient was close to zero (-0.60 ; 95% HPDI: -0.47 , 0.33), suggesting little evidence of an association. Posterior probability (0.018) of a clinically meaningful effect was low.

Across outcomes, bayesian and frequentist estimates were highly concordant. Model comparison using change in LOOIC favored models including hard drug use for CD4+ count and physical quality-of-life outcomes, while improvements were minimal for mental quality of life and viral load, consistent with weaker evidence for these associations.

Table 2: Main analysis: hard drug use coefficient by outcome (Frequentist vs Bayesian).

Outcome	Frequentist			Bayesian			
	Estimate (SE)	95% CI	p-value	Post. mean (SD)	95% HPDI	Pr(> clin)	Δ LOOIC (red-full)
Δ AGG_MENT	-0.29 (1.76)	[-3.75, 3.17]	0.8699	-0.29 (1.77)	[-3.78, 3.16]	0.261	-2.9
Δ AGG_PHYS	-3.34 (1.39)	[-6.06, -0.61]	0.0166	-3.34 (1.39)	[-6.05, -0.60]	0.834	3.6
Δ CD4 (LEU3N)	-163.88 (31.79)	[-226.35, -101.41]	<0.001	-148.85 (30.36)	[-209.06, -90.29]	1.000	23.6
log10(VLOAD2/VLOAD0)	-0.06 (0.20)	[-0.46, 0.34]	0.7698	-0.06 (0.20)	[-0.47, 0.33]	0.018	-1.9

Secondary analysis: adjustment for adherence to treatment

Descriptive measures from Table 1 demonstrate there is a higher percentage (97.4%) of hard drug users with $\geq 95\%$ adherence compared to non hard drug users (89.5%), indicating a potential relationship between hard drug use and treatment adherence. Table 3 presents results from models additionally adjusting for treatment adherence at year 2. The primary goal was to assess how the hard drug use coefficient changed after inclusion of adherence. The frequentist results are again shown primarily as a reference. For mental quality of life, the hard drug use coefficient changed modestly from -0.29 to -0.53 after adjustment for adherence, indicating little meaningful change in effect magnitude. Similarly, for physical quality of life, the estimate changed only slightly (approximately -3.34 to -3.57), suggesting that adherence did not substantially explain the association between hard drug use and physical quality of life change. For CD4+ cell count, adjustment for adherence led to a modest increase in magnitude (approximately -148.85 to -153.79), again indicating minimal change of the hard drug use effect. In contrast, the viral load coefficient moved closer to zero after adjustment (from -0.06 to -0.01), though both estimates remained small and highly uncertain. Overall, these findings suggest limited evidence that adherence substantially explained the association between baseline hard drug use and treatment response.

Table 3: Secondary analysis (naive mediation): hard drug use coefficient by outcome (Frequentist vs Bayesian).

Outcome	Frequentist				Bayesian				
	Estimate (SE)	95% CI	p-value	Δ (with-no)	Post. mean (SD)	95% HPDI	Pr(> clin)	Δ (with-no)	Δ LOOIC (no-with)
Δ AGG_MENT	-0.53 (1.77)	[-4.00, 2.94]	0.7639	-0.24	-0.53 (1.77)	[-3.99, 2.96]	0.279	-0.24	-0.1
Δ AGG_PHYS	-3.57 (1.39)	[-6.30, -0.84]	0.0106	-0.23	-3.57 (1.39)	[-6.30, -0.85]	0.871	-0.23	1.5
Δ CD4 (LEU3N)	-169.68 (31.82)	[-232.21, -107.15]	<0.001	-5.80	-153.79 (30.42)	[-213.89, -94.54]	1.000	-4.93	2.2
log10(VLOAD2/VLOAD0)	-0.01 (0.20)	[-0.41, 0.38]	0.9500	0.05	-0.01 (0.20)	[-0.42, 0.37]	0.014	0.05	3.8

Model diagnostics

Diagnostics indicated good convergence and sampling performance for all Bayesian models. Maximum $\hat{R}$ values were approximately 1.00, effective sample sizes were large (all > 26,000), and no divergent transitions were observed. Trace plots demonstrated good chain mixing,

and autocorrelation decreased rapidly across lags. These diagnostics support the stability and reliability of the posterior estimates.

Discussion

This analysis evaluated whether hard drug use was associated with treatment response two years after HAART initiation, using both bayesian and frequentist regression models. Across outcomes, results suggested that hard drug use was associated with poorer treatment response for physical quality of life and CD4+ cell count, while little evidence was observed for associations with mental quality of life or viral load change. Bayesian estimates were consistent with frequentist results, giving us reassurance that model specification was appropriate. Model comparison using changes in LOOIC supported stronger evidence for associations involving physical QoL and immunologic recovery than for the other outcomes.

The posterior probabilities of clinically meaningful effects provided interpretable measure of clinical relevance. These probabilities suggested meaningful evidence of clinically important differences in physical quality of life and CD4+ cell count for hard drug users, whereas clinically meaningful effects were unlikely for mental QoL and viral load.

Several limitations should be considered when interpreting these results. First, this was a complete case analysis. Although we believe missingness is at random for the outcomes, bias may still arise if missingness depended on unobserved factors. Second, covariates were collapsed into binary categories due to sparse sample sizes in some levels, which may have reduced the ability to detect nuanced associations. Third, the secondary analysis evaluating adherence to treatment was not a formal mediation analysis. While our results from this naive “mediation” analysis may provide descriptive insight, this approach does not estimate direct and indirect effects. Future work could extend this analysis using a formal mediation framework to rigorously evaluate whether adherence mediates the relationship between hard drug use and treatment response. Overall, these results suggest that baseline hard drug

use may be an important predictor of poor physical and immunologic outcomes following HAART initiation. Future studies incorporating longitudinal methods, alternative missing data methods, and better measurements of behavioral factors can help clarify the mechanisms behind these associations.

Reproducible Research Information

<https://github.com/eromo96/BIOS6624.git>

Data Location: /Users/eduardo/Eduardo/UColorado PhD Bios/Spring 26/BIOS6624

Advanced Statistical Methods/BIOS6624/Project1/Data