PYQCTools Documentation

Release 1.0

Enrico Ronca

May 17, 2016

1 Contents **3**
1.1 Tools for Omega Space Green's Functions 3
1.2 Tools for Real-Time Green's Functions 3
1.3 Tools for Integrals Dumping 4

PYQCTools is a collection of python scripts useful to dump quantum chemistry integrals and to perform data post-processing for different quantum chemistry methods.

PYQCTools requires the following prerequisites to work:

- Python 2.6, 2.7, 3.2, 3.3, 3.4
- Numpy 1.6.2 or higher
- Scipy 0.10 or higher (0.12.0 or higher for python 3.3, 3.4)
- PySCF for Integrals Calculation in Integrals Dumpings scripts.

CHAPTER

ONE

CONTENTS

1.1 Tools for Omega Space Green's Functions

`gf_trace.py`: Calculate the Density of States (DOS) value from an ω-dependent Green's Function. It makes the trace of the Green's Function associated with a certain frequency value.

Example :

```
>>> python gf_trace.py /PATH/green.txt omega_value
```

`green.txt`: formatted text file containing the Green's function.

`omega_value`: double frequency value.

1.2 Tools for Real-Time Green's Functions

`rtgf.py`: Calculate the Density of States (DOS) values during a time propagation. It makes the trace of time-dependent Green's Functions calculated along a time propagation and return the DOS values as a function of time both for the real and for the imaginary part of the Green's Function.

Example :

```
>>> python rtgf.py prop_time time_step /PATH/scratch
```

`prop_time`: double value of the full propagation time (period).

`time_step`: double value of the time-step.

`scratch`: directory containing text files of the real (green.$t.$t.txt) and imaginary (green.30000+$t.30000+$t.txt) Green's Functions, where $t indicate the specific time-step.

The script save two output files, `rt_real.txt` and `rt_imag.txt`, containing the real and imaginary part of the time-dependent DOS respectively.

`fft.py`: Perform the fourier transform of the time-dependent Density of States (DOS). It reads the `rt_real.txt` and `rt_imag.txt` generated by the `rtgf.py` script and produces the `ldos.out` and `real_part.txt` files containing the imaginary and real parts of the *omega*-dependent DOS respectively.

Example :

```
>>> python fft.py broad rem_add
```

`broad`: double value of the imaginary broadening.

`rem_add`: string specifying if we are working with the addition or removal part of the Green's Function. It can assume only the values 'add' or 'rem'.

`extrapolation.py`: Perform linear prediction to extend the total propagation time of a time propagation.

It reads N points of time-dependent DOS inside the files `rt_real.txt` and `rt_imag.txt` and use the last N/2 data to predict the following N points.

Example :

```
>>> python extrapolation.py full_range
```

`full_range`: boolean variable. If it is true is return the full range of calculated and predicted values, if it is false it returns only the predicted values.

The script produces `new_full_data.out` files if full_range = true otherwise it produces `predicted.out` output files.

`iter_extrapolation.py`: Perform an interative linear prediction to extend the total propagation time of a time propagatio

It reads 4 points of the time-dependent DOS inside the files `rt_real.txt` and `rt_imag.txt` and use the last 2 of them to predict the following N points.

Example :

```
>>> python extrapolation.py N
```

`N`: integer variable specifying the total number of points that need to be predicted.

The script produces `new_full_real.out` and `new_full_imag.out` output files with the real and imaginary parts of the exteded time-dependent DOS respectively.

1.3 Tools for Integrals Dumping

`Integrals_dump.py`: Dump 1 and 2-elctron integrals in the MO basis inside a CASCI space in FCIDUMP format.
The PySCF input to calculate integrals is already included in the script.

`DipoleIntegrals_dump.py`: Dump dipole integrals in the MO basis in a CASCI space in FCIDUMP format.
The PySCF input to calculate integrals is already included in the script.

`LowdinOrtho_Integrals.py`: Dump 1 and 2-electron integrals in the Localized basis obtained by Lowdin Orthogonalizatio
The PySCF input to calculate integrals is already included in the script.

You can also download the PDF version of this manual.