

Wetland_emissions_r2.R

August 9, 2024

Contents

SOCCR_Wetlands	1
Index	4

SOCCR_Wetlands	<i>Create SOCCR based wetland methane maps</i>
----------------	--

Description

‘SOCCR_Wetlands’ writes 2 netcdf files of gridded wetland methane emissions, 1 for SOCCR1 and 1 for SOCCR2. May include freshwater wetland emissions using Rosentretet et al. too. Includes optional visuals as well.

Usage

```
SOCCR_Wetlands(  
  output_directory,  
  plot_directory,  
  domain,  
  Use_SOCCR1,  
  Use_SOCCR2,  
  Include_freshwater,  
  Wetland_EFs,  
  verbose,  
  County_Tigerlines,  
  State_Tigerlines,  
  focus_city_tigerlines  
)
```

Arguments

- output_directory Character providing the full filepath to save processed data
- plot_directory Character providing the full filepath to save figures. Only relevant if verbose = TRUE.
- domain SpatRaster providing the desired output grid, including the desired resolution and coordinate reference system

Use_SOCCR1	Logical. Pulled from config file. Indicating whether or not to calculate emissions using SOCCR1.
Use_SOCCR2	Logical. Pulled from config file. Indicating whether or not to calculate emissions using SOCCR2.
Include_freshwater	Logical. Pulled from config file. Indicating whether or not to calculate emissions for freshwater wetlands using Rosentreter et al.
Wetland_EFs	Data frame. Pulled from config file. Emission factors to use for all wetlands - including SOCCR1, SOCCR2, and Rosentreter et al.
verbose	Logical indicating whether to save additional output. This includes monthly plots of the gridded methane emissions on log scales, saved separately for each model subset and each land cover dataset used.
County_Tigerlines	SpatVector. United States Census Bureau county shapefile. Available at https://www.census.gov/geographies/mapping-files/time-series/geo/tiger-line-file.html . Only relevant if verbose=TRUE.
State_Tigerlines	SpatVector. United States Census Bureau county shapefile. Available at https://www.census.gov/geographies/mapping-files/time-series/geo/tiger-line-file.html . Only relevant if verbose=TRUE.
focus_city_tigerlines	SpatVector. United States Census Bureau county shapefile. Available at https://www.census.gov/geographies/mapping-files/time-series/geo/tiger-line-file.html . Only relevant if a focus city was set in main and verbose=TRUE.

Details

This function takes the output of 'NWI_Wetland_fraction', which is per pixel fractional coverage of wetlands separated by wetland type, and applies emission factors to convert coverage to methane emissions. It is simply applying SOCCR1 or SOCCR2 average emissions from wetlands to the NWI activity data. Additionally, freshwater emissions are calculated using emissions from Rosentreter et al.

SOCCR1 values are based on the arithmetic averages of Table F5, SOCCR2 values are based on the arithmetic averages of Tables 13B.8 to 13B.11 for PFO and PNF and Table 15A.2 for M2 and E2, and Lakes and Rivers (L1, L2, and R1 - R4) are from Rosentreter et al. using the median flux from rivers and the largest lake class (>1 km). This lake flux was chosen as McDonald et al. show that large lakes (>1 km²) constitute 71% of the total lake area in CONUS (rising to 90% if including the Great Lakes).

SOCCR1 is available at <https://www.carboncyclescience.us/state-carbon-cycle-report-soccr> and SOCCR2 is available at <https://carbon2018.globalchange.gov/>.

See references [McDonald et al.](#) and [Rosentreter et al.](#)

Value

Nothing is returned from the function, but the main outputs are 2 netcdf files of the methane emissions from wetlands with 1 file for SOCCR1 and 1 file for SOCCR2 based emissions. Lakes and rivers from Rosentreter et al. emissions can also be included. They are titled "SOCCR1.nc" and "SOCCR2.nc".

If verbose is set to TRUE, then multiple figures are also saved. Log scale plots with consistent axes are saved for the 2 SOCCR emissions. They are saved as "SOCCR1.png" and "SOCCR2.png".

Author(s)

Joe Pitt, <madeup@wisc.edu>

Kris Hajny, <blank@fake.edu>

Israel Lopez-Coto, <test@test.edu>

Examples

```
library(terra)
grid_bbox=cbind(c(-76.65,-73.65),c(38.97,40.97))
grid_res=0.01
grid_crs="epsg:4326"
grid <- rast(nrows=diff(range(grid_bbox[,2]))/grid_res,
             ncols=diff(range(grid_bbox[,1]))/grid_res, xmin=min(grid_bbox[,1]),
             xmax=max(grid_bbox[,1]), ymin=min(grid_bbox[,2]), ymax=max(grid_bbox[,2]),
             crs=grid_crs)
EFs <- data.frame("E2"=c(10.3,15.29*16.043/12.011),
                  "M2"=c(10.3,15.29*16.043/12.011),
                  "PFO"=c(36,18.52*16.043/12.011),
                  "PNF"=c(36,24.92*16.043/12.011),
                  "L1"=5,
                  "L2"=5,
                  "R1"=7.88,
                  "R2"=7.88,
                  "R3"=7.88,
                  "R4"=7.88)
rownames(EFs) <- c("SOCCR1","SOCCR2")

# convert from g CH4 per m2 per yr to nmol/m2/s
EFs=EFs*1E9/(16.043*365.25*24*60*60)

SOCCR_Wetlands(output_directory=~/.//Desktop/",
               plot_directory=~/.//Desktop/plots/",
               domain=grid,
               Use_SOCCR1=TRUE,
               Use_SOCCR2=TRUE,
               Include_freshwater=TRUE,
               Wetland_EFs=EFs,
               verbose=TRUE,
               County_Tigerlines=vect("~/.//Desktop/County_Tigerlines/tl_2018_us_county.shp"),
               State_Tigerlines=vect("~/.//Desktop/State_Tigerlines/tl_2018_us_state.shp"),
               focus_city_tigerlines=focus_city)
```

Index

SOCCR_Wetlands, [1](#)