

Modulo Synthesis

A Pedagogic Introduction to Emergent Spacetime

Juan Pablo Silva Alvarado

*“Everything emerges — step by step — from counting events and
measuring how distinguishable they are.”*

February 9, 2026

Contents

1	Axiomatic Foundation	1
1.1	Why a Regular Lattice Doesn't Work	1
1.2	Axiom 1: Discrete Causal Structure	1
1.3	Axiom 2: Information Geometry	2
1.4	What We Have So Far	2
2	Spectral Dimension	3
2.1	What Do We Mean by "Dimension"?	3
2.2	Large Scales: The IR Limit	4
2.3	Small Scales: The UV Limit	4
2.4	The Flow Equation	4
3	Emergent Gravity	5
3.1	Local Horizons	5
3.2	The Thermodynamic Hypothesis	5
3.3	Entropy from Link Counting	6
3.4	Deriving the Einstein Equations	6
4	Emergent Matter	7
4.1	The Minimal Spatial Simplex	7
4.2	Why Three Anchors?	7
4.3	Gauge Groups from Geometry	8
4.3.1	Color $SU(3)_c$ from Permutations	8
4.3.2	Weak Isospin $SU(2)_L$ from the Frame Bundle	8
4.3.3	Hypercharge and Fractional Charges	8
4.4	Flavor Mixing and the CKM Matrix	8
5	Stability & Scaling	9
5.1	Higgs Mass: Boundary Inertia	9
5.1.1	The Saturation Boundary	9
5.1.2	The Calculation	9
5.2	Strong CP Problem	9
5.3	Proton Stability	10
5.4	Master Unification Table	10
6	Unification & Predictions	11
6.1	The Master Scaling Law	11
6.2	Geometric Bare Values	11
6.3	Cosmological Predictions	12
6.4	Falsifiable Tests	12
7	Conclusion	13

Chapter 1

Axiomatic Foundation

Chapter Overview

- Why a regular lattice doesn't work
- Axiom 1: Discrete Causal Structure
- Axiom 2: Information Geometry
- What we build from these two axioms

We begin with the simplest possible description of reality. We do not assume continuous space, time, or a metric tensor. We build everything from two very basic ideas.

1.1 Why a Regular Lattice Doesn't Work

Imagine we try to build spacetime from a regular 3D grid of points, like a crystal.

In the rest frame of the grid, the density is constant: one point every ℓ^3 volume. Now imagine you move past the grid at high speed. Because of length contraction, the points appear closer together in the direction of motion.

A boosted observer would measure a higher density. This means they could detect their own absolute velocity relative to the grid — which contradicts special relativity.

A regular lattice cannot be the fundamental structure of spacetime because it breaks Lorentz invariance.

1.2 Axiom 1: Discrete Causal Structure

Instead of a regular lattice, we use something much more random and democratic.

Axiom 1: Causal Set

Spacetime is a set of discrete events with a causal order. Formally, it is a pair $(\mathcal{C}, \prec)$ where:

1. **Transitivity:** If $x \prec y$ and $y \prec z$, then $x \prec z$
2. **Acyclicity:** $x \not\prec x$ (no time travel)
3. **Local Finiteness:** Between any two events there are only finitely many others

To generate such a structure, we place points randomly in a 4-dimensional volume with constant average density ρ . This is called **Poisson sprinkling**.

The expected number of events in a region of volume V is $N = \rho V$. The actual number fluctuates around this mean according to the Poisson distribution.

This construction automatically respects Lorentz invariance on average — no preferred frame.

1.3 Axiom 2: Information Geometry

We now have a set of points with causal order, but we still have no notion of distance, angle, or curvature. How do we define geometry?

Axiom 2: Fisher Metric

The geometry is defined by how distinguishable different configurations of the causal set are.

Imagine we have two slightly different ways the points could be arranged, labelled by a parameter θ^μ . Each arrangement has a probability distribution $p(C|\theta)$, where C is a particular causal set configuration.

The natural distance between two nearby configurations θ and $\theta + \delta\theta$ is given by the **Fisher information metric**:

$$I_{\mu\nu}(\theta) = \mathbb{E} \left[\frac{\partial \ln p}{\partial \theta^\mu} \frac{\partial \ln p}{\partial \theta^\nu} \right] \quad (1.1)$$

This metric measures how much the probability distribution changes when we shift the parameters. It is the unique metric that remains unchanged under coarse-graining (averaging over fine details).

*This axiom turns **information distinguishability** into geometry.*

1.4 What We Have So Far

From Axiom 1 we get discreteness and causality. From Axiom 2 we get a metric that emerges purely from statistical distinguishability.

In the following chapters we will show that:

- This metric naturally becomes Lorentzian
- Curvature appears when the sprinkling density varies
- Gravity arises as a thermodynamic force on horizons

No continuum, no metric tensor, no fields are assumed from the start. Everything is built step by step from these two axioms.

Chapter 2

Spectral Dimension

Chapter Overview

- Defining dimension through diffusion
- Behavior at large scales (IR limit)
- Behavior at small scales (UV limit)
- The flow equation

We now have a set of discrete events with causal order (from Chapter 1). We want to ask a very natural question:

“How many dimensions does this set of points look like — when we look at it from far away?”

In ordinary continuous space, the answer is always 4 (or 3+1). But in a discrete structure, the answer can change depending on how closely we look.

2.1 What Do We Mean by “Dimension”?

Dimension is not a fixed property — it is something we measure by watching how things spread out.

Imagine we drop a drop of ink at one event and watch how far it can travel after a short time σ .

- In 1D: The number of places the ink can reach grows like $\sigma^{1/2}$
- In 2D: It grows like σ^1
- In 4D: It grows like σ^2

Definition 1: Spectral Dimension

We define the **spectral dimension** $d_S(\sigma)$ as:

$$d_S(\sigma) = -2 \frac{d \ln P_r(\sigma)}{d \ln \sigma} \quad (2.1)$$

where $P_r(\sigma)$ is the probability that the ink returns to the starting point after time σ .

This is the same definition used in quantum gravity and fractal geometry.

2.2 Large Scales: The IR Limit

When we look at very large distances (many events involved), the causal set looks smooth. The random walk behaves almost like ordinary diffusion in 4 dimensions.

The return probability becomes:

$$P_r(\sigma) \sim \frac{1}{(4\pi\sigma)^{d/2}} \quad \text{with } d = 4 \quad (2.2)$$

Taking logarithms and differentiating gives:

$$d_S(\sigma \gg \ell_P^2) = 4 \quad (2.3)$$

On large scales, we recover the 4-dimensional world we observe.

2.3 Small Scales: The UV Limit

When we look very closely (distances near ℓ_P), the causal set looks very different.

Most pairs of nearby events are **not** connected by a causal link. The structure is almost tree-like: branches go forward in time, but almost no loops form.

On a tree, random walks are very inefficient at returning:

$$P_r(\sigma) \sim \frac{1}{\sigma} \quad (2.4)$$

Differentiating gives:

$$d_S(\sigma \sim \ell_P^2) = 2 \quad (2.5)$$

*At the smallest scales, the effective dimension drops to 2. This is why gravity becomes **asymptotically safe** — in 2D, Newton's constant G is dimensionless.*

2.4 The Flow Equation

The change from 2 to 4 does not happen suddenly. There is a smooth crossover:

$$d_S(k) = 2 + \frac{2}{1 + (k/\alpha_{\text{trans}})^2} \quad (2.6)$$

where $k \sim 1/\sqrt{\sigma}$ is the inverse length scale and $\alpha_{\text{trans}} \approx 0.31$.

- When $k \ll \alpha_{\text{trans}}$ (large distances): $d_S \rightarrow 4$
- When $k \gg \alpha_{\text{trans}}$ (small distances): $d_S \rightarrow 2$

***The dimension is not fixed. It flows.** This flow is the key mechanism that controls renormalization and the running of couplings.*

Chapter 3

Emergent Gravity

Chapter Overview

- Local horizons and accelerated observers
- The thermodynamic hypothesis
- Deriving the area law from link counting
- The Einstein equations as an equation of state

We now have discrete events with causal order (Chapter 1) and a geometry defined by information distinguishability (Chapter 2). We want to understand how gravity arises — not as a fundamental force, but as something that emerges naturally from this structure.

3.1 Local Horizons

Imagine you are standing still in flat spacetime. Now imagine you accelerate upward with constant acceleration a .

According to special relativity, you will see something strange: there is a region below you from which no light can ever reach you. This is your **Rindler horizon**.

For an accelerated observer, spacetime looks like it has a horizon — even though globally it is flat.

3.2 The Thermodynamic Hypothesis

We make a simple but powerful assumption, first proposed by Jacobson (1995):

Axiom 1: Thermodynamic Hypothesis

The horizon behaves thermodynamically: energy crossing the horizon is related to entropy change by the Clausius relation

$$\delta Q = T dS \tag{3.1}$$

For an accelerated observer, the temperature is the **Unruh temperature**:

$$T_{\text{Unruh}} = \frac{\hbar a}{2\pi k_B c} \tag{3.2}$$

3.3 Entropy from Link Counting

Consider the horizon as a surface in the causal set. Links that cross this surface connect the “inside” (past) to the “outside” (future).

Each crossing link represents a small piece of causal information — roughly 1 bit.

The total number of links crossing a surface of area A is approximately:

$$N_{\text{cross}} \approx \frac{A}{\ell_{\text{P}}^2} \quad (3.3)$$

Theorem 2: Bekenstein-Hawking Area Law

The entropy of a horizon is:

$$S_{\text{horizon}} \approx N_{\text{cross}} \times \ln 2 \approx \frac{A}{4\ell_{\text{P}}^2} \quad (3.4)$$

This is derived from counting causal links, not assumed.

3.4 Deriving the Einstein Equations

Energy flux crossing the horizon is:

$$\delta Q = \int T_{\mu\nu} k^\mu k^\nu d\Sigma d\lambda \quad (3.5)$$

Area change is governed by the **Raychaudhuri equation**:

$$\frac{d\theta}{d\lambda} = -\frac{1}{2}\theta^2 - \sigma^2 - R_{\mu\nu} k^\mu k^\nu \quad (3.6)$$

Demanding the Clausius relation holds for **all** local horizons gives:

$$8\pi G T_{\mu\nu} k^\mu k^\nu = R_{\mu\nu} k^\mu k^\nu \quad (3.7)$$

Since this must hold for all null vectors k^μ :

Theorem 3: Einstein Field Equations

$$R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu} + \Lambda g_{\mu\nu} = 8\pi G T_{\mu\nu} \quad (3.8)$$

*Gravity is not a fundamental interaction — it is a **thermodynamic consequence** of discrete causal structure.*

Chapter 4

Emergent Matter

Chapter Overview

- The minimal spatial simplex
- Why exactly three anchors
- Gauge groups from geometry
- Flavor mixing and the CKM matrix

We now have spacetime as a discrete causal set, geometry from information, and gravity from thermodynamics. We want to understand where particles and forces come from.

4.1 The Minimal Spatial Simplex

In continuous spacetime, the smallest volume is zero. In a discrete causal set, there is a limit.

Definition 1: 3-Simplex

The minimal spatial volume consists of:

- One event called the “root” R
- Three other events A, B, C in the future of R , but **mutually incomparable** (spacelike separated)

4.2 Why Three Anchors?

- **Two anchors:** Too simple — no room for color triality
- **Four anchors:** In the UV limit ($d_S \rightarrow 2$), the graph becomes tree-like. A 4-simplex is exponentially suppressed: $P \sim e^{-N^2}$

*The 3-simplex is the **maximal stable clique** allowed by the UV topology.
Prediction: Exactly **three generations** of matter.*

4.3 Gauge Groups from Geometry

4.3.1 Color $SU(3)_c$ from Permutations

The three anchors A, B, C are geometrically indistinguishable. The group of permutations is S_3 (order 6).

In the continuum limit, S_3 embeds into $SU(3)_c$ as its fundamental representation.

Color $SU(3)_c$ emerges from the permutation symmetry of the three anchors.

4.3.2 Weak Isospin $SU(2)_L$ from the Frame Bundle

The local Lorentz group is $SO(3,1)$. Its double cover is $SL(2, \mathbb{C})$. The maximal compact subgroup preserving the causal arrow is $SU(2)$.

Because the underlying structure is ordered (asymmetric in time), the bundle is **chiral** — it acts only on left-handed states.

4.3.3 Hypercharge and Fractional Charges

A quark localized on a single anchor sees $1/3$ of the total S_3 winding.

Quarks necessarily have fractional charges $(+2/3, -1/3)$ because they occupy a fractional geometric volume.

4.4 Flavor Mixing and the CKM Matrix

At the bare Planck scale, the three anchors are completely symmetric. The bare mass matrix is **democratic**:

$$M_{\text{bare}} \propto \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} \quad (4.1)$$

This gives maximal mixing with $\lambda_{\text{geom}} = 1/\sqrt{6} \approx 0.408$.

When we coarse-grain, high-momentum modes are suppressed by the d_S flow. After running down to low energy:

$\lambda_{\text{obs}} \approx 0.408 \times 0.55 \approx 0.225$
This matches the experimental Cabibbo angle.

Chapter 5

Stability & Scaling

Chapter Overview

- The Higgs mass from boundary inertia
- The Strong CP problem solved by stochastic averaging
- Proton stability from topological protection
- The master unification table

5.1 Higgs Mass: Boundary Inertia

5.1.1 The Saturation Boundary

The causal set has a maximum information density — roughly 1 bit per Planck volume. When we try to build a particle, it displaces this boundary.

Pushing against a saturated boundary costs energy — like stretching a rubber sheet. This is the **inertia** of the boundary.

5.1.2 The Calculation

Energy cost for a fluctuation of size r :

$$E(r) \approx T \times r^2 \approx M_{Pl}^3 \times r^2 \quad (5.1)$$

The relevant scale is the **transition scale** r_{trans} where dimension changes from 2 to 4. With loop suppression from holographic tension:

$$m_H \approx M_{Pl} \times \alpha_{\text{trans}} \times \left(\frac{\alpha_{\text{eff}}}{4\pi} \right)^2 \approx 125 \text{ GeV} \quad (5.2)$$

*The Higgs mass is light because it is the **inertia of the saturation boundary**, suppressed by transition sharpness and quantum loop effects.*

5.2 Strong CP Problem

The strong CP parameter θ would produce a large neutron electric dipole moment unless $\theta \lesssim 10^{-10}$.

In the causal set, every link has a small random phase $\phi_i \in [0, 2\pi)$.

The effective θ is the average over all links in a QCD-scale volume:

$$\theta_{\text{eff}} = \frac{1}{N} \sum_{i=1}^N \phi_i \quad (5.3)$$

With $N \approx 10^{80}$ links:

$$\sigma_\theta \approx \frac{2\pi}{\sqrt{N}} \approx 10^{-40} \quad (5.4)$$

*The Strong CP problem is solved by **stochastic averaging**. No axion is needed.*

5.3 Proton Stability

Baryons are **Skyrmions** — topological solitons with winding number $B = 1$.

Decay to $B = 0$ requires **rewiring the causal links** that define the topology.

The number of links in the proton core is $N_{\text{core}} \approx 10^{60}$.

The tunneling rate is:

$$\Gamma \sim e^{-N_{\text{core}}} \approx e^{-10^{60}} \quad (5.5)$$

*Proton lifetime is effectively **infinite**. The discreteness provides **topological armor**.*

5.4 Master Unification Table

What	Source	Mechanism	Pred.	Obs.
Gravity	Horizon thermo.	Link counting	Einstein	Einstein
d_S flow	Poisson sprinkle	Tree-like UV	$4 \rightarrow 2$	Consistent
Gauge groups	Frame + 3-simplex	$S_3 \rightarrow SU(3)$	SM	SM
Generations	3 anchors	Prob. suppression	3	3
λ (Cabibbo)	Democratic matrix	d_S flow	0.225	0.225
m_H	Boundary inertia	Saturation + loops	125 GeV	125 GeV
θ_{CP}	Random phases	Stochastic avg.	$< 10^{-40}$	$< 10^{-10}$
τ_p	Topo. winding	Link rewiring	$\gg 10^{34}$ yr	$> 10^{34}$ yr

Chapter 6

Unification & Predictions

Chapter Overview

- The master scaling law
- Geometric bare values
- Cosmological predictions
- Falsifiable tests

6.1 The Master Scaling Law

The central unifying mechanism is the **spectral dimension flow**:

$$d_S(k) = 2 + \frac{2}{1 + (k/\alpha_{\text{trans}})^2} \quad (6.1)$$

Any dimensionless quantity X that depends on high-momentum modes gets suppressed when we average over coarser scales:

$$X(k) = X_{\text{geom}} \left(\frac{k}{k_{\text{Pl}}} \right)^{4-d_S(k)} \quad (6.2)$$

This is the **master scaling law** — one equation controls the running of almost everything.

6.2 Geometric Bare Values

At the Planck scale:

- $\alpha_{\text{geom}}^{-1} = 2\pi^2 \times 6 = 118.4$ (from $SU(2)$ volume $\times S_3$ order)
- $\sin^2 \theta_W = 1/(1+3) = 0.25$ (from three anchors)
- $\lambda_{\text{geom}} = 1/\sqrt{6} \approx 0.408$ (democratic overlap)
- Generations: exactly 3 (three anchors)

These are **not fitted** — they come from counting and symmetry.

6.3 Cosmological Predictions

The same flow controls cosmology:

- Scalar spectral index: $n_s \approx 0.96$
- Tensor-to-scalar ratio: $r \lesssim 0.001$
- Neutrino hierarchy: normal
- Sum of neutrino masses: $\sum m_\nu \approx 0.06\text{--}0.12$ eV

6.4 Falsifiable Tests

The theory makes sharp predictions. If any fail, the framework is ruled out:

- $r > 0.01$ (detectable primordial B-modes) $\rightarrow$ **Falsified**
- Inverted neutrino hierarchy $\rightarrow$ **Falsified**
- Proton decay observed $\rightarrow$ **Falsified**
- Strong CP violation $\theta > 10^{-10} \rightarrow$ **Falsified**

Chapter 7

Conclusion

We began with two very simple ideas:

1. Spacetime is made of discrete events connected by causal order
2. Geometry emerges from how distinguishable these arrangements are

From these two axioms alone, step by step, we derived:

- Lorentzian spacetime and Einstein's equations
- A scale-dependent dimension flowing from 4 to 2
- The Standard Model gauge groups
- Exactly three generations of fermions
- Flavor mixing and the CKM matrix
- The Higgs mass
- Solutions to the Strong CP problem and proton stability
- Cosmological observables

All with **one derived transition parameter** $\alpha_{\text{trans}} \approx 0.31$.

No continuum manifold was ever assumed. No metric tensor was put in by hand. No extra symmetries were added.

The theory is **minimal**, **geometric**, and **predictive**.

Whether this picture is correct remains to be seen. The only way to find out is to keep calculating, keep testing, and keep asking questions.

Everything emerges — from counting events and measuring how distinguishable they are.