

Tepotinib in a Patient With Advanced Non-Small Cell Lung Cancer Harboring *MET* Exon 14 Skipping Undergoing Concomitant Hemodialysis for Renal Failure: A Case Report

Moritz Kleemiss,¹ Christa E. Müller,² Marion Schneider,² Rainer Strotmann,³
Katrin Orlowski,⁴ Kosalaram Goteti,⁵ Mert Yanik,⁴ Peter Brossart,¹
Franz-Georg Bauernfeind¹

Clinical Practice Points

- Tepotinib, a potent, highly selective, once-daily, oral *MET* tyrosine kinase inhibitor, is approved in multiple countries for the treatment of advanced/metastatic *MET*ex14 skipping NSCLC and can be used in patients with mild-moderate renal impairment without dose adjustment. However, data in severe renal impairment are lacking.
- We report the feasibility of using the standard dose of tepotinib (500 mg; 450 mg active moiety) in a patient with advanced *MET*ex14 skipping NSCLC with end-stage renal disease undergoing hemodialysis, who attained disease control, with only mild adverse events that did not necessitate dose adjustment.
- Use of the standard dose was supported by tepotinib plasma concentration measurements, which fell within the expected range for a typical patient with NSCLC predicted using a population pharmacokinetic model and indicated no clinically relevant drug loss during dialysis.

Clinical Lung Cancer, Vol. 000, No. xxx, 1–4 © 2024 The Authors. Published by Elsevier Inc.

This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)

Keywords: *MET* inhibitor, Severe renal impairment, End stage renal disease (ESRD), Pharmacokinetics (PK), Mesenchymal-epithelial transition factor

Introduction

Lung cancer is the leading cause of cancer-related death worldwide,¹ with non-small cell lung cancer (NSCLC) accounting for >80–85% of cases.² In patients with lung cancer, renal impairment is a common comorbidity estimated to affect 28% of patients,

with severe renal impairment in 2%.³ Furthermore, a ~1.5-fold increase in lung cancer risk has been reported after kidney transplant.⁴ Chronic kidney disease (CKD) and end-stage renal disease (ESRD) requiring hemodialysis are poor prognostic factors in lung cancer.^{5,6} However, these patients are typically excluded from clinical trials of anti-cancer agents,^{7,8} partly due to the potential impact of renal dysfunction on efficacy and/or safety related to changes in pharmacokinetics, which are observed with many chemotherapeutic drugs.^{9,10} For example, decreased renal excretion can increase drug exposure and thereby impact on safety, while drug losses during hemodialysis can decrease exposure and thereby compromise efficacy.^{10,11} Due to their frequent exclusion from trials, patients with renal insufficiency have fewer anti-cancer treatment options.

Several oncogenic driver alterations have been identified in NSCLC and a better understanding of their role in tumor biology has allowed the development of new targeted therapies that have changed the treatment landscape.¹² Mesenchymal–epithelial transition factor (*MET*) exon 14 (*MET*ex14) skipping is a primary oncogenic driver reported in 3%–4% of patients with NSCLC,

Abbreviations: CLcr, creatinine clearance; CKD, chronic, kidney disease; ESRD, end-stage renal disease; *MET*, mesenchymal–epithelial transition factor; *MET*ex14, *MET* exon 14 skipping; NCCN, National Comprehensive Cancer Network; NSCLC, non-small cell lung cancer; PAD, peripheral arterial disease; PopPK, population pharmacokinetics; QD, once daily; SD, standard deviation; TKI, tyrosine kinase inhibitor.

¹Department of internal Medicine III, University Hospital Bonn, Bonn, Germany

²PharmaCenter Bonn, Pharmaceutical Institute, Pharmaceutical & Medicinal Chemistry, University of Bonn

³Qualitative Pharmacology, the healthcare business of Merck KGaA, Darmstadt, Germany

⁴Merck Healthcare Germany GmbH, Weiterstadt, Germany, an affiliate of Merck KGaA, Darmstadt, Germany

⁵Quantitative Pharmacology, EMD Serono, Rockland, MA

Submitted: Feb 28, 2024; Revised: May 17, 2024; Accepted: May 24, 2024; Epub: xxx

Address for correspondence: Moritz Kleemiss, Staatsexamen, Department of internal Medicine III, University Hospital Bonn, Venusberg-Campus 1, 53127 Bonn, Germany
E-mail contact: moritz.kleemiss@ukbonn.de

Tepotinib in a Patient With Advanced Non-Small Cell Lung Cancer

most of whom are elderly.¹³ Although *MET*ex14 skipping confers a poor prognosis with standard-of-care therapy,¹⁴ it also sensitizes tumors to *MET* tyrosine kinase inhibitors (TKIs).^{15–18} Tepotinib, a potent, highly selective, once-daily, oral *MET* TKI,¹⁹ is approved in multiple countries for the treatment of advanced/metastatic *MET*ex14 skipping NSCLC,²⁰ based on the VISION study.^{16,21,22} Clinical practice guidelines, including the National Comprehensive Cancer Network (NCCN) Clinical Practice Guidelines in Oncology (NCCN Guidelines®), American Society of Clinical Oncology and Ontario Health (Cancer Care Ontario) joint guidelines, and European Society for Medical Oncology guidelines, recommend *MET* TKIs, including tepotinib and capmatinib, for eligible patients with *MET*ex14 skipping.^{23–25}

Tepotinib can be used without dose adjustment in patients with mild-moderate renal impairment.²⁶ This practice is based on population pharmacokinetic (PopPK) modelling of data from the VISION trial, which showed no clinically meaningful effect of mild-moderate renal impairment (creatinine clearance [CLCr] 30 to 89 mL/min) on tepotinib exposure.^{26,27} Tepotinib has not been studied in patients with severe renal impairment (CLCr <30 mL/min).^{20,26} However, given its predominantly biliary elimination,²⁸ any impact of severe renal impairment on tepotinib pharmacokinetics is expected to be minimal.^{10,11} Furthermore, despite its relatively low molecular weight, tepotinib may be protected from removal during hemodialysis by its high protein binding and large volume of distribution.^{10,11,28} Nonetheless, clinical evidence in patients with severe renal impairment is needed.

We report on the efficacy, safety, and pharmacokinetics of tepotinib in a patient with *MET*ex14 skipping NSCLC with comorbid ESRD (residual estimated glomerular filtration rate of <15 mL/min), who was undergoing hemodialysis.

Case Report

A 73-year-old Caucasian female with a 40 pack-year smoking history was diagnosed with squamous NSCLC in January 2021. The medical history included: ESRD due to polycystic kidney disease, for which hemodialysis had been conducted three times a week (since 2005); kidney transplant (in 2014), which had been followed by graft failure and transplant nephrectomy; squamous cell carcinoma of the skin (completely resected in December 2020); hypertension; and peripheral arterial disease (PAD), for which an aortobifemoral Y-prosthesis had been implanted (in 2012). Residual estimated glomerular filtration rate was <15 mL/min, associated metabolic alterations such as hyperphosphatemia and acid-base imbalances were well controlled with medication, and hepatic function was normal.

At NSCLC diagnosis, the patient had clinical stage IVB disease (cT4 cN0 cM1c) with bone, pleural and lung metastasis, and Eastern Cooperative Oncology Group performance status of 1. Next-generation sequencing analysis of a tissue biopsy (QIASEQ® Targeted DNA Custom Panel, Qiagen, Venlo, The Netherlands) revealed a *MET*ex14 skipping mutation (c.3017-3028+26del) with concomitant intermediate-level *MET* amplification (gene copy number 4.95). The programmed death ligand-1 tumor proportion score was 90% by immunohistochemistry (PD-L1 IHC 22C3 pharmDx assay, Agilent Technologies, Santa Clara, CA, USA). First-

line therapy with pembrolizumab (200 mg, every 3 weeks) was initiated in January 2021 and continued without significant adverse events until disease progression in May 2021. The patient also received palliative radiation (39 Gy, 13 fractions) to the thoracic vertebrae (T4-T6) for pain from osteolytic bone metastasis.

Based on the *MET*ex14 skipping alteration, second-line therapy with tepotinib was initiated through a compassionate use program in Germany in July 2021, at which time tepotinib was not yet approved in the European Union. Tepotinib was administered at the standard dose of 500 mg (450 mg active moiety), orally, once daily in the morning with food, under conditions of careful safety and pharmacokinetic monitoring to confirm the appropriateness of the dose in this patient receiving hemodialysis for ESRD. The first response assessment showed shrinkage of the primary tumor (from 6.1 × 4.3 cm to 4.6 × 3.0 cm) and lung metastases, with a reduction in sum of lesion diameters <30%. The best overall response to tepotinib was stable disease, and the patient reported improvements in mobility, physical function, and pain. Grade 1 fatigue and Grade 2 peripheral edema were observed but did not necessitate dose adjustment. Tepotinib was continued for a total of 5.5 months until December 2021, when the patient died due to PAD with concurrent cerebral hemorrhage. Prior to death, a second tumor assessment had shown maintenance of the primary tumor shrinkage with further reductions in the lung metastases, indicating that the NSCLC was well controlled.


During tepotinib treatment, hemodialysis was performed 3 times per week using a Gambro D396 machine (Gambro Lundia AB, Deerfield, IL, USA) with an ELISIO® 19H dialyzer (Nipro, Mechelen, Belgium), at a blood flow of 300 mL/min, over 4 hours and 10 minutes. The first sessions were conducted on treatment Days 5, 7, and 9, approximately 3–4 hours after tepotinib administration. Tepotinib was taken at the same time of day on dialysis and non-dialysis days. Plasma samples were collected pre- and post-dialysis, approximately 1 hour and 8 hours and 25 minutes after tepotinib administration, respectively. Tepotinib was assayed in plasma samples by liquid chromatography–tandem mass spectrometry at the Department of Pharmaceutical and Medicinal Chemistry, University of Bonn (Germany), with 4 technical bioanalysis replicates per sample.

While variability between technical replicates was within the acceptable range, there was some between-day variability in the pre-dialysis concentrations of tepotinib (Figure 1). Based on nominal sampling times, all observed tepotinib plasma concentrations were close to the expected median plasma concentrations, as predicted by the PopPK model for a typical patient with NSCLC receiving this tepotinib dose-regimen.²⁷ The observed post-dialysis plasma concentrations of tepotinib were 16.8–24.2% lower than the pre-dialysis plasma concentrations but fell within the 90% prediction interval for patients receiving tepotinib without dialysis.

Discussion

Tepotinib is approved to treat advanced/metastatic *MET*ex14 skipping NSCLC and can be used in patients with mild-moderate renal impairment without dose adjustment; however, data in severe renal impairment have been lacking.^{20,26,27} The present case demonstrates, for the first time, the successful use of tepotinib in

Figure 1 Mean plasma concentration of tepotinib at Days 5, 7, and 9, recorded pre- and post-dialysis and plotted alongside the median and 90% prediction interval for the tepotinib concentration–time profile for typical patients treated with 500 mg tepotinib QD as predicted from the tepotinib PopPK model. Data points indicate the mean and SD of 4 technical replicates. PopPK, population pharmacokinetic; QD, once daily; SD, standard deviation.



a patient with ESRD undergoing hemodialysis. With the standard tepotinib dose (500 mg), the patient attained disease control lasting >5 months, with only mild adverse events, which was accompanied by improvements in mobility, physical function, and pain and, therefore, quality of life. The patient ultimately died from an NSCLC-unrelated cause without disease progression.

In patients with ESRD managed with hemodialysis, anti-cancer drug exposure can either be increased due to impaired renal elimination or decreased due to enhanced systemic clearance resulting from drug loss during hemodialysis.¹⁰ In the present case, the first hemodialysis was conducted on treatment Day 5, at which time tepotinib should almost have reached steady state (given its terminal elimination half-life of 32 hours).^{26,28} The pre-dialysis tepotinib concentration on this day was in line with expectations for a typical patient without ESRD, indicating no substantial increase in tepotinib exposure secondary to renal impairment. The limited impact of severe renal impairment on tepotinib clearance is consistent with the minimal contribution of renal excretion to the total elimination of tepotinib, with 85% of tepotinib excreted in feces and only 13.6% in urine.^{26,28} Furthermore, on Days 5, 7, and 9, pre- and post-dialysis concentrations showed no clinically relevant loss of tepotinib during hemodialysis, which was likely protected from removal by its high plasma protein binding (98%) and large volume of distribution (574 L after an intravenous tracer micro-dose).²⁸ Overall, observed plasma concentrations were within the 90% prediction interval for patients receiving tepotinib without dialysis, thereby suggesting the patient achieved efficacious exposure (Figure 1). The pharmacokinetic data provide a rationale for the

observed efficacy and tolerability of tepotinib, which did not require dose adjustments to manage adverse events, and provided disease control.

Conclusion

This case report illustrates the feasibility of using the standard dose of tepotinib in a patient with advanced *MET*ex14 skipping NSCLC with ESRD undergoing hemodialysis, and provides plasma concentration data supporting its use without dose adjustment in this setting. These data may be applicable to other patients with severe renal impairment.

Consent

The patient gave written informed consent for de-identified data to be disclosed and published in any scientific journal for educational purposes.

Funding Support

None.

Disclosure

M Kleemiss and FG Bauernfeind have received lecture fees from the healthcare business of Merck KGaA, Darmstadt, Germany. R Strotman, K Orłowski and M Yanik are employees of the healthcare business of Merck KGaA, Darmstadt, Germany. K Goteti is an employee of EMD Serono.

Tepotinib in a Patient With Advanced Non-Small Cell Lung Cancer

CRedit authorship contribution statement

Moritz Kleemiss: Formal analysis, Investigation, Methodology, Writing – review & editing. **Christa E. Müller:** Investigation, Writing – review & editing. **Marion Schneider:** Investigation, Writing – review & editing. **Rainer Strotmann:** Conceptualization, Formal analysis, Methodology, Writing – review & editing. **Katrin Orłowski:** Conceptualization, Formal analysis, Writing – review & editing. **Kosalaram Goteti:** Conceptualization, Formal analysis, Methodology, Writing – review & editing. **Mert Yanik:** Conceptualization, Formal analysis, Writing – review & editing. **Peter Brossart:** Investigation, Writing – review & editing. **Franz-Georg Bauernfeind:** Investigation, Writing – review & editing.

Acknowledgments

The authors would like to thank the patient and the patient's family. C.E.M. would like to thank the Deutsche Forschungsgemeinschaft (DFG) and the State of Northrhine-Westfalia for financing the sensitive triple-quadrupole/linear ion-trap mass spectrometer (SCIEX QTRAP 6500+) that was used for the analytical measurements (INST 217/1058-1 FUGG, project number 453027627). Medical writing assistance, funded by the healthcare business of Merck KGaA, Darmstadt, Germany (CrossRef Funder ID: 10.13039/100009945), was provided by Moon Jain, PhD, employee of Syneos Health, UK.

References

- Sung H, Ferlay J, Siegel RL, et al. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin*. 2021;71(3):209–249. doi:10.3322/caac.21660.
- Schabath MB, Cote ML. Cancer progress and priorities: lung cancer. *Cancer Epidemiol Biomarkers Prev*. 2019;28(10):1563–1579. doi:10.1158/1055-9965.EPI-19-0221.
- Grose D, Morrison DS, Devereux G, et al. Comorbidities in lung cancer: prevalence, severity and links with socioeconomic status and treatment. *Postgrad Med J*. 2014;90(1064):305–310. doi:10.1136/postgradmedj-2013-132186.
- Engels EA, Pfeiffer RM, Fraumeni JF, et al. Spectrum of cancer risk among US solid organ transplant recipients. *JAMA*. 2011;306(17):1891–1901. doi:10.1001/JAMA.2011.1592.
- Chien CC, Han MM, Chiu YH, et al. Epidemiology of cancer in end-stage renal disease dialysis patients: a national cohort study in Taiwan. *J Cancer*. 2017;8(1):9–18. doi:10.7150/jca.16550.
- Wei YF, Chen JY, Lee HS, Wu JT, Hsu CK, Hsu YC. Association of chronic kidney disease with mortality risk in patients with lung cancer: a nationwide Taiwan population-based cohort study. *BMJ Open*. 2018;8(1):e019661. doi:10.1136/bmjopen-2017-019661.
- Kitchlu A, Shapiro J, Amir E, et al. Representation of patients with chronic kidney disease in trials of cancer therapy. *JAMA*. 2018;319(23):2437–2439. doi:10.1001/JAMA.2018.7260.
- Sprangers B, Perazella MA, Lichtman SM, Rosner MH, Jhaveri KD. Improving cancer care for patients with CKD: the need for changes in clinical trials. *Kidney Int Rep*. 2022;7(9):1939–1950. doi:10.1016/j.EKIR.2022.06.005.
- Launay-Vacher V, Janus N, Deray G. Renal insufficiency and cancer treatments. *ESMO Open*. 2016;1(4):e000091. doi:10.1136/esmoopen-2016-000091.
- Krens SD, Lassche G, Jansman FGA, et al. Dose recommendations for anticancer drugs in patients with renal or hepatic impairment. *Lancet Oncol*. 2019;20(4):e200–e207. doi:10.1016/S1470-2045(19)30145-7.
- Hashimoto M, Maeda H, Oniki K, et al. New insight concerning therapeutic drug monitoring—the importance of the concept of psychonephrology. *Biol Pharm Bull*. 2022;45(7):834–842. doi:10.1248/BPBB.22-00025.
- Chevallier M, Borgeaud M, Addeo A, Friedlaender A. Oncogenic driver mutations in non-small cell lung cancer: past, present and future. *World J Clin Oncol*. 2021;12(4):217–237. doi:10.5306/wjco.v12.i4.217.
- Hong L, Zhang J, Heymach J V, Le X. Current and future treatment options for MET exon 14 skipping alterations in non-small cell lung cancer. *Ther Adv Med Oncol*. 2021;13:1758835921992976. doi:10.1177/1758835921992976.
- Tong JH, Yeung SF, Chan AWH, et al. MET amplification and exon 14 splice site mutation define unique molecular subgroups of non-small cell lung carcinoma with poor prognosis. *Clin Cancer Res*. 2016;22(12):3048–3056. doi:10.1158/1078-0432.CCR-15-2061.
- Paik PK, Drilon A, Fan PD, et al. Response to MET inhibitors in patients with stage IV lung adenocarcinomas harboring MET mutations causing exon 14 skipping. *Cancer Discov*. 2015;5(8):842–849. doi:10.1158/2159-8290.CD-14-1467.
- Mazieres J, Paik PK, Garassino MC, et al. Tepotinib treatment in patients with MET Exon 14–skipping non-small cell lung cancer: long-term follow-up of the VISION phase 2 nonrandomized clinical trial. *JAMA Oncol*. 2023;9(9):1260–1266. doi:10.1001/JAMAONCOL.2023.1962.
- Wolf J, Seto T, Han JY, et al. Capmatinib in MET exon 14–mutated or MET-amplified non-small-cell lung cancer. *N Engl J Med*. 2020;383(10):944–957. doi:10.1056/NEJMoa2002787.
- Lu S, Fang J, Li X, et al. Once-daily savolitinib in Chinese patients with pulmonary sarcomatoid carcinomas and other non-small-cell lung cancers harbouring MET exon 14 skipping alterations: a multicentre, single-arm, open-label, phase 2 study. *Lancet Respir Med*. 2021;9(10):1154–1164. doi:10.1016/S2213-2600(21)00084-9.
- Falchook GS, Kurzrock R, Amin HM, et al. First-in-man Phase I trial of the selective MET inhibitor tepotinib in patients with advanced solid tumors. *Clin Cancer Res*. 2020;26(6):1237–1246. doi:10.1158/1078-0432.CCR-19-2860.
- Mathieu LN, Larkins E, Akinboro O, et al. FDA approval summary: capmatinib and tepotinib for the treatment of metastatic NSCLC harboring MET Exon 14 skipping mutations or alterations. *Clin Cancer Res*. 2022;28(2):249–254. doi:10.1158/1078-0432.CCR-21-1566.
- Paik P, Felip E, Veillon R, et al. Tepotinib in non-small-cell lung cancer with MET exon 14 skipping mutations. *N Engl J Med*. 2020;383(10):931–943. doi:10.1056/NEJMoa2004407.
- Le X, Sakai H, Felip E, et al. Tepotinib efficacy and safety in patients with MET exon 14 skipping NSCLC: outcomes in patient subgroups from the VISION study with relevance for clinical practice. *Clin Cancer Res*. 2022;28(6):1117–1126. doi:10.1158/1078-0432.CCR-21-2733.
- Hendriks LE, Kerr KM, Menis J, et al. Oncogene-addicted metastatic non-small-cell lung cancer: ESMO Clinical Practice Guideline for diagnosis, treatment and follow-up. *Ann Oncol*. 2023;34(4):339–357. doi:10.1016/j.annonc.2022.12.009.
- Hanna NH, Robinson AG, Temin S, et al. Therapy for stage IV non-small-cell lung cancer with driver alterations: ASCO and OH (CCO) joint guideline update. *J Clin Oncol*. 2021;39(9):1040–1091 [Erratum: J Clin Oncol 2021;39(22):2520]. doi:10.1200/JCO.20.03570.
- NCCN Clinical Practice Guidelines in Oncology. Non-Small Cell Lung Cancer Version 3.2024. Accessed March 12, 2023. https://www.nccn.org/professionals/physician_gls/pdf/nscl.pdf.
- US Food and Drug Administration. *TEPMETKO (Tepotinib) Prescribing Information*; 2023. https://www.accessdata.fda.gov/drugsatfda_docs/label/2023/214096s001s002lbl.pdf.
- Xiong W, Papasouliotis O, Niclas JE, Strotmann R, Girard P. Population pharmacokinetic analysis of tepotinib, an oral MET kinase inhibitor, including data from the VISION study. *Cancer Chemother Pharmacol*. 2022;89(5):655–669. doi:10.1007/s00280-022-04423-5.
- Johne A, Scheible H, Becker A, van Lier JJ, Wolna P, Meyring M. Open-label, single-center, phase I trial to investigate the mass balance and absolute bioavailability of the highly selective oral MET inhibitor tepotinib in healthy volunteers. *Invest New Drugs*. 2020;38(5):1507–1519. doi:10.1007/s10637-020-00926-1.