

Trigonometric Coefficients for Regular S-Sided Polygons

Eli Ruminer

Table of Contents

1	Introduction	2
2	Trigonometric Equations	2
2.1	Identities	3
3	Polygonal Equations	4
3.1	Intersections	5
4	Proof	5

1 Introduction

This paper creates and proves a function that finds the distance between the center of a regular n-sided polygon and a point on its perimeter at a given angle without knowing that point's position. This when multiplied by sine or cosine allows finding those points on the perimeter of the polygon. This allows for the creation of a general equation for any regular n-sided polygon similar to the equation of a circle. This equation can then be used to find areas of intersection between polygons.

Following this, the modified sine and cosine functions are then proved to always generate a point on the perimeter of an s-sided polygon, given s and an angle.

2 Trigonometric Equations

Let $s = \{x : x \in \mathbb{Z}^+ \text{ and } x \geq 3\}$, and

$$\begin{aligned} d_s(\theta) &= \cos\left(\frac{\pi}{s}\right) \sec\left(\frac{\pi - \arccos(\cos(s\theta))}{s}\right) \\ \cos_s(\theta) &= d_s(\theta) \cos(\theta) \\ \sin_s(\theta) &= d_s(\theta) \sin(\theta) \end{aligned}$$

This is derived through the original functions:

$$\begin{aligned} h_s(\theta) &= \sin\left(\frac{\pi}{2} - \frac{\pi}{s} + (\theta \bmod \frac{2\pi}{s})\right) - \sin\left(\frac{\pi}{2} - \frac{\pi}{s}\right) \\ d_s(\theta) &= \frac{h_s(\theta)}{\sin\left(\frac{\pi}{2} - \frac{\pi}{s} + (\theta \bmod \frac{2\pi}{s})\right)} \\ \cos_s(\theta) &= (1 - d_s(\theta)) \cos(\theta) \\ \sin_s(\theta) &= (1 - d_s(\theta)) \sin(\theta) \end{aligned}$$

These functions follow the premise that any s-sided regular polygon (P_s) centered at (0,0) has a circumcircle where each vertex touches the circumference of the circumcircle.

Arc a can be constructed which is $\frac{1}{s^{\text{th}}}$ of the circumcircle, its diameter is the same length as any line which constructs P_s , and it's circumference equals to the circumference of the section of the circumcircle spanning from adjacent vertices on P_s .

Thus, the height of a at a given point on a circle subtracted from the radius of the circle will result in the distance from the center of P_s to the point on the perimeter of P_s which intersects the ray of angle θ drawn in standard position.

$h_s(\theta)$ serves to find the height of a at a position by splitting the circumcircle into s^{th} s (resulting in a period of $\frac{2\pi}{s}$), finding the height of a point on a through sine and changing from the "left" vertex to the "right" vertex over the period of $\frac{2\pi}{s}$ (the length of a). The use of mod makes the function periodic. It then subtracts by the height of a vertex so that

$h_s(\theta) = 0$ when $\theta = 0$

$d_s(\theta)$ divides the height, $h_s(\theta)$, by the angle between the assumed intersection point on P_s and the intersection point on the circumcircle to get the actual distance.

These functions then simplify to:

$$\begin{aligned} d_s(\theta) &= \cos\left(\frac{\pi}{s}\right) \sec\left(\frac{\pi}{s} - (\theta \bmod \frac{2\pi}{s})\right) \\ \cos_s(\theta) &= d_s(\theta) \cos(\theta) \\ \sin_s(\theta) &= d_s(\theta) \sin(\theta) \end{aligned}$$

Which is not algebraic due to modulo, yet it can be noted that due to the nature of the original $h_s(\theta)$, $\theta \bmod \frac{2\pi}{s}$ can be substituted for an equivalent function which has an equal period of $\frac{2\pi}{s}$, increases linearly from 0 to $\frac{\pi}{s}$ over $[0, \frac{\pi}{s}]$ and decreases linearly at the same rate over $[\frac{\pi}{s}, \frac{2\pi}{s}]$.

These requirements can be met by $\frac{\arccos(\cos(n\theta))}{n}$ ¹, which is periodic over $\frac{2\pi}{s}$, has a domain of $n\theta \in \mathbb{R}$, and thus $\theta \in \mathbb{R}$, and a range of $[0, \frac{\pi}{s}]$, thus the above equations can be changed to:

$$\begin{aligned} d_s(\theta) &= \cos\left(\frac{\pi}{s}\right) \sec\left(\frac{\pi - \arccos(\cos(s\theta))}{s}\right) \\ \cos_s(\theta) &= d_s(\theta) \cos(\theta) \\ \sin_s(\theta) &= d_s(\theta) \sin(\theta) \end{aligned}$$

2.1 Identities

Identity 1.1 Let $n = \mathbb{Z}$:

$$\begin{aligned} \cos_s\left(\theta + n \frac{2\pi}{s}\right) &= d_s(\theta) \cos\left(\theta + n \frac{2\pi}{s}\right) \\ \sin_s\left(\theta + n \frac{2\pi}{s}\right) &= d_s(\theta) \sin\left(\theta + n \frac{2\pi}{s}\right) \end{aligned}$$

Identity 1.2 When $0 \leq \theta \leq \frac{2\pi}{s}$ then $d_s(\theta) = \cos\left(\frac{\pi}{s}\right) \sec\left(\frac{\pi}{s} - \theta\right)$

When θ only spans one period, $\frac{\arccos(\cos(s\theta))}{s}$ is not required to make the function periodic.

Identity 1.3 Let $n = \mathbb{Z}$ then:

$$\begin{aligned} \cos_s\left(n \frac{2\pi}{s}\right) &= (1) \cos\left(n \frac{2\pi}{s}\right) = \cos\left(n \frac{2\pi}{s}\right) \\ \sin_s\left(n \frac{2\pi}{s}\right) &= (1) \sin\left(n \frac{2\pi}{s}\right) = \sin\left(n \frac{2\pi}{s}\right) \end{aligned}$$

This is true because $\arccos(\cos(s\theta))$ evaluates to 0 whenever $\theta = n \frac{2\pi}{s}$, which means $\sec\left(\frac{\pi - \arccos(\cos(s\theta))}{s}\right)$ simplifies to $\sec\left(\frac{\pi}{s}\right)$ so $d_s\left(n \frac{2\pi}{s}\right) = \sec\left(\frac{\pi}{s}\right) \cos\left(\frac{\pi}{s}\right) = 1$

¹J. L. Spence, "Periodic absolute value functions".

3 Polygonal Equations

From these formulas, the equation for any regular polygon can be derived. The equation is found through a mutation of the ellipse equation by $d_s(\theta)$:

$$\frac{(x-h)^2}{a^2} + \frac{(y-k)^2}{b^2} = d_s(\theta)^2 * 1$$

To derive theta, one would start with $\theta = \arctan(\frac{(y-k)a}{(x-h)b})$, which would find the angle at the correct point offset from the center, but due to x being squared, the shape is mirrored over $x = h$, which will work for even sided polygons as they are symmetric over $x = h$, but not for odd sided polygons due to their anti-symmetry (this is also the case over the line $y = k$ but all shapes, regardless of side parity, are mirrored over $y = k$). This can be fixed by noting that the angle is calculated correctly in Quadrants I and IV (relative to (h, k) as the origin), and π radians off from the expected angle in Quadrants II and III. Thus when $x < h$, π must be added to θ . Using the sign function and modifying it to equal 1 when $x < h$ and 0 when $x \geq h$ results in the formula:

$$-\frac{1}{2}(\frac{x-h}{|x-h|} - 1)$$

which when multiplied by π and added to θ becomes:

$$\arctan(\frac{(y-k)a}{(x-h)b}) - \frac{\pi}{2}(\frac{x-h}{|x-h|} - 1)$$

meaning that the standard equation is:

$$\frac{(x-h)^2}{a^2} + \frac{(y-k)^2}{b^2} = d_s(\arctan(\frac{(y-k)a}{(x-h)b}) - \frac{\pi}{2}(\frac{x-h}{|x-h|} - 1))^2$$

As d_s takes an angle, a constant value can be added to θ to make any shape rotate, allowing for easy rotation calculations.

3.1 Intersections

Let the regular Polygon P_s with s sides (where $(s \in \mathbb{Z} \text{ and } s \geq 3)$) be represented by the equation

$$P_s(x, y) = \frac{(x-h)^2}{a^2} + \frac{(y-k)^2}{b^2} - d_s(\arctan(\frac{(y-k)a}{(x-h)b}) - \frac{\pi}{2}(\frac{x-h}{|x-h|} - 1))^2$$

A point (x, y) is on the perimeter of P_s if $P_s(x, y) = 0$, inside the area of P_s if $P_s(x, y) < 0$, and outside the area of P_s if $P_s(x, y) > 0$.

Thus (x, y) is inside the intersection of two polygons P_{s_1} and P_{s_2} if $P_{s_1}(x, y) < 0$ and $P_{s_2}(x, y) < 0$. This can be made into one inequality of:

$$|(P_{s_1}(x, y) - |P_{s_1}(x, y)||)(P_{s_2}(x, y) - |P_{s_2}(x, y)||)| > 0$$

More generally for any set of Polygons C , a point (x, y) is inside the intersection of all polygons in the set (the "addition" of the polygons) if

$$\left| \prod_{i=1}^{|C|} (C_i(x, y) - |C_i(x, y)|) \right| > 0$$

4 Proof

Let $n, s = \mathbb{Z}, s \geq 3$. Shape P_s is made of s vertices and s line segments with the n^{th} vertex (n_i) at the point $(\cos(n \frac{2\pi}{s}), \sin(n \frac{2\pi}{s}))$. The n^{th} line segment, L_n , spans between v_n and v_{n+1} and is represented by the equation

$$L_n = \frac{\sin((n+1) \frac{2\pi}{s}) - \sin(n \frac{2\pi}{s})}{\cos((n+1) \frac{2\pi}{s}) - \cos(n \frac{2\pi}{s})} x + \sin(n \frac{2\pi}{s}) - \frac{\sin((n+1) \frac{2\pi}{s}) - \sin(n \frac{2\pi}{s})}{\cos((n+1) \frac{2\pi}{s}) - \cos(n \frac{2\pi}{s})} \cos(n \frac{2\pi}{s})$$

which simplifies to

$$L_n = -\cot((2n+1) \frac{\pi}{s}) x + \sin(n \frac{2\pi}{s}) + \cot((2n+1) \frac{\pi}{s}) \cos(n \frac{2\pi}{s})$$

Let $0 \leq a < b \leq \frac{2\pi}{s}$, $\alpha = a + n \frac{2\pi}{s}$, and $\beta = b + n \frac{2\pi}{s}$: a "modified" line can be drawn with the equation $\sin_s(\theta)$ and $\cos_s(\theta)$ which would be written as

$$y_m = \frac{\sin_s(\beta) - \sin_s(\alpha)}{\cos_s(\beta) - \cos_s(\alpha)} x + \sin_s(\alpha) - \frac{\sin_s(\beta) - \sin_s(\alpha)}{\cos_s(\beta) - \cos_s(\alpha)} \cos_s(\alpha)$$

the slope of y_m can be expanded to

$$\frac{\sin(\beta)(\cos(a) + \tan(\frac{\pi}{s}) \sin(a)) - \sin(\alpha)(\cos(b) + \tan(\frac{\pi}{s}) \sin(b))}{\cos(\beta)(\cos(a) + \tan(\frac{\pi}{s}) \sin(a)) - \cos(\alpha)(\cos(b) + \tan(\frac{\pi}{s}) \sin(b))}$$

Which simplifies to

$$\frac{\sin(a-b)(\sin(n \frac{2\pi}{s}) \tan(\frac{\pi}{s}) - \cos(n \frac{2\pi}{s}))}{\sin(a-b)(\cos(n \frac{2\pi}{s}) \tan(\frac{\pi}{s}) + \sin(n \frac{2\pi}{s}))} = -\cot((2n+1) \frac{\pi}{s})$$

thus the equation of the "modified" line becomes:

$$y_m = -\cot\left((2n+1)\frac{\pi}{s}\right)x + \sin_s(\alpha) + \cot\left((2n+1)\frac{\pi}{s}\right)\cos_s(\alpha)$$

This leads to the conclusion that when $n\frac{2\pi}{s} \leq a < b \leq (n+1)\frac{2\pi}{s}$ the slopes of y_m and L_n are equal.

If $a = 0$, due to **Identity 1.3**, then for any applicable value of b , y_m becomes:

$$y_m = -\cot\left((2n+1)\frac{\pi}{s}\right)x + \sin\left(n\frac{2\pi}{s}\right) + \cot\left((2n+1)\frac{\pi}{s}\right)\cos\left(n\frac{2\pi}{s}\right)$$

Meaning that $y_m = L_n$ when $a = 0$ (denoted $y_{m,a=0}$) and due to **Identity 1.3**: $\sin_s(a + n\frac{2\pi}{s}) = \sin(n\frac{2\pi}{s})$ and $\cos_s(a + n\frac{2\pi}{s}) = \cos(n\frac{2\pi}{s})$ thus the point $A(\cos_s(a + n\frac{2\pi}{s}), \sin_s(a + n\frac{2\pi}{s}))$, is equal to v_n .

Because $y_{m,a=0} = L_n$, the point $B(\cos_s(b + n\frac{2\pi}{s}), \sin_s(b + n\frac{2\pi}{s}))$ is always on line segment L_n , thus when $a \neq 0$, A must also always be on line segment L_n .

This means that for any value x where $0 \leq x \leq \frac{2\pi}{s}$, the point $X(\cos_s(x + n\frac{2\pi}{s}), \sin_s(x + n\frac{2\pi}{s}))$, falls on line L_n of P_s , and since $n \in \mathbb{Z}$, any point $P(\cos_s(\theta), \sin_s(\theta))$, will always be on the perimeter of P_s .

References

- [1] J. L. Spence, “Periodic absolute value functions,” *School Science and Mathematics*, vol. 61, no. 9, pp. 664–666, 1961. DOI: <https://doi.org/10.1111/j.1949-8594.1961.tb08220.x>. eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1949-8594.1961.tb08220.x>. [Online]. Available: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1949-8594.1961.tb08220.x>.