



DESK No.

--	--	--

FAMILY NAME: _____

GIVEN NAMES: _____

SIGNATURE: _____

STUDENT NUMBER:

--	--	--	--	--	--	--	--

SEMESTER 2, 2021 EXAMINATIONS - ATTACHMENTS**Engineering**

Electrical, Electronic and Computer Engineering

ENSC3015**Signals and Systems**This paper contains: **12 Pages (including title page)**Time Allowed: **2:00** hours**INSTRUCTIONS:**

Please use the included tables of pairs and properties and useful formulas as required.

THIS IS A CLOSED BOOK EXAMINATION (SEE ALLOWABLE ITEMS)**SUPPLIED STATIONERY****1 x Answer Booklet 18 Pages****ALLOWABLE ITEMS****UWA Approved Calculator with Sticker
Open Book with Student Notes****PLEASE NOTE**

Examination candidates may only bring authorised materials into the examination room. If a supervisor finds, during the examination, that you have unauthorised material, in whatever form, in the vicinity of your desk or on your person, whether in the examination room or the toilets or en route to/from the toilets, the matter will be reported to the head of school and disciplinary action will normally be taken against you. This action may result in your being deprived of any credit for this examination or even, in some cases, for the whole unit. This will apply regardless of whether the material has been used at the time it is found.

Therefore, any candidate who has brought any unauthorised material whatsoever into the examination room should declare it to the supervisor immediately. Candidates who are uncertain whether any material is authorised should ask the supervisor for clarification.

Candidates must comply with the Examination Rules of the University and with the directions of supervisors.

No electronic devices are permitted during the examination.

All question papers and answer booklets are the property of the University and remain so at all times.

This page has been left intentionally blank

Zero-input Response

Real Roots (continuous-time systems)

Assume r of the N roots are real but identical, that is $\lambda_1 = \lambda_2 = \dots = \lambda_r = \lambda$ with the remaining $(N - r)$ roots being distinct, that is $\lambda_{r+1}, \lambda_{r+2}, \dots, \lambda_N$:

$$y_0(t) = (c_1 + c_2 t + \dots + c_r t^{r-1})e^{\lambda t} + c_{r+1}e^{\lambda_{r+1}t} + c_{r+2}e^{\lambda_{r+2}t} + \dots + c_N e^{\lambda_N t}$$

Real Roots (discrete-time systems)

Assume r of the N roots are real but identical ($= \lambda$), with the remaining $(N - r)$ roots being distinct:

$$y_0[n] = (c_1 + c_2 n + \dots + c_r n^{r-1})\lambda^n + \sum_{k=r+1}^N c_k \lambda_k^n$$

Convolution

$$x_1(t) * x_2(t) = \int_{-\infty}^{\infty} x_1(\tau)x_2(t - \tau)d\tau$$

$$x_1[n] * x_2[n] = \sum_{m=-\infty}^{\infty} x_1[m]x_2[n - m]$$

Partial Fraction Expansion

Simple, distinct poles, $d_i \neq d_j$ for $X(s)$ with $M < N$

$$X(s) = \sum_{k=1}^N \frac{A_k}{(s - d_k)}$$

where:

$$A_k = (s - d_k)X(s)|_{s=d_k} \quad \text{and} \quad \frac{A_k}{(s - d_k)} \leftrightarrow A_k e^{d_k t} u(t)$$

Simple, distinct poles, $d_i \neq d_j$ for $X(z)$ with $M < N$

$$X(z) = \sum_{k=1}^N \frac{A_k}{(1 - d_k z^{-1})}$$

where:

$$A_k = (1 - d_k z^{-1})X(z)|_{z=d_k} \quad \text{and} \quad \frac{A_k}{(1 - d_k z^{-1})} \leftrightarrow A_k (d_k)^n u[n]$$


Causal and Stable Systems

For a continuous-time system with $M \leq N$ **all the poles of $H(s)$ must lie in the left half of the s -plane**, or left hand plane (LHP), that is $\text{Re}(d_k) < 0$, since the ROC is to the right of all poles (causal) and the $j\omega$ -axis is included (stable). **If $M > N$ the system is deemed unstable.**

For a discrete-time system **all the poles of $H(z)$ must lie inside the unit circle**, that is $|d_k| < 1$, since the ROC is to the exterior of all poles (causal) and the unit circle is included (stable).

Direct Form II Realisation of Discrete-Time Systems

$$H(z) = \frac{Y(z)}{X(z)} = \frac{\sum_{k=0}^N B_k z^{-k}}{1 + \sum_{k=1}^N A_k z^{-k}}$$



D.1 Basic Laplace Transforms

Signal $x(t) = \frac{1}{2\pi j} \int_{\sigma-j\infty}^{\sigma+j\infty} X(s)e^{st} ds$	Transform $X(s) = \int_{-\infty}^{\infty} x(t)e^{-st} dt$	ROC
$u(t)$	$\frac{1}{s}$	$\text{Re}\{s\} > 0$
$tu(t)$	$\frac{1}{s^2}$	$\text{Re}\{s\} > 0$
$\delta(t - \tau), \tau \geq 0$	$e^{-s\tau}$	for all s
$e^{-at}u(t)$	$\frac{1}{s + a}$	$\text{Re}\{s\} > -a$
$te^{-at}u(t)$	$\frac{1}{(s + a)^2}$	$\text{Re}\{s\} > -a$
$[\cos(\omega_1 t)]u(t)$	$\frac{s}{s^2 + \omega_1^2}$	$\text{Re}\{s\} > 0$
$[\sin(\omega_1 t)]u(t)$	$\frac{\omega_1}{s^2 + \omega_1^2}$	$\text{Re}\{s\} > 0$
$[e^{-at} \cos(\omega_1 t)]u(t)$	$\frac{s + a}{(s + a)^2 + \omega_1^2}$	$\text{Re}\{s\} > -a$
$[e^{-at} \sin(\omega_1 t)]u(t)$	$\frac{\omega_1}{(s + a)^2 + \omega_1^2}$	$\text{Re}\{s\} > -a$

Signal	Bilateral Transform	ROC
$\delta(t - \tau), \tau < 0$	$e^{-s\tau}$	for all s
$-u(-t)$	$\frac{1}{s}$	$\text{Re}\{s\} < 0$
$-tu(-t)$	$\frac{1}{s^2}$	$\text{Re}\{s\} < 0$
$-e^{-at}u(-t)$	$\frac{1}{s + a}$	$\text{Re}\{s\} < -a$
$-te^{-at}u(-t)$	$\frac{1}{(s + a)^2}$	$\text{Re}\{s\} < -a$

D.2 Laplace Transform Properties

Signal	Unilateral Transform $x(t) \xleftrightarrow{\mathcal{L}_u} X(s)$ $y(t) \xleftrightarrow{\mathcal{L}_u} Y(s)$	Bilateral Transform $x(t) \xleftrightarrow{\mathcal{L}} X(s)$ $y(t) \xleftrightarrow{\mathcal{L}} Y(s)$	ROC $s \in R_x$ $s \in R_y$
$ax(t) + by(t)$	$aX(s) + bY(s)$	$aX(s) + bY(s)$	At least $R_x \cap R_y$
$x(t - \tau)$	$e^{-s\tau}X(s)$ if $x(t - \tau)u(t) = x(t - \tau)u(t - \tau)$	$e^{-s\tau}X(s)$	R_x
$e^{s_0 t}x(t)$	$X(s - s_0)$	$X(s - s_0)$	$R_x + \text{Re}\{s_0\}$
$x(at)$	$\frac{1}{a}X\left(\frac{s}{a}\right), a > 0$	$\frac{1}{ a }X\left(\frac{s}{a}\right)$	$\frac{R_x}{ a }$
$x(t) * y(t)$	$X(s)Y(s)$ if $x(t) = y(t) = 0$ for $t < 0$	$X(s)Y(s)$	At least $R_x \cap R_y$
$-tx(t)$	$\frac{d}{ds}X(s)$	$\frac{d}{ds}X(s)$	R_x
$\frac{d}{dt}x(t)$	$sX(s) - x(0^-)$	$sX(s)$	At least R_x
$\int_{-\infty}^t x(\tau) d\tau$	$\frac{1}{s} \int_{-\infty}^{0^-} x(\tau) d\tau + \frac{X(s)}{s}$	$\frac{X(s)}{s}$	At least $R_x \cap \{\text{Re}\{s\} > 0\}$

Unilateral Transform: $\frac{d^n}{dt^n}x(t) \leftrightarrow s^n X(s) - \sum_{k=1}^n s^{n-k}x^{(k-1)}(0^-)$ where $x^{(r)}(0^-)$ is $\left.\frac{d^r x}{dt^r}\right|_{t=0^-}$

E.1 Basic z-Transforms

Signal	Transform	
$x[n] = \frac{1}{2\pi j} \oint X(z) z^{n-1} dz$	$X[z] = \sum_{n=-\infty}^{\infty} x[n] z^{-n}$	ROC
$\delta[n]$	1	All z
$u[n]$	$\frac{1}{1 - z^{-1}}$	$ z > 1$
$\alpha^n u[n]$	$\frac{1}{1 - \alpha z^{-1}}$	$ z > \alpha $
$n\alpha^n u[n]$	$\frac{\alpha z^{-1}}{(1 - \alpha z^{-1})^2}$	$ z > \alpha $
$[\cos(\Omega_1 n)]u[n]$	$\frac{1 - z^{-1} \cos \Omega_1}{1 - z^{-1} 2 \cos \Omega_1 + z^{-2}}$	$ z > 1$
$[\sin(\Omega_1 n)]u[n]$	$\frac{z^{-1} \sin \Omega_1}{1 - z^{-1} 2 \cos \Omega_1 + z^{-2}}$	$ z > 1$
$[r^n \cos(\Omega_1 n)]u[n]$	$\frac{1 - z^{-1} r \cos \Omega_1}{1 - z^{-1} 2r \cos \Omega_1 + r^2 z^{-2}}$	$ z > r$
$[r^n \sin(\Omega_1 n)]u[n]$	$\frac{z^{-1} r \sin \Omega_1}{1 - z^{-1} 2r \cos \Omega_1 + r^2 z^{-2}}$	$ z > r$

Signal	Bilateral Transform	ROC
$u[-n - 1]$	$\frac{1}{1 - z^{-1}}$	$ z < 1$
$-\alpha^n u[-n - 1]$	$\frac{1}{1 - \alpha z^{-1}}$	$ z < \alpha $
$-n\alpha^n u[-n - 1]$	$\frac{\alpha z^{-1}}{(1 - \alpha z^{-1})^2}$	$ z < \alpha $

E.2 z-Transform Properties

Signal	Unilateral Transform $x[n] \xleftrightarrow{z_u} X(z)$ $y[n] \xleftrightarrow{z_u} Y(z)$	Bilateral Transform $x[n] \xleftrightarrow{z} X(z)$ $y[n] \xleftrightarrow{z} Y(z)$	ROC $z \in R_x$ $z \in R_y$
$ax[n] + by[n]$	$aX(z) + bY(z)$	$aX(z) + bY(z)$	At least $R_x \cap R_y$
$x[n - k]$	See below	$z^{-k}X(z)$	R_x , except possibly $ z = 0, \infty$
$\alpha^n x[n]$	$X\left(\frac{z}{\alpha}\right)$	$X\left(\frac{z}{\alpha}\right)$	$ \alpha R_x$
$x[-n]$	—	$X\left(\frac{1}{z}\right)$	$\frac{1}{R_x}$
$x[n] * y[n]$	$X(z)Y(z)$ if $x[n] = y[n] = 0$ for $n < 0$	$X(z)Y(z)$	At least $R_x \cap R_y$
$nx[n]$	$-z \frac{d}{dz} X(z)$	$-z \frac{d}{dz} X(z)$	R_x , except possibly addition or deletion of $z = 0$

E.2.1 UNILATERAL Z-TRANSFORM TIME-SHIFT PROPERTY

$$x[n - k] \xleftrightarrow{z_u} x[-k] + x[-k + 1]z^{-1} + \dots + x[-1]z^{-k+1} + z^{-k}X(z) \quad \text{for } k > 0$$

$$x[n + k] \xleftrightarrow{z_u} -x[0]z^k - x[1]z^{k-1} - \dots - x[k - 1]z + z^kX(z) \quad \text{for } k > 0$$

Euler's Relation

$$e^{j(\omega t + \phi)} + e^{-j(\omega t + \phi)} = 2 \cos(\omega t + \phi)$$

$$e^{j(\omega t + \phi)} - e^{-j(\omega t + \phi)} = 2j \sin(\omega t + \phi)$$

Trigonometric Formulas

$$\begin{aligned} 2 \sin A \sin B &= \cos(A - B) - \cos(A + B) & 2 \cos A \cos B &= \cos(A + B) + \cos(A - B) \\ 2 \cos A \sin B &= \sin(A + B) - \sin(A - B) & 2 \sin A \cos B &= \sin(A + B) + \sin(A - B) \\ \sin(A \pm B) &= \sin A \cos B \pm \cos A \sin B & \cos(A \pm B) &= \cos A \cos B \mp \sin A \sin B \end{aligned}$$

Sifting Property

$$x(t) = \int_{-\infty}^{\infty} x(\tau) \delta(t - \tau) d\tau = x(t) * \delta(t) \quad x[n] = \sum_{k=-\infty}^{\infty} x[k] \delta[n - k] = x[n] * \delta[n]$$

Definite Integrals

$$\int_a^b x^n dx = \frac{1}{n+1} x^{n+1} \Big|_a^b, \quad n \neq -1$$

$$\int_a^b e^{cx} dx = \frac{1}{c} e^{cx} \Big|_a^b$$

$$\int_a^b x e^{cx} dx = \frac{1}{c^2} e^{cx} (cx - 1) \Big|_a^b$$

$$\int_a^b \cos(cx) dx = \frac{1}{c} \sin(cx) \Big|_a^b$$

$$\int_a^b \sin(cx) dx = -\frac{1}{c} \cos(cx) \Big|_a^b$$

$$\int_a^b x \cos(cx) dx = \frac{1}{c^2} (\cos(cx) + cx \sin(cx)) \Big|_a^b$$

$$\int_a^b x \sin(cx) dx = \frac{1}{c^2} (\sin(cx) - cx \cos(cx)) \Big|_a^b$$

$$\int_a^b e^{gx} \cos(cx) dx = \frac{e^{gx}}{g^2 + c^2} (g \cos(cx) + c \sin(cx)) \Big|_a^b$$

$$\int_a^b e^{gx} \sin(cx) dx = \frac{e^{gx}}{g^2 + c^2} (g \sin(cx) - c \cos(cx)) \Big|_a^b$$

Geometric Series

$$\sum_{n=0}^{M-1} \beta^n = \begin{cases} \frac{1 - \beta^M}{1 - \beta}, & \beta \neq 1 \\ M, & \beta = 1 \end{cases}$$

$$\sum_{n=k}^l \beta^n = \begin{cases} \frac{\beta^k - \beta^{l+1}}{1 - \beta}, & \beta \neq 1 \\ l - k + 1, & \beta = 1 \end{cases}$$

$$\sum_{n=0}^{\infty} \beta^n = \frac{1}{1 - \beta}, \quad |\beta| < 1$$

$$\sum_{n=k}^{\infty} \beta^n = \frac{\beta^k}{1 - \beta}, \quad |\beta| < 1$$

$$\sum_{n=-k}^{-\infty} \beta^n = \beta^{-k} \left(\frac{\beta}{\beta - 1} \right), \quad |\beta| > 1$$

$$\sum_{n=0}^{\infty} n \beta^n = \frac{\beta}{(1 - \beta)^2}, \quad |\beta| < 1$$

Quadratic formula

The solution to:

$$ax^2 + bx + c = 0$$

is:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

In the following table you can use any convenient period rather than the default

\int_0^T for the Fourier Series and $\sum_{n=0}^{N-1}$ for the Discrete-Time Fourier Series

TABLE 3.2 The Four Fourier Representations.

Time Domain	Periodic (t, n)	Non periodic (t, n)	
Continuous (t)	<p>Fourier Series</p> $x(t) = \sum_{k=-\infty}^{\infty} X[k]e^{jk\omega_o t}$ $X[k] = \frac{1}{T} \int_0^T x(t)e^{-jk\omega_o t} dt$ <p>$x(t)$ has period T</p> $\omega_o = \frac{2\pi}{T}$	<p>Fourier Transform</p> $x(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} X(j\omega)e^{j\omega t} d\omega$ $X(j\omega) = \int_{-\infty}^{\infty} x(t)e^{-j\omega t} dt$	Non periodic (k, ω)
Discrete (n)	<p>Discrete-Time Fourier Series</p> $x[n] = \sum_{k=0}^{N-1} X[k]e^{jk\Omega_o n}$ $X[k] = \frac{1}{N} \sum_{n=0}^{N-1} x[n]e^{-jk\Omega_o n}$ <p>$x[n]$ and $X[k]$ have period N</p> $\Omega_o = \frac{2\pi}{N}$	<p>Discrete-Time Fourier Transform</p> $x[n] = \frac{1}{2\pi} \int_{-\pi}^{\pi} X(e^{j\Omega})e^{j\Omega n} d\Omega$ $X(e^{j\Omega}) = \sum_{n=-\infty}^{\infty} x[n]e^{-j\Omega n}$ <p>$X(e^{j\Omega})$ has period 2π</p>	Periodic (k, Ω)
	Discrete (k)	Continuous (ω, Ω)	Frequency Domain

$$g(t) \xleftrightarrow{\text{FS}; \omega_o = 2\pi/T} G[k]$$

$$v[n] \xleftrightarrow{\text{DTFT}} V(e^{j\Omega})$$

$$w[n] \xleftrightarrow{\text{DTFS}; \Omega_o = 2\pi/N} W[k]$$

■ **C.8.1 FT REPRESENTATION FOR A CONTINUOUS-TIME PERIODIC SIGNAL**

$$g(t) \xleftrightarrow{\text{FT}} G(j\omega) = 2\pi \sum_{k=-\infty}^{\infty} G[k]\delta(\omega - k\omega_o)$$

■ **C.8.2 DTFT REPRESENTATION FOR A DISCRETE-TIME PERIODIC-SIGNAL**

$$w[n] \xleftrightarrow{\text{DTFT}} W(e^{j\Omega}) = 2\pi \sum_{k=-\infty}^{\infty} W[k]\delta(\Omega - k\Omega_o)$$

■ **C.8.3 FT REPRESENTATION FOR A DISCRETE-TIME NONPERIODIC SIGNAL**

$$v_\delta(t) = \sum_{n=-\infty}^{\infty} v[n]\delta(t - nT_s) \xleftrightarrow{\text{FT}} V_\delta(j\omega) = V(e^{j\Omega}) \Big|_{\Omega=\omega T_s}$$

■ **C.8.4 FT REPRESENTATION FOR A DISCRETE-TIME NONPERIODIC SIGNAL**

$$w_\delta(t) = \sum_{n=-\infty}^{\infty} w[n]\delta(t - nT_s) \xleftrightarrow{\text{FT}} W_\delta(j\omega) = \frac{2\pi}{T_s} \sum_{k=-\infty}^{\infty} W[k]\delta\left(\omega - \frac{k\Omega_o}{T_s}\right)$$

Fourier Transform Pairs

<i>Time Domain</i>	<i>Frequency Domain</i>
$x(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} X(j\omega) e^{j\omega t} d\omega$	$X(j\omega) = \int_{-\infty}^{\infty} x(t) e^{-j\omega t} dt$
$x(t) = \begin{cases} 1, & t \leq T_o \\ 0, & \text{otherwise} \end{cases}$	$X(j\omega) = \frac{2 \sin(\omega T_o)}{\omega}$
$x(t) = \frac{1}{\pi t} \sin(Wt)$	$X(j\omega) = \begin{cases} 1, & \omega \leq W \\ 0, & \text{otherwise} \end{cases}$
$x(t) = \delta(t)$	$X(j\omega) = 1$
$x(t) = 1$	$X(j\omega) = 2\pi\delta(\omega)$
$x(t) = u(t)$	$X(j\omega) = \frac{1}{j\omega} + \pi\delta(\omega)$
$x(t) = e^{-at}u(t), \quad \text{Re}\{a\} > 0$	$X(j\omega) = \frac{1}{a + j\omega}$
$x(t) = te^{-at}u(t), \quad \text{Re}\{a\} > 0$	$X(j\omega) = \frac{1}{(a + j\omega)^2}$
$x(t) = e^{-a t }, \quad a > 0$	$X(j\omega) = \frac{2a}{a^2 + \omega^2}$
$x(t) = \frac{1}{\sqrt{2\pi}} e^{-t^2/2}$	$X(j\omega) = e^{-\omega^2/2}$

<i>Periodic Time-Domain Signal</i>	<i>Fourier Transform</i>
$x(t) = \sum_{k=-\infty}^{\infty} X[k] e^{jk\omega_o t}$	$X(j\omega) = 2\pi \sum_{k=-\infty}^{\infty} X[k] \delta(\omega - k\omega_o)$
$x(t) = \cos(\omega_o t)$	$X(j\omega) = \pi\delta(\omega - \omega_o) + \pi\delta(\omega + \omega_o)$
$x(t) = \sin(\omega_o t)$	$X(j\omega) = \frac{\pi}{j}\delta(\omega - \omega_o) - \frac{\pi}{j}\delta(\omega + \omega_o)$
$x(t) = e^{j\omega_o t}$	$X(j\omega) = 2\pi\delta(\omega - \omega_o)$
$x(t) = \sum_{n=-\infty}^{\infty} \delta(t - nT_s)$	$X(j\omega) = \frac{2\pi}{T_s} \sum_{k=-\infty}^{\infty} \delta\left(\omega - k\frac{2\pi}{T_s}\right)$
$x(t) = \begin{cases} 1, & t \leq T_o \\ 0, & T_o < t < T/2 \end{cases}$ $x(t + T) = x(t)$	$X(j\omega) = \sum_{k=-\infty}^{\infty} \frac{2 \sin(k\omega_o T_o)}{k} \delta(\omega - k\omega_o)$


Fourier Transform Properties

	<p style="text-align: center;"><i>Fourier Transform</i></p> $x(t) \xleftrightarrow{FT} X(j\omega)$ $y(t) \xleftrightarrow{FT} Y(j\omega)$
Property	
Linearity	$ax(t) + by(t) \xleftrightarrow{FT} aX(j\omega) + bY(j\omega)$
Time shift	$x(t - t_0) \xleftrightarrow{FT} e^{-j\omega t_0} X(j\omega)$
Frequency shift	$e^{j\gamma t} x(t) \xleftrightarrow{FT} X(j(\omega - \gamma))$
Scaling	$x(at) \xleftrightarrow{FT} \frac{1}{ a } X\left(\frac{j\omega}{a}\right)$
Differentiation in time	$\frac{d}{dt} x(t) \xleftrightarrow{FT} j\omega X(j\omega)$
Differentiation in frequency	$-jtx(t) \xleftrightarrow{FT} \frac{d}{d\omega} X(j\omega)$
Integration/Summation	$\int_{-\infty}^t x(\tau) d\tau \xleftrightarrow{FT} \frac{X(j\omega)}{j\omega} + \pi X(j0)\delta(\omega)$
Convolution	$\int_{-\infty}^{\infty} x(\tau)y(t - \tau) d\tau \xleftrightarrow{FT} X(j\omega)Y(j\omega)$
Multiplication	$x(t)y(t) \xleftrightarrow{FT} \frac{1}{2\pi} \int_{-\infty}^{\infty} X(j\nu)Y(j(\omega - \nu)) d\nu$
Parseval's Theorem	$\int_{-\infty}^{\infty} x(t) ^2 dt = \frac{1}{2\pi} \int_{-\infty}^{\infty} X(j\omega) ^2 d\omega$
Duality	$X(jt) \xleftrightarrow{FT} 2\pi x(-\omega)$
Symmetry	$x(t) \text{ real} \xleftrightarrow{FT} X^*(j\omega) = X(-j\omega)$ $x(t) \text{ imaginary} \xleftrightarrow{FT} X^*(j\omega) = -X(-j\omega)$ $x(t) \text{ real and even} \xleftrightarrow{FT} \text{Im}\{X(j\omega)\} = 0$ $x(t) \text{ real and odd} \xleftrightarrow{FT} \text{Re}\{X(j\omega)\} = 0$

Effect of Sampling

$$\sum_n x(nT)\delta(t - nT) \xleftrightarrow{FT} \frac{1}{T} \sum_{k=-\infty}^{\infty} X\left(\omega - k\frac{2\pi}{T}\right)$$

Sampling in time and frequency (digital signal processing)



$$N_0 = \frac{T_0}{T} = \frac{f_s}{f_0}, \quad \text{where } N_0 \text{ is the number of samples over } T_0 \text{ or } f_s$$

Discrete-Time Fourier Transform Pairs

Time Domain	Frequency Domain
$x[n] = \frac{1}{2\pi} \int_{-\pi}^{\pi} X(e^{j\Omega}) e^{j\Omega n} d\Omega$	$X(e^{j\Omega}) = \sum_{n=-\infty}^{\infty} x[n] e^{-j\Omega n}$
$x[n] = \begin{cases} 1, & n \leq M \\ 0, & \text{otherwise} \end{cases}$	$X(e^{j\Omega}) = \frac{\sin\left[\Omega\left(\frac{2M+1}{2}\right)\right]}{\sin\left(\frac{\Omega}{2}\right)}$
$x[n] = \alpha^n u[n], \quad \alpha < 1$	$X(e^{j\Omega}) = \frac{1}{1 - \alpha e^{-j\Omega}}$
$x[n] = \delta[n]$	$X(e^{j\Omega}) = 1$
$x[n] = u[n]$	$X(e^{j\Omega}) = \frac{1}{1 - e^{-j\Omega}} + \pi \sum_{p=-\infty}^{\infty} \delta(\Omega - 2\pi p)$
$x[n] = \frac{1}{\pi n} \sin(Wn), \quad 0 < W \leq \pi$	$X(e^{j\Omega}) = \begin{cases} 1, & \Omega \leq W \\ 0, & W < \Omega \leq \pi \end{cases} \quad X(e^{j\Omega}) \text{ is } 2\pi \text{ periodic}$
$x[n] = (n+1)\alpha^n u[n]$	$X(e^{j\Omega}) = \frac{1}{(1 - \alpha e^{-j\Omega})^2}$

Periodic Time-Domain Signal	Discrete-Time Fourier Transform
$x[n] = \sum_{k=0}^{N-1} X[k] e^{jk\Omega_0 n}$	$X(e^{j\Omega}) = 2\pi \sum_{k=-\infty}^{\infty} X[k] \delta(\Omega - k\Omega_0)$
$x[n] = \cos(\Omega_1 n)$	$X(e^{j\Omega}) = \pi \sum_{k=-\infty}^{\infty} \delta(\Omega - \Omega_1 - k2\pi) + \delta(\Omega + \Omega_1 - k2\pi)$
$x[n] = \sin(\Omega_1 n)$	$X(e^{j\Omega}) = \frac{\pi}{j} \sum_{k=-\infty}^{\infty} \delta(\Omega - \Omega_1 - k2\pi) - \delta(\Omega + \Omega_1 - k2\pi)$
$x[n] = e^{j\Omega_1 n}$	$X(e^{j\Omega}) = 2\pi \sum_{k=-\infty}^{\infty} \delta(\Omega - \Omega_1 - k2\pi)$
$x[n] = \sum_{k=-\infty}^{\infty} \delta(n - kN)$	$X(e^{j\Omega}) = \frac{2\pi}{N} \sum_{k=-\infty}^{\infty} \delta\left(\Omega - \frac{k2\pi}{N}\right)$

Discrete-Time Fourier Transform Properties

Property	<p>Discrete-Time FT</p> $x[n] \xleftrightarrow{DTFT} X(e^{j\Omega})$ $y[n] \xleftrightarrow{DTFT} Y(e^{j\Omega})$
Linearity	$ax[n] + by[n] \xleftrightarrow{DTFT} aX(e^{j\Omega}) + bY(e^{j\Omega})$
Time shift	$x[n - n_o] \xleftrightarrow{DTFT} e^{-j\Omega n_o} X(e^{j\Omega})$
Frequency shift	$e^{j\Gamma n} x[n] \xleftrightarrow{DTFT} X(e^{j(\Omega - \Gamma)})$
Scaling	$x_z[n] = 0, \quad n \neq 0, \pm p, \pm 2p, \pm 3p, \dots$ $x_z[pn] \xleftrightarrow{DTFT} X_z(e^{j\Omega/p})$
Differentiation in time	—
Differentiation in frequency	$-jnx[n] \xleftrightarrow{DTFT} \frac{d}{d\Omega} X(e^{j\Omega})$
Integration/Summation	$\sum_{k=-\infty}^n x[k] \xleftrightarrow{DTFT} \frac{X(e^{j\Omega})}{1 - e^{-j\Omega}}$ $+ \pi X(e^{j0}) \sum_{k=-\infty}^{\infty} \delta(\Omega - k2\pi)$
Convolution	$\sum_{l=-\infty}^{\infty} x[l]y[n - l] \xleftrightarrow{DTFT} X(e^{j\Omega})Y(e^{j\Omega})$
Multiplication	$x[n]y[n] \xleftrightarrow{DTFT} \frac{1}{2\pi} \int_{-\pi}^{\pi} X(e^{j\Gamma})Y(e^{j(\Omega - \Gamma)}) d\Gamma$
Parseval's Theorem	$\sum_{n=-\infty}^{\infty} x[n] ^2 = \frac{1}{2\pi} \int_{-\pi}^{\pi} X(e^{j\Omega}) ^2 d\Omega$
Duality	$x[n] \xleftrightarrow{DTFT} X(e^{j\Omega})$ $X(e^{j\Omega}) \xleftrightarrow{FS;1} x[-k]$
Symmetry	$x[n] \text{ real} \xleftrightarrow{DTFT} X^*(e^{j\Omega}) = X(e^{-j\Omega})$ $x[n] \text{ imaginary} \xleftrightarrow{DTFT} X^*(e^{j\Omega}) = -X(e^{-j\Omega})$ $x[n] \text{ real and even} \xleftrightarrow{DTFT} \text{Im}\{X(e^{j\Omega})\} = 0$ $x[n] \text{ real and odd} \xleftrightarrow{DTFT} \text{Re}\{X(e^{j\Omega})\} = 0$