

# Spatiotemporal model-based index development for Bering Sea and Aleutian Islands crab stocks

Update for Crab Plan Team modeling workshop

Caitlin Stern<sup>1,2</sup>, Emily Ryznar<sup>3,4</sup>, and Jon Richar<sup>3,5</sup>

<sup>1</sup> Alaska Department of Fish and Game

<sup>2</sup>caitlin.stern@alaska.gov

<sup>3</sup>NOAA Fisheries

<sup>4</sup>emily.ryznar@noaa.gov

<sup>5</sup>jon.richar@noaa.gov

December 2024

## Introduction

The goal of this investigation was to develop spatiotemporal model-based indices of abundance for three Bering Sea and Aleutian Islands (BSAI) crab stocks: Tanner crab (*Chionoecetes bairdi*), Norton Sound red king crab (*Paralithodes camtschaticus*), and St. Matthew Island blue king crab (*Paralithodes platypus*). Research suggests that spatiotemporal model-based indices can be more robust to survey changes than are design-based indices, though the models must be well-specified (Yalcin et al. 2023). Spatiotemporal model-based indices are used in North Pacific Fishery Management Council (NPFMC) groundfish stock assessments for species including Eastern Bering Sea (EBS) walleye pollock (*Gadus chalcogrammus*) and EBS Pacific cod (*Gadus macrocephalus*), both of which use the vector-autoregressive spatial temporal (VAST) approach (Thorson 2019) to produce indices used in the assessments (Ianelli et al. 2024; Barbeaux et al. 2024). Previous BSAI crab stock assessments have presented models using spatiotemporal model-based indices (e.g., Ianelli et al. 2017), although these models were not accepted for harvest specifications (SSC 2017).

We generated biomass and abundance estimates using the R package *sdmTMB* (Anderson et al. 2022), which uses geostatistical time series data to estimate spatial and spatiotemporal generalized linear mixed effects models. This approach allows for index standardization when the set of stations surveyed is not consistent across years: one can generate a spatial grid that covers the area of interest, predict from the model onto that grid, and sum the predicted biomass to obtain an area-weighted biomass index that is independent of sampling locations (Anderson et al. 2022).

All three stock assessments for the crab stocks presented here use data from the National Marine Fisheries Service (NMFS) EBS bottom trawl survey (Stockhausen 2024; Hamazaki 2024; Stern and Palof 2024). The St. Matthew Island blue king crab stock assessment also uses data from the Alaska Department of Fish and Game (ADF&G) St. Matthew Island blue king crab pot survey, while the Norton Sound red king crab stock assessment uses data from the NMFS Northern Bering Sea bottom trawl survey and the ADF&G Norton Sound red king crab trawl survey.

Spatiotemporal model-based index development is expected to confer distinct advantages for each of the three stocks. For the St. Matthew Island blue king crab stock, standardizing the survey indices could allow the assessment to use the existing survey data more rigorously. The NMFS EBS trawl survey is undergoing changes including dropping the high sampling density “corner stations” near St. Matthew Island from 2024 onward (DePhilippo et al. 2023; Stern & Palof 2024); index standardization will allow the assessment to

continue using the full time series of data despite changes in the spatial footprint of the survey. For Norton Sound red king crab, a model-based approach could provide a more consistent way to combine the three existing trawl survey data sets into a single index of abundance.

## Methods

We fit models using the R package *sdmTMB*. For all models fitted using *sdmTMB*, we used the R package *DHARMA* (Hartig 2022) for model diagnostics. After calculating the DHARMA residuals using the function `DHARMA::dharma_residuals()`, we tested for quantile deviations, under/overdispersion, outliers, and zero inflation using the functions `DHARMA::testQuantiles()`, `DHARMA::testDispersion()`, `DHARMA::testOutliers()`, and `DHARMA::testZeroInflation()`, respectively.

### Tanner crab

#### Norton Sound red king crab

We combined data from the NMFS trawl survey (1976-1991), ADF&G trawl survey (1996-2024), and NMFS NBS trawl survey (2010-2023) into a single data set to which we fit models in *sdmTMB*. We filtered the data set to ensure that it included only observations with coordinates falling within the Norton Sound Section of Statistical Area Q. For model fitting, we used spatial meshes at three resolutions, specified in terms of the number of knots (vertices), where a higher number of knots indicates a higher resolution mesh: 100 knots, 50 knots, and 30 knots (Figures 6 - 4). We used a prediction grid with resolution of 5 km<sup>2</sup> (Figure 3).

#### St. Matthew Island blue king crab

For model fitting, we used spatial meshes at three resolutions, specified in terms of the number of knots (vertices), where a higher number of knots indicates a higher resolution mesh: 120 knots, 90 knots, and 50 knots (Figures 17 - 15). We used a prediction grid with resolution of 4 km<sup>2</sup> (Figure 18).

## Results

### Tanner crab

#### Norton Sound red king crab

##### Model diagnostics

The DHARMA residuals diagnostic plots show evidence of quantile deviations for all three NSRKC models (Figures 7 - 9). The models with 100 knots and 50 knots showed evidence of underdispersion, with observed data less dispersed than expected under the fitted models, while the model with 30 knots did not. None of the models showed evidence of outliers or zero inflation.

##### Predicted abundance

Heat maps of predicted NSRKC abundance for the three models are show in figures 10 - 12.




Figure 1: Estimated abundance (millions) for Eastern Bering Sea Tanner crab (*Chionoecetes bairdi*). Colored lines represent abundance ( $\pm 95\%$  CI) estimated by sdmTMB, with pink and blue denoting models fit with a 50- and 120-knot mesh, respectively. Black points represent abundance ( $\pm 95\%$  CI) estimated by the NMFS summer bottom trawl survey.

## EBS Tanner estimated biomass




Figure 2: Estimated biomass (tons) for Eastern Bering Sea Tanner crab (*Chionoecetes bairdi*). Colored lines represent abundance ( $\pm 95\%$  CI) estimated by sdmTMB, with pink and blue denoting models fit with a 50- and 120-knot mesh, respectively. Black points represent abundance ( $\pm 95\%$  CI) estimated by the NMFS summer bottom trawl survey.

## **St. Matthew Island blue king crab**

### **Model diagnostics**

Examination of DHARMA residuals showed similar patterns for the three SMBKC models (Figures 19 - 21). All three models showed evidence of underdispersion, with observed data less dispersed than expected under the fitted models. None of the models showed evidence of outliers. All three models showed evidence of quantile deviations. The model with 120 knots showed evidence of zero inflation, with the observed data containing more zeros than would be expected under the fitted model, but the models with 50 and 90 knots did not show evidence of zero inflation.

### **Predicted abundance**

Heat maps of predicted SMBKC abundance for the three models are show in figures 22 - 24.

### **Predicted index fits to observations**

The model-predicted indices varied in their fits to the survey biomass observations, with the model fit using a mesh with an intermediate number of knots seeming to fit the survey observations more closely than the models fit to meshes with either higher or lower numbers of knots (Figure 25).

## **Conclusions**

## **Acknowledgements**

The authors thank Katie Palof and Mike Litzow for their support and feedback on this work.

## References

- Anderson, S.C.; Ward, E.J.; English, P.A.; Barnett, L.A.K. 2022. sdmTMB: An R package for fast, flexible, and user-friendly generalized linear mixed effects models with spatial and spatiotemporal random fields. *bioRxiv*.
- Barbeaux, S.J., L. Barnett, P. Hulson, J. Nielsen, S.K. Shotwell, E. Siddon, I. Spies. 2024. Assessment of the Pacific cod stock in the Eastern Bering Sea. In: Stock Assessment and Fishery Evaluation report for the Groundfish Resources of the Bering Sea/Aleutian Islands regions. North Pacific Fishery Management Council, Anchorage, AK.
- DeFilippo, L., S. Kotwicki, L. Barnett, J. Richar, M.A. Litzow, W.T. Stockhausen, K. Palof. 2023. Evaluating the impacts of reduced sampling density in a systematic fisheries-independent survey design. *Frontiers in Marine Science* 10:1219283.
- Hamazaki, T. 2024. Norton Sound red king crab stock assessment for the fishing year 2024. In: Stock Assessment and Fishery Evaluation Report for the King and Tanner Crab Fisheries of the Bering Sea and Aleutian Islands: 2024 Final Crab SAFE. North Pacific Fishery Management Council, Anchorage AK.
- Hartig, F. 2022. DHARMA: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression Models. R package version 0.4.6, <https://CRAN.R-project.org/package=DHARMA>.
- Ianelli, J., D. Webber, J. Zheng, A. Letaw. 2017. Saint Matthew Island blue king crab stock assessment 2017. In: Stock Assessment and Fishery Evaluation Report for the King and Tanner Crab Fisheries of the Bering Sea and Aleutian Islands: 2017 Final Crab SAFE. North Pacific Fishery Management Council, Anchorage AK.
- Ianelli, J., T. Honkalehto, S. Wassermann, A. McCarthy, S. Steinessen, C. McGilliard, E. Siddon. 2024. Assessment of walleye pollock in the eastern Bering Sea. In: Stock Assessment and Fishery Evaluation report for the Groundfish Resources of the Bering Sea/Aleutian Islands regions. North Pacific Fishery Management Council, Anchorage, AK.
- SSC. 2017. Scientific and Statistical Committee report to the North Pacific Fishery Management Council, October 2nd-4th, 2017. North Pacific Fishery Management Council, Anchorage, AK.
- Stern, C., K. Palof. 2024. Saint Matthew Island blue king crab stock assessment 2024. In: Stock Assessment and Fishery Evaluation Report for the King and Tanner Crab Fisheries of the Bering Sea and Aleutian Islands: 2024 Final Crab SAFE. North Pacific Fishery Management Council, Anchorage AK.
- Stockhausen, W.T. 2024. 2024 stock assessment and fishery evaluation report for the Tanner crab fisheries of the Bering Sea and Aleutians Islands regions. In: Stock Assessment and Fishery Evaluation Report for the King and Tanner Crab Fisheries of the Bering Sea and Aleutian Islands: 2024 Final Crab SAFE. North Pacific Fishery Management Council, Anchorage AK.
- Thorson, J.T. 2019. Guidance for decisions using the vector autoregressive spatio-temporal (VAST) package in stock, ecosystem, habitat, and climate assessments. *Fisheries Research* 210:143-161. <https://doi.org/10.1016/j.fishres.2018.10.013>
- Yalcin, S., S.C. Anderson, P.M. Regular, P.A. English. 2023. Exploring the limits of spatiotemporal and design-based index standardization under reduced survey coverage. *ICES Journal of Marine Sciences* 80:2368-2379. <https://doi.org/10.1093/icesjms/fsad155>

## Tables

## Figures




Figure 3: Prediction grid used for Norton Sound red king crab spatial abundance predictions. Spatial resolution is  $5 \text{ km}^2$ .




Figure 4: Spatial mesh with 30 knots used for fitting Norton Sound red king crab spatial models. Points represent observations and vertices represent knot locations.




Figure 5: Spatial mesh with 50 knots used for fitting Norton Sound red king crab spatial models. Points represent observations and vertices represent knot locations.




Figure 6: Spatial mesh with 100 knots used for fitting Norton Sound red king crab spatial models. Points represent observations and vertices represent knot locations.




Figure 7: Model diagnostic plots using DHARMA residuals for the Norton Sound red king crab model with 30 knots.




Figure 8: Model diagnostic plots using DHARMA residuals for the Norton Sound red king crab model with 50 knots.




Figure 9: Model diagnostic plots using DHARMA residuals for Norton Sound red king crab model with 100 knots.

## NSRKC predicted abundance, 30 knots



Figure 10: Heat map of Norton Sound red king crab predicted abundance generated using the model with 50 knots.

## NSRKC predicted abundance, 50 knots



Figure 11: Heat map of Norton Sound red blue king crab predicted abundance generated using the model with 90 knots.

## NSRKC predicted abundance, 100 knots




Figure 12: Heat map of Norton Sound red blue king crab predicted abundance generated using the model with 120 knots.




Figure 13: Estimated abundance in number of crab for Norton Sound red king crab. Colored points represent abundance ( $\pm 95\%$  CI) estimated by the trawl surveys.




Figure 14: Estimated abundance in number of crab for Norton Sound red king crab. Colored lines represent abundance ( $\pm 95\%$  CI) estimated using sdmTMB. Black points represent abundance ( $\pm 95\%$  CI) estimated by the trawl surveys.




Figure 15: Spatial mesh with 50 knots used for fitting St. Matthew Island blue king crab spatial models. Points represent observations and vertices represent knot locations.




Figure 16: Spatial mesh with 90 knots used for fitting St. Matthew Island blue king crab spatial models. Points represent observations and vertices represent knot locations.




Figure 17: Spatial mesh with 120 knots used for fitting St. Matthew Island blue king crab spatial models. Points represent observations and vertices represent knot locations.




Figure 18: Prediction grid used for St. Matthew Island blue king crab spatial abundance predictions. Spatial resolution is  $4 \text{ km}^2$ .




Figure 19: Model diagnostic plots using DHARMA residuals for the model with 50 knots.




Figure 20: Model diagnostic plots using DHARMA residuals for the model with 90 knots.




Figure 21: Model diagnostic plots using DHARMA residuals for the model with 120 knots.

### BKC predicted abundance, 50 knots




Figure 22: Heat map of St. Matthew Island blue king crab predicted abundance generated using the model with 50 knots.

### BKC predicted abundance, 90 knots




Figure 23: Heat map of St. Matthew Island blue king crab predicted abundance generated using the model with 90 knots.

### BKC predicted abundance, 120 knots




Figure 24: Heat map of St. Matthew Island blue king crab predicted abundance generated using the model with 120 knots.




Figure 25: Estimated biomass (t) for St. Matthew Island blue king crab. Colored lines represent biomass ( $\pm 95\%$  CI) estimated using sdmTMB. Black points represent biomass ( $\pm 95\%$  CI) estimated by the NMFS EBS bottom trawl survey.