

Refer to the provided table and answer the question

. Question : What percentage of patients had vomiting ?. Table : \{" table\\_title ": Table 2, " table\\_caption ": Distribution of associated symptoms among the ADL cases., " table\\_headers ": ['Symptom', 'Prevalence among episodes', 'Percentage'], " table\\_sub\\_headers ": [], " table\\_rows ": [['Pain in', '', ''], ['Upper arms', '12', '10.62'], ['Legs', '83', '73.45'], ['Genitals', '29', '20.00'], ['Breasts', '1', '0.88'], ['Tenderness in', '', ''], ['Upper arms', '10', '8.85'], ['Legs', '13', '11.50'], ['Genitals', '24', '21.24'], ['Lymphangitis in', '', ''], ['Left inguinal region', '34', '30.09'], ['Right inguinal region', '38', '33.63'], ['Left papillary', '3', '2.65'], ['Right papillary', '3', '2.65'], ['Swelling', '92', '81.42'], ['Fever', '', ''], ['High', '61', '53.98'], ['Low', '27', '23.89'], ['Anorexia', '72', '63.72'], ['Nausea', '61', '53.98'], ['Vomiting', '32', '28.32'], ['Microfilariemia e', '13', '11.50 a']], " table\\_footnote ": apercentage is to total number of patients\}.