

Refer to the provided table and answer the question

. Question : What percentage of patients had vomiting ?. Table : `\{"table_title": Table 2, "table_caption": Distribution of associated symptoms among the AD L cases., "table_headers": ['Symptom', 'Prevalence among episodes', 'Percentage'], "table_subheaders": [], "table_rows": [['Pain in', '', ''], ['Upper arms', '12', '10.62'], ['Leg s', '83', '73.45'], ['Gen itals', '29', '20.00'], ['Breasts', '1', '0.88'], ['Tenderness in', '', ''], ['Upper arms', '10', '8.85'], ['Leg s', '13', '11.50'], ['Gen itals', '24', '21.24'], ['Lymphangitis in', '', ''], ['Left inguinal region', '34', '30.09'], ['Right inguinal region', '38', '33.63'], ['Left papillary', '3', '2.65'], ['Right papillary', '3', '2.65'], ['Swelling', '92', '81.42'], ['Fever', '', ''], ['High', '61', '53.98'], ['Low', '27', '23.89'], ['Anorexia', '72', '63.72'], ['Nausea', '61', '53.98'], ['Vomiting', '32', '28.32'], ['Microfilaraemiae', '13', '11.50a']], "table_footnote": a percentage is to total number of patients\}.`