Hypocoercivity-preserving schemes for kinetic equations

Yi Cai, School of Mathematical Sciences, University of Science and Technology

In collaboration with Francis Filbet (IMT), Tao Xiong (USTC) and Alain Blaustein (UDL)

Talk at CSIAM, Lanzhou, China (Oct. 11, 2026)

Contents

1. Hypocoercivity theory for kinetic equations

2. Discontinuous Galerkin/Hermite spectral methods for VPFP system

3. Conclusions and perspectives

Contents

1. Hypocoercivity theory for kinetic equations

1.1 Overview

1.2 L^2 hypocoercivity

1.3 Discrete hypocoercivity

2. Discontinuous Galerkin/Hermite spectral methods for VPFP system


3. Conclusions and perspectives

Kinetic theory: bridging microscopic and macroscopic worlds

Microscopic




Motion




Interaction

Kinetic




Distribution


Macroscopic




Rarefied gas



Plasma



Radiation



Collective behavior

Applications



Aerospace



Controlled fusion



Neuroscience




Active matter

Kinetic equation: coupling of transport and collision

$$\partial_t f + \mathscr{T}(f) = \mathscr{C}(f)$$

Transport


Transports information along characteristics, rearranging the distribution in phase space.



Coupling

Collision

Dissipates nonequilibrium perturbations, driving the distribution toward local equilibrium.



Long-time behavior: convergence toward global equilibrium

Starting from a nonequilibrium, what happens as $t \to +\infty$?

Distribution

$$f(t, x, v) \xrightarrow[\text{transport + collision}]{t \to +\infty} f_\infty(x, v)$$

Global equilibrium

Convergence

Does the solution converge?

Equilibrium

Which equilibrium is selected?

Rate

How fast does the solution converge?

A simple example for difficulties in analysis

Consider a simple but representative model

$$\partial_t f + v\partial_x f - \partial_x \Phi \partial_v f = \rho M - f, \quad (x, v) \in \mathbb{T} \times \mathbb{R},$$

where $\Phi(x) \in W^{1,\infty}$, $M(v) = (2\pi)^{-1/2} e^{-|v|^2/2}$ and $\rho(t, x) = \int_{\mathbb{R}} f(t, x, v) \mathrm{d}v$.

Qualitative convergence

A **compactness method**[1] proves the convergence to the global equilibrium

$$f_\infty(x, v) = \rho_\infty(x) M(v),$$

where ρ_∞ is determined by the initial data and conservation law

$$\rho_\infty(x) = c_\infty e^{-\Phi(x)}, \quad c_\infty \int e^{-\Phi(x)} \mathrm{d}x = \int \rho^{in}(x) \mathrm{d}x.$$

[1] Bouchut and Dolbeault (1995)

Obstacle from qualitative to quantitative: degenerate dissipation

Measure the distance between f and $f_\infty = \rho_\infty M$ by

$$\mathcal{E}(t) = \frac{1}{2} \iint |f(t) - f_\infty|^2 f_\infty^{-1} \mathrm{d}x\mathrm{d}v.$$

Time differentiation gives

$$\frac{\mathrm{d}\mathcal{E}}{\mathrm{d}t} = -\mathcal{D} \le 0, \quad \mathcal{D}(t) = \iint |f(t) - \rho(t)M|^2 f_\infty^{-1} \mathrm{d}x\mathrm{d}v.$$

Coercive estimate

A coercive estimate turns energy dissipation into exponential decay

$$\mathcal{D}(t) \overset{?}{\ge} \lambda \mathcal{E}(t) \overset{\lambda>0}{\Longrightarrow} \mathcal{E}(t) \le e^{-\lambda t} \mathcal{E}(0).$$

Counterexample

$$\mathcal{D} = 0 \iff f = \rho M.$$

Take $f_* = \rho_* M$ with $\rho_* \neq \rho_\infty$ implies

$$\mathcal{D} = 0 \quad \text{but} \quad \mathcal{E} > 0.$$

Degenerate dissipation beyond the simple example

Let $\mathcal{E}$ measures the distance to the global equilibrium, and $\mathcal{D}$ represents dissipation induced by the collision.

$$\frac{\mathrm{d}\mathcal{E}}{\mathrm{d}t} = -\mathcal{D}, \qquad \mathcal{D} = 0 \iff \mathscr{C}(f) = 0.$$

Degenerate dissipation beyond the simple example

Let $\mathcal{E}$ measures the distance to the global equilibrium, and $\mathcal{D}$ represents dissipation induced by the collision.

$$\frac{\mathrm{d}\mathcal{E}}{\mathrm{d}t} = -\mathcal{D}, \qquad \mathcal{D} = 0 \iff \mathscr{C}(f) = 0.$$

More entropies[2]

- Relative entropy
- Boltzmann entropy
- φ-entropy

[2] Desvillettes and Villani (2005)
Evans (2021)
Villani (2025)

Degenerate dissipation beyond the simple example

Let $\mathcal{E}$ measures the distance to the global equilibrium, and $\mathcal{D}$ represents dissipation induced by the collision.

$$\frac{\mathrm{d}\mathcal{E}}{\mathrm{d}t} = -\mathcal{D}, \qquad \mathcal{D} = 0 \iff \mathscr{C}(f) = 0.$$

More entropies[2]

- Relative entropy
- Boltzmann entropy
- φ-entropy

More collisions[3]

- Fokker–Planck
- Boltzmann
- Scattering
- Landau
- BGK / ES-BGK

[2] Desvillettes and Villani (2005)
Evans (2021)
Villani (2025)

[3] Desvillettes and Villani (2001)
Villani (2003)
Dolbeault, Mouhot, and Schmeiser (2009)
Guillen and Silvestre (2025)

Hypocoercivity theory: global overview

Underlying mechanism

Collision only produces dissipation in some of variables, transport propagates the degenerate dissipation to the remaining variables.

Hypocoercivity theory: global overview

Underlying mechanism

Collision only produces dissipation in some of variables, transport propagates the degenerate dissipation to the remaining variables.

A brief summary of hypocoercivity theories

- H^1 hypocoercivity → Villani (2009)
- L^2 hypocoercivity → Dolbeault, Mouhot, and Schmeiser (2015)
- φ-hypocoercivity → Dolbeault and Li (2018)
- H^{-1} hypocoercivity → Albritton et al. (2024); Brigati, Lörler, and Wang (2025)

Contents

1. Hypocoercivity theory for kinetic equations
1.1 Overview
1.2 L^2 hypocoercivity
1.3 Discrete hypocoercivity


2. Discontinuous Galerkin/Hermite spectral methods for VPFP system

3. Conclusions and perspectives

A toy example for L^2 hypocoercivity

Damping oscillator

$$\begin{cases} \dfrac{dx}{dt} = +v, \\ \dfrac{dv}{dt} = -x - \beta v. \end{cases}$$



Energy $H = \frac{1}{2}(|x|^2 + |v|^2)$ satisfies

$$\frac{dH}{dt} = -\beta |v|^2 \leq 0.$$

Observation: lack of dissipation in x!

Key idea: add a **correction term** to recover missing dissipation

$$G = H + \alpha xv.$$

Hypocoercive estimate

$$\frac{dG}{dt} = -\beta |v|^2 + \alpha |v|^2 - \alpha |x|^2 - \alpha\beta xv$$

$$\leq -\left(\beta - \alpha - \frac{\beta^2}{2}\alpha\right)|v|^2 - \frac{\alpha}{2}|x|^2.$$

Taking $\alpha = \frac{2\beta}{3+\beta^2}$ and $\lambda = \frac{\alpha}{2(1+\alpha)}$, it yields

- **Equivalence**: $(1-\alpha)H \leq G \leq (1+\alpha)H$;
- **Exponential decay**: $\frac{dG}{dt} \leq -\lambda G$.

Let $C = \frac{1+\alpha}{1-\alpha}$, it follows that

$$H(t) \leq Ce^{-\lambda t}H(0).$$

L^2 hypocoercivity: linear BGK equation

Model description

$$\partial_t f + v\partial_x f = \rho M - f.$$

Velocity integration after multiplying 1 and v,

$$\begin{cases} \partial_t \rho + \partial_x j = 0, \\ \partial_t j + D\partial_x \rho + R = -j, \end{cases}$$

where $j = \int v f \mathrm{d}v$, $D = \int |v|^2 M \mathrm{d}v$ and

$$R = \int v^2 \partial_x (f - \rho M) \mathrm{d}v.$$

Energy decay

$$\frac{1}{2}\frac{\mathrm{d}}{\mathrm{d}t}\|f(t) - f_\infty\|^2_{L^2_{x,v}(f_\infty^{-1})} = -\|f - \rho M\|^2_{L^2_{x,v}(f_\infty^{-1})}.$$

Hypocoercive energy[4]

$$\mathcal{H} = \frac{1}{2}\|f - f_\infty\|^2_{L^2_{x,v}(f_\infty^{-1})} + \alpha\langle \partial_x \cdot j, \psi \rangle_{L^2_x(\rho_\infty^{-1})},$$

where

$$D\Delta_x \psi = \rho - \rho_\infty.$$

[4] Bessemoulin-Chatard, Herda, and Rey (2020)

L^2 hypocoercivity: linear BGK equation

Regularity estimates[5]

$$\|j\|_{L^2_x(\rho_\infty^{-1})} + \|R\|_{L^2_x(\rho_\infty^{-1})} \lesssim \|f - \rho M\|_{L^2_{x,v}(f_\infty^{-1})},$$

$$\|\psi\|_{L^2_x(\rho_\infty^{-1})} + \|\partial_x \psi\|_{L^2_x(\rho_\infty^{-1})} \lesssim \|\rho - \rho_\infty\|_{L^2_x(\rho_\infty^{-1})}.$$

Computation

$$\frac{\mathrm{d}\mathcal{H}}{\mathrm{d}t} = -\|f - \rho M\|^2_{L^2_{x,v}(f_\infty^{-1})} + \alpha(\mathcal{J}_1 + \mathcal{J}_2 + \mathcal{J}_3)$$

$$\mathcal{J}_1 = -D\langle \Delta_x \rho, \psi \rangle_{L^2_x(\rho_\infty^{-1})} = -\|\rho - \rho_\infty\|^2_{L^2_x(\rho_\infty^{-1})}$$

$$\mathcal{J}_2 = -\langle \nabla_x \cdot j, \psi \rangle_{L^2_x(\rho_\infty^{-1})} \lesssim +\|f - \rho M\|_{L^2_{x,v}(f_\infty^{-1})} \|\rho - \rho_\infty\|_{L^{-1}_x(\rho_\infty^{-1})}$$

$$\mathcal{J}_3 = -\langle \nabla_x \cdot R, \psi \rangle_{L^2_x(\rho_\infty^{-1})} \lesssim +\|f - \rho M\|_{L^2_{x,v}(f_\infty^{-1})} \|\rho - \rho_\infty\|_{L^{-1}_x(\rho_\infty^{-1})}$$

Hypocoercive estimate

There exists $C > 0$ such that for all $\alpha \in [0, 1/2C]$

$$\frac{1}{2}\mathcal{E} \le \mathcal{H} \le \frac{3}{2}\mathcal{E},$$

and

$$\frac{\mathrm{d}\mathcal{H}}{\mathrm{d}t} \le -\frac{\alpha}{2}\|\rho - \rho_\infty\|^2_{L^2_x(\rho_\infty^{-1})} - (1 - 2\alpha C)\|f - \rho M\|^2_{L^2_{x,v}(f_\infty^{-1})}.$$

Take $\alpha = 2/(4C + 1)$, then

$$\frac{\mathrm{d}\mathcal{H}}{\mathrm{d}t} \le -\frac{2}{3}\alpha\mathcal{H}.$$

[5] Dolbeault, Mouhot, and Schmeiser (2015); Bessemoulin-Chatard, Herda, and Rey (2020)

L^2 hypocoercivity: diffusive limit


Model description

Kinetic model:

$$\varepsilon \partial_t f + v \cdot \nabla_x f = \frac{1}{\varepsilon}(\rho M - f).$$

Diffusion model:

$$\partial_t \rho^0 = D \Delta_x \rho^0.$$



Hypocoercive estimate

There are constants $C_1, C_2 \geq 1$ and $\lambda_1, \lambda_2 > 0$ such that for all $\varepsilon \in (0,1)$, the solution satisfies[6]:

$$\|f(t) - \rho M\|_{L^2_{x,v}(f_\infty^{-1})} \leq e^{-\lambda_1 t/(4\varepsilon^2)} \|f(0) - \rho M\|_{L^2_{x,v}(f_\infty^{-1})} + \varepsilon C_1 e^{-\lambda_1 t} \|\rho(0) - \rho_\infty\|_{H^1_x(\rho_\infty^{-1})},$$

and

$$\|\rho(t) - \rho^0(t)\|_{H^{-1}_x(\rho_\infty^{-1})} \leq C_2 e^{-\lambda_2 t} \left(\|\rho(0) - \rho^0(0)\|_{H^{-1}_x(\rho_\infty^{-1})} + \varepsilon \|\rho(0) - \rho_\infty\|_{H^{-1}_x(\rho_\infty^{-1})} \right)$$

[6] Blaustein and Filbet (2024)

Contents

1. Hypocoercivity theory for kinetic equations
1.1 Overview
1.2 L^2 hypocoercivity
1.3 Discrete hypocoercivity

2. Discontinuous Galerkin/Hermite spectral methods for VPFP system

3. Conclusions and perspectives

From energy stability to discrete hypocoercivity

Energy stability:

$$\mathcal{E}_{h,N}(t) \le \mathcal{E}_{h,N}(0), \quad \forall t \ge 0.$$

Discrete hypocoercivity:

$$\mathcal{E}_{h,N}(t) \le Ce^{-\lambda t}\mathcal{E}_{h,N}(0), \quad \forall t \ge 0.$$

The constants $C > 0$ and $\lambda > 0$ are uniform in h and N.

Counterexample: energy stable does not imply discrete hypocoercivity

$$\frac{\mathrm{d}f_j}{\mathrm{d}t} + v\,\frac{f_{j+1} - f_{j-1}}{2h} = \rho_j M - f_j, \quad f_j(0) = (1 + (-1)^j/2)\rho_\infty M, \quad 1 \le j \le 2n.$$

Representative numerical studies

A brief summary of existing work

- Kolmogorov equation → Porretta and Zuazua (2017); Georgoulis (2021); Dong, Georgoulis, and Herbert (2025)
- Free transport + Linear BGK/Fokker-Planck collision → Bessemoulin-Chatard, Herda, and Rey (2020)
- Free transport + Nonlinear reaction collision → Bessemoulin-Chatard, Laidin, and Rey (2025)
- Vlasov transport + Linearized BGK collision → Grosse (2025)
- Vlasov transport + Linear Fokker-Planck collision → Blaustein and Filbet (2024)

Contents

1. Hypocoercivity theory for kinetic equations

2. Discontinuous Galerkin/Hermite spectral methods for VPFP system
2.1 Close-to-equilibrium framework
2.2 Discretization
2.3 Numerical simulations

3. Conclusions and perspectives

Vlasov-Poisson-Fokker-Planck model

1D VPFP model

$$\partial_t f + v\partial_x f + E\partial_v f = \frac{1}{\tau_0}\partial_v\left(vf + T_0\partial_v f\right),$$

$$E = -\partial_x \Phi, \quad -\partial_{xx}\Phi = \rho - 1.$$

- electric potential $\Phi(t, x)$.
- electric field $E(t, x)$.
- background temperature $T_0 > 0$.
- mean free time $\tau_0 > 0$.

Equilibria

Maxwellian:

$$M(v) = (2\pi T_0)^{-\frac{1}{2}} e^{-\frac{|v|^2}{2T_0}}.$$

Stationary state:

$$f_\infty(v) = M(v), \quad \rho_\infty \equiv 1, \quad E_\infty \equiv 0.$$

Close-to-equilibrium framework: linearized free energy

Logarithmic free energy estimate

$$\frac{d\mathcal{G}}{dt} = -\frac{1}{\tau_0}\mathcal{D}.$$

- logarithmic free energy

$$\mathcal{G} = \int f \log\left(\frac{f}{f_\infty}\right) dx dv + \frac{1}{2T_0}\|E\|_{L^2_x}^2.$$

- dissipation from collision

$$\mathcal{D} = 4T_0 \int \left|\partial_v \left(\sqrt{\frac{f}{f_\infty}}\right)\right|^2 f_\infty \, dx dv.$$

- **Key idea**: linearization near f_∞

$$f \log\left(\frac{f}{f_\infty}\right) \approx f - f_\infty + \frac{|f - f_\infty|^2}{2f_\infty}$$

Close-to-equilibrium framework: linearized free energy

Logarithmic free energy estimate

$$\frac{\mathrm{d}\mathcal{G}}{\mathrm{d}t} = -\frac{1}{\tau_0}\mathcal{D}.$$

- logarithmic free energy

$$\mathcal{G} = \int f \log\left(\frac{f}{f_\infty}\right)\mathrm{d}x\mathrm{d}v + \frac{1}{2T_0}\|E\|_{L^2_x}^2.$$

- dissipation from collision

$$\mathcal{D} = 4T_0 \int |\partial_v\left(\sqrt{\frac{f}{f_\infty}}\right)|^2 f_\infty \mathrm{d}x\mathrm{d}v.$$

- **Key idea**: linearization near f_∞

$$f\log\left(\frac{f}{f_\infty}\right) \approx f - f_\infty + \frac{|f-f_\infty|^2}{2f_\infty}$$

Linearized free energy estimate

$$\frac{\mathrm{d}\mathcal{E}}{\mathrm{d}t} = -\frac{1}{\tau_0}\mathcal{I} + \mathcal{R}.$$

- **linearized** free energy

$$\mathcal{E} = \frac{1}{2}\int \frac{|f-f_\infty|^2}{f_\infty}\mathrm{d}x\mathrm{d}v + \frac{1}{2T_0}\|E\|_{L^2_x}^2.$$

- dissipation from **collision**

$$\mathcal{I} = T_0 \int |\partial_v\left(\frac{f}{f_\infty}\right)|^2 f_\infty \mathrm{d}x\mathrm{d}v.$$

- residual from **nonlinear field interaction**

$$\mathcal{R} = \frac{1}{2}\int E(f-f_\infty)\partial_v\left(\frac{f}{f_\infty}\right)\mathrm{d}x\mathrm{d}v.$$

Close-to-equilibrium framework: from linearized to nonlinear model

Linearized model

$$\partial_t f + v\partial_x f + E\partial_v f_\infty = \frac{1}{\tau_0}\partial_v\left(vf + T_0\partial_v f\right),$$

$$E = -\partial_x \Phi, \quad -\partial_{xx}\Phi = \rho - 1.$$

- free energy estimate

$$\frac{\mathrm{d}\mathcal{E}}{\mathrm{d}t} = -\frac{1}{\tau_0}\mathcal{I}.$$

- $\mathcal{H}$ with $\lambda \sim \min(\tau_0, \tau_0^{-1})$

$$\mathcal{H} \sim \mathcal{E} \quad \textit{and} \quad \frac{\mathrm{d}\mathcal{H}}{\mathrm{d}t} \leq -\lambda\mathcal{H}.$$

- exponential decay

$$\mathcal{E}(t) \lesssim e^{-\lambda t}\mathcal{E}(0).$$

Close-to-equilibrium framework: from linearized to nonlinear model

Linearized model

$$\partial_t f + v\partial_x f + E\partial_v f_\infty = \frac{1}{\tau_0}\partial_v\left(vf + T_0\partial_v f\right),$$

$$E = -\partial_x \Phi, \quad -\partial_{xx}\Phi = \rho - 1.$$

- free energy estimate

$$\frac{\mathrm{d}\mathcal{E}}{\mathrm{d}t} = -\frac{1}{\tau_0}\mathcal{I}.$$

- $\mathcal{H}$ with $\lambda \sim \min(\tau_0, \tau_0^{-1})$

$$\mathcal{H} \sim \mathcal{E} \quad \textit{and} \quad \frac{\mathrm{d}\mathcal{H}}{\mathrm{d}t} \leq -\lambda\mathcal{H}.$$

- exponential decay

$$\mathcal{E}(t) \lesssim e^{-\lambda t}\mathcal{E}(0).$$

Nonlinear model

$$\partial_t f + v\partial_x f + E\partial_v f_\infty + E\partial_v(f - f_\infty) = \frac{1}{\tau_0}\partial_v\left(vf + T_0\partial_v f\right),$$

$$E = -\partial_x \Phi, \quad -\partial_{xx}\Phi = \rho - 1.$$

- linearized free energy estimate

$$\frac{\mathrm{d}\mathcal{E}}{\mathrm{d}t} = -\frac{1}{\tau_0}\mathcal{I} + \mathcal{R}.$$

- $\mathcal{H}$ with $\lambda \sim \min(\tau_0, \tau_0^{-1})$ and $\mu \sim \max(\tau_0, \tau_0^{-1})$

$$\mathcal{H} \sim \mathcal{E} \quad \textit{and} \quad \frac{\mathrm{d}\mathcal{H}}{\mathrm{d}t} \leq -\lambda\mathcal{H} + \mu\mathcal{H}^2.$$

- exponential decay when $\mathcal{E}(0) \lesssim \min(\tau_0^2, \tau_0^{-2})$

$$\mathcal{E}(t) \lesssim e^{-\lambda t}\mathcal{E}(0).$$

Contents

1. Hypocoercivity theory for kinetic equations

2. Discontinuous Galerkin/Hermite spectral methods for VPFP system

2.1 Close-to-equilibrium framework

2.2 Discretization

2.3 Numerical simulations

3. Conclusions and perspectives

Velocity discretization: scaled Hermite spectral method

Scaled Hermite spectral method

Scaled Hermite expansion

$$f(t,x,v) = \sum_{k\geq 0} D_k(t,x) H_k\left(\frac{v}{\sqrt{T_0}}\right) M(v).$$

Hermite system: $0 \leq k \leq N_H - 1$

$$\partial_t D_k + \sqrt{k}\mathcal{A} D_{k-1} - \sqrt{k+1}\mathcal{A}^* D_{k+1} - \sqrt{\frac{k}{T_0}} E D_{k-1} = -\frac{k}{\tau_0} D_k,$$

$$E = -\mathcal{B}\Phi, \quad -\mathcal{B}^* E = D_0 - D_{\infty,0}.$$

where $\mathcal{A} = \sqrt{T_0}\partial_x$, $\mathcal{A}^* = -\sqrt{T_0}\partial_x$, $\mathcal{B} = \partial_x$ and $\mathcal{B}^* = -\partial_x$.

Translation

Linearized free energy

$$\mathcal{E} = \frac{1}{2} \sum_{k=0}^{N_H-1} \|D_k - D_{\infty,k}\|_{L_x^2}^2 + \frac{1}{2T_0} \|E\|_{L_x^2}^2.$$

Dissipation from **collision**:

$$\mathcal{I} = \sum_{k=1}^{N_H-1} k \left\|D_k\right\|_{L_x^2}^2$$

Residual from **field interaction**:

$$\mathcal{R} = \sum_{k=0}^{N_H-1} \sqrt{\frac{k+1}{T_0}} \langle E_h(D_{h,k} - D_{\infty,k}), D_{h,k+1} \rangle_{L_x^2}$$

Spatial discretization: structure-preserving conditions

Semidiscrete scheme

$$\partial_t D_{h,0} - \mathcal{A}_h^* D_{h,1} = 0,$$

$$\partial_t D_{h,1} + \mathcal{A}_h D_{h,0} - \sqrt{2}\mathcal{A}_h^* D_{h,2} + \mathcal{A}_h \Phi_h - \frac{1}{T_0} E_h (D_{h,0} - 1) = -\frac{1}{\tau_0} D_{h,1},$$

$$\partial_t D_{h,k} + \sqrt{k}\mathcal{A}_h D_{h,k-1} - \sqrt{k+1}\mathcal{A}_h^* D_{h,k+1} - \frac{k}{T_0} E_h D_{h,k-1} = -\frac{k}{\tau_0} D_{h,k}, \; k \geq 2$$

$$E_h = -\mathcal{B}_h \Phi_h, \quad -\mathcal{B}_h^* E_h = D_{h,0} - D_{\infty,0}.$$

Preservation of free energy estimate

Sufficient conditions

(H1) Duality: $\langle \mathcal{A}_h u, v \rangle_{L_x^2} = \langle u, \mathcal{A}_h^* v \rangle_{L_x^2}$.

(H2) Kernel: $\mathcal{A}_h 1 = 0$.

(H3) Duality: $\langle \mathcal{B}_h u, v \rangle_{L_x^2} = \langle u, \mathcal{B}_h^* v \rangle_{L_x^2}$.

$$\frac{\mathrm{d}\mathcal{E}_h}{\mathrm{d}t} = -\frac{1}{\tau_0}\mathcal{I}_h + \mathcal{R}_h.$$

Preservation of hypocoercivity

Sufficient conditions: (H1)-(H3) and

(H4) Poincaré inequality:

$$\|u\|_{L_x^2} \lesssim \|\mathcal{A}_h u\|_{L_x^2}.$$

(H5) Primal-dual balance:

$$\|\mathcal{A}_h u\|_{L_x^2} \sim \|\mathcal{A}_h^* u\|_{L_x^2}.$$

(H6) L^∞ regularity:

$$\|E_h\|_{L_x^\infty} \lesssim \|D_{h,0} - D_{\infty,0}\|_{L_x^2}.$$

If $\mathcal{E}_h(0) \leq \lambda^2$, then

$$\mathcal{E}_h(t) \leq 3e^{-\lambda t}\mathcal{E}_h(0), \quad \forall t \geq 0.$$

Temporal discretization: splitting method

Linearized step

$$\frac{D_{h,0}^{n+\frac{1}{2}} - D_{h,0}^{n}}{\Delta t} - \mathcal{A}_h^* D_{h,1}^{n+\frac{1}{2}} = 0,$$

$$\frac{D_{h,1}^{n+\frac{1}{2}} - D_{h,1}^{n}}{\Delta t} + \mathcal{A}_h D_{h,0}^{n+\frac{1}{2}} - \sqrt{2}\mathcal{A}_h^* D_{h,2}^{n+\frac{1}{2}} + \mathcal{A}_h \Phi_h^{n+\frac{1}{2}} = -\frac{1}{\tau_0} D_{h,1}^{n+\frac{1}{2}},$$

$$\frac{D_{h,k}^{n+\frac{1}{2}} - D_{h,k}^{n}}{\Delta t} + \sqrt{k}\mathcal{A}_h D_{h,k-1}^{n+\frac{1}{2}} - \sqrt{k+1}\mathcal{A}_h^* D_{h,k+1}^{n+\frac{1}{2}} = -\frac{k}{\tau_0} D_{h,k}^{n+\frac{1}{2}}, k \geq 2,$$

$$E_h^{n+\frac{1}{2}} = -\mathcal{B}_h \Phi_h^{n+\frac{1}{2}}, \quad -\mathcal{B}_h^* E_h^{n+\frac{1}{2}} = D_{h,0}^{n+\frac{1}{2}} - D_{\infty,0}.$$

Nonlinear step

$$D_{h,0}^{n+1} = D_{h,0}^{n+\frac{1}{2}},$$

$$\frac{D_{h,1}^{n+1} - D_{h,1}^{n+\frac{1}{2}}}{\Delta t} - \frac{1}{T_0} E_h^{n+1}(D_{h,0}^{n+1} - 1) = 0,$$

$$\frac{D_{h,k}^{n+1} - D_{h,k}^{n+\frac{1}{2}}}{\Delta t} - \frac{k}{T_0} E_h^{n+1} D_{h,k-1}^{n+1} = 0, k \geq 2,$$

$$E_h^{n+1} = E_h^{n+\frac{1}{2}}, \ \Phi_h^{n+1} = \Phi_h^{n+\frac{1}{2}}.$$

Linear control

$$\frac{\mathcal{E}_h^{n+\frac{1}{2}} - \mathcal{E}_h^{n}}{\Delta t} \leq -\kappa \min(\tau_0, \tau_0^{-1}) \mathcal{E}_h^{n+\frac{1}{2}}.$$

Quadratic control

$$\frac{\mathcal{E}_h^{n+1} - \mathcal{E}_h^{n+\frac{1}{2}}}{\Delta t} \leq \kappa \max(\tau_0, \tau_0^{-1}) (\mathcal{E}_h^{n+1})^2.$$

Contents

1. Hypocoercivity theory for kinetic equations

2. Discontinuous Galerkin/Hermite spectral methods for VPFP system
2.1 Close-to-equilibrium framework
2.2 Discretization
2.3 Numerical simulations

3. Conclusions and perspectives

Landau damping

Problem setup

Let $x \in \mathbb{T} = [0, L]$, $v \in \mathbb{R}$, and

$$f^{in}(x, v) = \left(1 + \delta \cos \frac{2\pi x}{L}\right) M(v),$$

with length of domain L, amplitude of perturbation δ, and background temperature $T_0 = 1$. In the collisionless limit $\tau_0 \to \infty$, this is the **classical Landau damping** configuration.

Solver setup

- second-order LDG with alternating fluxes for both VFP and Poisson in space;
- **Strang splitting** with second-order DIRK in time;
- **Hou-Li filter** with 2/3 dealiasing;
- collisional regimes $\tau_0 \in \{10^1, 10^3, 10^5\}$.


Landau damping: long-time behavior across collisional regimes

Strong collision

Collision dominates and the solution **relaxes rapidly toward equilibrium**.

Weak collision

Transport dominates for a long transient and generates **filamentation** in phase space.




(a) $\tau_0 = 10^1$ (b) $\tau_0 = 10^3$ (c) $\tau_0 = 10^5$

Figure 1: Strong Landau damping ($\delta = 0.05$): snapshots of the distribution function f at $t \in \{4, 16, 40\}$ across the collisional regimes. $N_x = 128$, $N_H = 640$.


Landau damping: long-time convergence rate

Strong collision


The numerical decay follows the **exponential** trend predicted by the discrete hypocoercive estimate with **close-to-equilibrium** initial data.




(a) $\|E\|_{L^2_x}$



(b) $\left\|f - f_\infty\right\|_{L^2_{x,v}(f_\infty^{-1})}$



(c) $\left\|\rho - \rho_\infty\right\|_{L^2_x(\rho_\infty^{-1})}$



(d) $\|f - \rho\mathcal{M}\|_{L^2_{x,v}(f_\infty^{-1})}$

Figure 2: Strong Landau damping ($\delta = 0.05$): time evolution in logarithmic scale of distances to equilibria across the collisional regimes. $N_x = 128$, $N_H = 640$.

Landau damping: accuracy test

Table 1: L^1, L^2 and L^∞ errors and convergence order, $\delta = 0.05$, $\Delta t = 0.1$, $t_{final} = 1$.

τ_0	$N_x \times N_H$	L^1 error	Order	L^2 error	Order	L^∞ error	Order
10^1	64 × 64	5.52E-03	-	1.82E-03	-	1.10E-03	-
	128 × 128	1.43E-03	1.95	4.70E-04	1.95	2.85E-04	1.96
	256 × 256	3.63E-04	1.98	1.20E-04	1.98	7.20E-05	1.98
10^3	64 × 64	5.87E-03	-	1.96E-03	-	1.23E-03	-
	128 × 128	1.31E-03	2.16	4.45E-04	2.14	2.89E-04	2.09
	256 × 256	2.67E-04	2.29	8.98E-05	2.31	5.62E-05	2.36
10^5	64 × 64	5.87E-03	-	1.96E-03	-	1.23E-03	-
	128 × 128	1.31E-03	2.16	4.44E-04	2.14	2.90E-04	2.09
	256 × 256	2.68E-04	2.29	8.98E-05	2.31	5.64E-05	2.36

Model with non-uniform background density

VPFP Model

$$\partial_t f + v\partial_x f + E\partial_v f = \frac{1}{\tau_0}\partial_v(vf + T_0\partial_v f),$$

$$E = -\partial_x \Phi, -\partial_{xx}\Phi = \rho - \rho_0(x),$$

Let $F = E - E_\infty$, $\Psi = \Phi - \Phi_\infty$, then

$$\partial_t f + v\partial_x f + E_\infty \partial_v f + F\partial_v f = \frac{1}{\tau_0}\partial_v(vf + T_0\partial_v f),$$

$$F = -\partial_x \Psi, -\partial_{xx}\Psi = \rho - \rho_\infty,$$

Gibbs state

Poisson-Boltzmann equation:

$$\rho_\infty = c_\infty \exp(-\frac{\Phi_\infty}{T_0}), \quad -\Delta_x \Phi_\infty = \rho_\infty - \rho_0,$$

where c_∞ is determined by

$$c_\infty \int \exp(-\frac{\Phi_\infty}{T_0})dx = \int \rho_0(x)dx.$$

Landau damping with non-uniform ion density

Problem setup


Let $x \in \mathbb{T} = [-L, L]$ with $L = 6$, $v \in \mathbb{R}$, and consider initial data

$$f^{in}(x, v) = (\rho_\infty(x) + \delta cos(\frac{\pi}{L}x))M(v)),$$

where amplitude of perturbation $\delta = 0.01$ and background temperature $T_0 = 1$. Consider stationary density ρ_∞ be given by

$$\rho_\infty(x) = 0.2 \sin(\frac{\pi}{L}x).$$

Landau damping with non-uniform ion density



(a) $t = 8$ (b) $t = 30$ (c) $t = 70$

Figure 3: Landau damping with non-uniform ion density: snapshots of **distance** between distribution f and the equilibrium f_∞ at $t \in \{8, 30, 70\}$ with $(N_x, N_H) = (128, 400)$. The first line is for strong collision $\tau_0 = 10^2$ (max distance is about 10^{-5} at $t = 70$), while the second line is for weak collision $\tau_0 = 10^4$ (max distance is about 10^{-3} at $t = 70$).

Two-stream instability with non-uniform ion density

Problem setup


Let $x \in \mathbb{T} = [-L, L]$ with $L = 6$, $v \in \mathbb{R}$, and consider initial data

$$f^{in}(x, v) = (\frac{1}{6} + \frac{5}{6}|v|^2)(1 + \delta cos(\frac{\pi}{L}x))M(v),$$

where amplitude of perturbation $\delta = 0.01$ and background temperature $T_0 = 1$. Consider stationary potential Φ_∞ given by

$$\Phi_\infty(x) = 0.1(1 - \cos(\frac{\pi}{L}x)).$$

Two-stream instability with non-uniform ion density



(a) $\tau_0 = 10^2$ (b) $\tau_0 = 10^3$ (c) $\tau_0 = 10^4$

Figure 4: Two stream instability with non-uniform density: snapshots of distribution function f at $t \in \{8, 30, 60\}$ (from top to bottom) across the collisional regimes. $N_x = 128$, $N_H = 800$.

Contents

1. Hypocoercivity theory for kinetic equations
1.1 Overview
1.2 L^2 hypocoercivity
1.3 Discrete hypocoercivity

2. Discontinuous Galerkin/Hermite spectral methods for VPFP system
2.1 Close-to-equilibrium framework
2.2 Discretization
2.3 Numerical simulations

3. Conclusions and perspectives

Conclusions

Hypocoercivity theory:

- a mathematical framework for quantifying convergence to equilibrium in degenerate dissipative systems

Hypocoercivity-preserving DGHS solver for VPFP system:

- sufficient conditions for preserving hypocoercivity at the semidiscrete level.
- provable exponential relaxation for close-to-equilibrium initial data.

Perspectives

Analysis:

- nonlinear BGK equation, radiative transfer equation;
- Boltzmann entropy, φ-entropy;
- characteristic-based methods;

Algorithm:

- High accuracy hypocoercive time integrator;
- Hypocoercive dynamical compression.

References

- C. Villani.
  Hypocoercivity.
  American Mathematical Society, 2009.
- J. Dolbeault, C. Mouhot, and C. Schmeiser.
  Hypocoercivity for linear kinetic equations conserving mass.
  Transactions of the American Mathematical Society, 367(6):3807–3828, 2015.
- A. Blaustein and F. Filbet.
  On a discrete framework of hypocoercivity for kinetic equations.
  Mathematics of Computation, 93(345):163–202, 2024.
- **Y. Cai**, A. Blaustein, T. Xiong, and F. Filbet.
  Discrete hypocoercive estimates for discontinuous Galerkin methods: application to the Vlasov–Poisson–Fokker–Planck system.
  arXiv preprint arXiv:2603.26218, 2026.

Thank you for your attention!