

---

# Suche nach Elektroweakinos mit dem ATLAS Detektor

---



LUDWIG-MAXIMILIANS-UNIVERSITÄT MÜNCHEN  
FAKULTÄT FÜR PHYSIK

DISSERTATION

Eric Schanet  
March 2021

Supervisor: PD Dr. Jeanette Lorenz





## Part I

# Fundamental concepts



## Part II

# The 1-lepton analysis







## Part III

# Reinterpretation



# Chapter 11

## Reinterpretation in the pMSSM

After having discussed to some extent efforts and methods to reinterpret ATLAS searches for Supersymmetry (SUSY), this chapter presents a reinterpretation of the 1-lepton analysis in the phenomenological Minimal Supersymmetric Standard Model (pMSSM). The truth analysis and simplified likelihoods discussed in chapters 9 and 10, respectively, are instrumental for the following sections.

### 11.1 Motivation

In today's searches for beyond the Standard Model (BSM) physics, it is common to use simplified models as a way of avoiding to necessity to deal with high-dimensional parameter spaces that are extremely challenging to sample and compare to data in an exhaustive way. The simplified model approach has also been used in the second part of this work, where results of the interpretation of the 1-lepton analysis in the  $\tilde{\chi}_1^\pm \tilde{\chi}_2^0 \rightarrow Wh\tilde{\chi}_1^0 \tilde{\chi}_1^0$  model have been presented. As has been discussed in section 1.2.7, simplified models are however by no means complete SUSY models and only serve as proxies for more complex and realistic SUSY scenarios. As such, simplified model limits cannot trivially be translated into limits on model parameters of a more complete SUSY model. Large-scale reinterpretations are necessary to understand the constraints today's SUSY searches set on realistic SUSY scenarios.

One class of more complete models, focussing on phenomenologically viable models, is the pMSSM, introduced in section 1.2.6. With its 19 parameters it offers much more complex SUSY scenarios while still being of somewhat manageable dimensionality. Still, large-scale reinterpretations in the pMSSM are computationally challenging and require a set of approximation as those introduced in chapters 9 and 10.

Large-scale reinterpretations in the pMSSM using a collection of relevant ATLAS SUSY searches not only allow to assess the sensitivity of the ATLAS SUSY search program towards more realistic SUSY scenarios, but can also potentially reveal interesting regions of the parameter space not yet covered by the current search programme. Moreover, such reinterpretations allow to demonstrate the sensitivity of simplified model searches beyond the simplified models they are originally interpreted in, thereby justifying the use of simplified models as proxies for more complete SUSY scenarios. In addition, reinterpretations in the pMSSM can be used to connect

the ATLAS SUSY searches with dark matter constraints from non-collider experiments, as well as Higgs and flavour measurements.

Although the following sections will be restricted to a reinterpretation of the 1-lepton search presented in the second part of this thesis, efforts are ongoing in ATLAS to perform large-scale reinterpretations using a majority of the full Run 2 ATLAS SUSY searches. These efforts will most likely result in one of the most comprehensive set of ATLAS constraints on SUSY yetope.

## 11.2 Truth analysis

As discussed in chapter 9, the reinterpretation of an analysis involves re-executing the analysis pipeline in order to derived signal rate estimates in all regions. In large-scale reinterpretations, running a RECAST implementation on all signal models considered is not computationally feasible and instead a *truth-level* analysis is performed for most signal models sampled. The truth-level analysis skips the detector simulation and uses generator-level objects instead. Any detector-level effects and inefficiencies will thus not be reflected in truth-level observables. In order to reproduce the kinematic distributions observed in the full analysis (using reconstruction-level objects), a dedicated *truth smearing*—discussed in detail in section 11.2.2—is applied.

### 11.2.1 Truth selection

All signal and control regions considered in the original 1-lepton search are implemented at truth-level using SIMPLEANALYSIS. The exact implementation is publicly available at Ref. [258] and was already used in chapter 7 for the derivation of a number of the theory uncertainties in the full analysis.

Although a number of object definitions introduced in section 4.4 are not well-defined at truth-level, they are nonetheless specified in the truth-level implementation as the subsequent truth smearing may depend on them. Additionally, as discussed in section 9.1, the full specification of the original analysis event selection allows for simpler reinterpretations by efforts outside of the ATLAS collaboration that generally do not have access to the full original analysis software.

All observables used in the analysis are computed using truth-level quantities. An overlap removal procedure following the same prescription as described for the reconstruction-level analysis is performed, i.e. especially also using the same shrinking cone definitions introduced in section 4.5. Overlap removal step removing electrons sharing a track with a muon is approximated by using a distance parameter of  $\Delta R = 0.01$  between the objects.

### 11.2.2 Truth smearing

The general assumption of the truth smearing applied in the following is that the detector response roughly factorises into the responses of single particles. This allows to use detector performance results provided by the *combined performance groups* in ATLAS in order to construct detector response maps parameterised in different observables for each physics object. Detector response maps include object reconstruction and identification efficiencies as well as scale factors to correct for differences between Monte Carlo (MC) and observed data. Likewise,

effects from the finite resolution of energy measurements in the detector are modelled through energy resolution maps. In the following, the 4-vector components of electrons, muons, jets and  $E_T^{\text{miss}}$  are smeared.

In the case of truth electrons, the identification efficiencies considered are parameterised in  $\eta$  and  $p_T$  as well as the identification working point used. In  $\eta$ , nine fixed-width bins are used. In  $p_T$ , six bins are implemented and a linear interpolation between two adjacent  $p_T$ -bins is used to get the efficiency for the given  $p_T$  of each truth electron. The probability of finding a fake electron in a truth jet is estimated through a similar two-dimensional map depending on the truth jet  $\eta$  and  $p_T$ , again using fixed-width bins in  $\eta$  and linear interpolation in  $p_T$ . The range of the  $p_T$  interpolation for identification efficiencies and fake rates extends from 7 GeV to 120 GeV. If the truth  $p_T$  of the electron is outside of that range, the identification efficiency and fake rate from the respective bound of the corresponding  $\eta$ -bin are used. The probability for misidentifying an electron as a photon is estimated using different fixed values for the barrel and end-cap regions. Finally, the transverse energy of the electron is smeared using a random number drawn from a Gaussian distribution with mean corresponding to the truth value and a standard deviation corresponding to the  $\eta$ - and  $p_T$ -dependent energy resolution.

For truth muons, the identification efficiencies are also parameterised in  $\eta$  and  $p_T$  as well as the identification working point used. Similar to truth electrons, the  $p_T$  of the muon is smeared using a random number drawn from a Gaussian distribution with mean corresponding to the true transverse momentum, and standard deviation corresponding to the momentum resolution. The momentum resolution of combined truth muons is computed from the measured resolutions in the inner detector (ID) ( $\sigma_{\text{ID}}$ ) and muon spectrometer (MS) ( $\sigma_{\text{MS}}$ ) as

$$\sigma_{\text{CB}} = \frac{\sigma_{\text{ID}}\sigma_{\text{MS}}}{\sqrt{\sigma_{\text{ID}}^2 + \sigma_{\text{MS}}^2}}, \quad (11.1)$$

where  $\sigma_{\text{ID}}$  and  $\sigma_{\text{MS}}$  are parameterised in  $\eta$  and  $p_T$ .

The transverse momentum of truth jets is smeared using a Gaussian with standard deviation equal to the jet energy resolution (JER), provided in a map parameterised in five bins in  $\eta$  ranging from  $|\eta| = 0$  to  $|\eta| = 4.5$ . Following [216], jet energy resolutions are provided using parameterisations of a noise  $N$ , stochastic  $S$  and constant  $C$  term for each of the seven bins in  $|\eta|$ , such that the resolution can be computed as

$$\frac{\sigma(p_T)}{p_T} = \frac{N}{p_T} \oplus \frac{S}{\sqrt{p_T}} \oplus C. \quad (11.2)$$

Only truth jets with  $10 \text{ GeV} < p_T < 1.5 \text{ TeV}$  are smeared. For truth jets with  $p_T > 20 \text{ GeV}$ , the flavour tagging efficiency is considered using efficiencies parameterised in  $\eta$ ,  $p_T$  and the MV2c10 working point (introduced in section 4.4) used, measured in fully reconstructed simulated  $t\bar{t}$  events [222].

Finally, the smeared missing transverse energy is computed using the the transverse momenta of all smeared truth objects in the event, including an approximation for the track soft term. The latter is approximated using results from  $Z \rightarrow e^+e^-$  events, allowing to infer a distribution of the mean soft term projected in the direction longitudinal to the total transverse momentum of all hard objects in an event,  $\mathbf{p}_T^{\text{hard}}$ . The measured resolution parallel and perpendicular to  $\mathbf{p}_T^{\text{hard}}$  is then used to smear the nominal soft track value.

## 11.3 Validation of truth analysis

### 11.3.1 Validation at loose preselection

The performance of the truth smearing is illustrated in a loose preselection for a single exemplary benchmark signal point in fig. 11.1. The loose preselection applied requires exactly one lepton,  $E_T^{\text{miss}} > 50$  GeV,  $m_T > 50$  GeV, and 2–3 jets, two of which need to be  $b$ -tagged. The truth-level distributions are compared with the reconstruction-level distributions before and after truth smearing. It can clearly be observed that the truth smearing noticeably improves the agreement between the truth- and reconstruction-level distributions. While the lepton and jet reconstruction and identification efficiencies are—due to their general dependence on  $\eta$ ,  $p_T$  and individual working points—crucial for the overall agreement in shape, the inclusion of flavour-tagging efficiencies significantly improves the overall agreement in normalisation.

Although some minor differences remain, overall a good agreement is observed across all relevant kinematic distributions at loose preselection level. Most of the differences between smeared truth-level and reconstruction-level distributions in individual bins are well within the MC statistical uncertainties arising from the relatively limited MC statistics available.


### 11.3.2 Validation in signal regions

As the expected signal rates in the signal regions are ultimately what is entering the (simplified) likelihood, it is important that the good agreement observed at preselection is still present in the kinematically tight selections of the signal regions. Additionally, it is worth investigating the agreement across all signal models considered in the original analysis. A comparison of the truth-level and reconstruction-level event rates before and after smearing in the signal regions SR-LM, SR-MM and SR-HM is shown in fig. 11.2 for all signal models considered in the 1-lepton analysis. Only the cumulative  $m_{CT}$  bins are shown in each signal region (SR) in fig. 11.2. The agreement in the individual  $m_{CT}$  bins in each SR-LM, SR-MM and SR-HM is shown in figs. D.1 to D.3.

It can clearly be seen that the truth smearing drastically improves the agreement in event rate estimates at truth- and reconstruction-level across all SR bins considered. While the event rates are generally overestimated at truth-level before smearing, compared to reconstruction-level, both tend to agree well within statistical uncertainties after smearing.


### 11.3.3 Validation using simplified likelihood

Using the expected event rates at smeared truth-level for every signal model in the original signal grid considered in the 1-lepton analysis, expected and observed  $CL_s$  values can be computed and exclusion contours can be derived. Figure 11.3 compares the expected and observed exclusion contours obtained using the simplified likelihood and smeared truth-level signal inputs with those obtained using the full likelihood and reconstruction-level signal inputs. Even though—with the simplified likelihood and the smeared truth-level inputs—a significant two-fold approximation is made, the agreement is overall quite good and the published exclusion contour can be reproduced to a relatively high degree of precision.



**Figure 11.1:** Comparisons of the kinematic distributions of key observables at (smeared) truth- and reconstruction-level. The exemplary benchmark signal point with  $m(\tilde{\chi}_1^\pm/\tilde{\chi}_2^0), m(\tilde{\chi}_1^0) = 700, 150$  GeV is shown. The ratio pad shows the ratio between smeared and unsmeared truth-level distributions (green and blue) to reconstruction-level distributions (red). Only MC statistical uncertainty is included in the error bars. All distributions are shown in a loose preselection requiring exactly one lepton,  $E_T^{\text{miss}} > 50$  GeV,  $m_T > 50$  GeV, and 2–3 jets, two of which need to be  $b$ -tagged. The latter requirement is dropped for the  $b$ -jet multiplicity distribution.





**Figure 11.2:** Comparison of the event rates at truth- and reconstruction-level before (left) and after (right) truth smearing. From top to bottom, the SR-LM, SR-MM and SR-HM signal regions are shown, with cumulative (integrated)  $m_{CT}$  bins. Every single point in the scatter plots represents a single signal model considered in the original 1-lepton analysis. Uncertainties include MC statistical uncertainties.




Figure 11.3

In summary, this validation process shows that the signal pipeline in fig. 9.1 can be efficiently approximated using truth-level analysis and a simplified treatment of the statistical model, allowing a considerably faster evaluation of BSM models while still offering reliable results.

## 11.4 Model sampling and processing

### 11.4.1 Sampling

All signal models considered in the following are sampled from the pMSSM using the parameter ranges shown in table 11.1. Flat probability distributions are used to draw random values within the given ranges for each parameter and each unique set of pMSSM parameters generated that way is referred to as an independent SUSY model.

As this work discusses a search for electroweakinos, the SUSY models drawn from the pMSSM are sampled with a special focus on said supersymmetric particles. This is achieved by setting the mass parameters of the first and second generation squarks as well as those of the sleptons to values much higher than those accessible at Large Hadron Collider (LHC) energies, effectively decoupling them. For naturalness arguments, third generation squarks and the gluino are not strictly decoupled but set to sufficiently high values such as not to affect the electroweak sector too much. The lower and upper bounds on the 12 scanned parameters are chosen to yield a high density of models with electroweakino masses accessible at LHC energies.

Once a value for each of the 19 pMSSM parameters has been chosen, a number of publicly available software packages are executed in order to compute the properties of each model point. In a first step, SPHENO v4.0.5 [274, 275] is used to calculate the spectrum of the sparticles. The result of SPHENO is used to determine the masses and mixings of the Higgs bosons using FEYNHIGGS v2.16.0 [276–278], after which SPHENO is re-executed in order to update the rest of the sparticle spectrum. An additional SUSY spectrum calculation is performed with SOFTSUSY v4.1.10 [279]. Although the masses, mixings and branching fractions from SOFTSUSY will not directly be used in the following, the program is still required to complete successfully in order to reduce the number of pMSSM models with pathological properties.

**Table 11.1:** Scan ranges used for each of the 19 pMSSM parameters. For parameters written with a modulus sign, both the positive and negative values are allowed. The term “gen(s)” refers to generation(s).

Parameter	min	max	Note
$m_{\tilde{L}_1} (= m_{\tilde{L}_2})$	10 TeV	10 TeV	Left-handed slepton (first two gens.) mass
$m_{\tilde{e}_1} (= m_{\tilde{e}_2})$	10 TeV	10 TeV	Right-handed slepton (first two gens.) mass
$m_{\tilde{L}_3}$	10 TeV	10 TeV	Left-handed stau doublet mass
$m_{\tilde{e}_3}$	10 TeV	10 TeV	Right-handed stau mass
$m_{\tilde{Q}_1} (= m_{\tilde{Q}_2})$	10 TeV	10 TeV	Left-handed squark (first two gens.) mass
$m_{\tilde{u}_1} (= m_{\tilde{u}_2})$	10 TeV	10 TeV	Right-handed up-type squark (first two gens.) mass
$m_{\tilde{d}_1} (= m_{\tilde{d}_2})$	10 TeV	10 TeV	Right-handed down-type squark (first two gens.) mass
$m_{\tilde{Q}_3}$	2 TeV	5 TeV	Left-handed squark (third gen.) mass
$m_{\tilde{u}_3}$	2 TeV	5 TeV	Right-handed top squark mass
$m_{\tilde{d}_3}$	2 TeV	5 TeV	Right-handed bottom squark mass
$ M_1 $	0 TeV	2 TeV	Bino mass parameter
$ M_2 $	0 TeV	2 TeV	Wino mass parameter
$ \mu $	0 TeV	2 TeV	Bilinear Higgs mass parameter
$M_3$	1 TeV	5 TeV	Gluino mass parameter
$ A_t $	0 TeV	8 TeV	Trilinear top coupling
$ A_b $	0 TeV	2 TeV	Trilinear bottom coupling
$ A_\tau $	0 TeV	2 TeV	Trilinear $\tau$ lepton coupling
$M_A$	0 TeV	5 TeV	Pseudoscalar Higgs boson mass
$\tan \beta$	1	60	Ratio of the Higgs vacuum expectation values

After the complete model spectrum has calculated, additional properties are determined. The dark matter relic abundance of each model is calculated with MICROMEGAS v5.2.1 [280, 281]. Finally, flavour physics and precision electroweak observables like  $\Delta\rho$ ,  $\Delta(g-2)_\mu$ ,  $\text{BR}(b \rightarrow s\gamma)$  and  $\text{BR}(B_s \rightarrow \mu^+\mu^-)$  are determined using GM2CALC v1.7.1 [282] and SUPERISO v4.0 [283].

#### 11.4.2 Selection and processing

A total of 200,000 unique models are sampled from the pMSSM using the above prescription. After requiring all spectrum generators and properties calculations to finish without errors as well as models to produce a dark matter relic density satisfying  $\Omega_c h^2 < 0.12$ , a total of 106,854 models survive. The cross section for all surviving models is computed at next-to-leading order (NLO) using PROSPINO v2.1 [284, 285]. Models with an inclusive cross sections for all electroweak production processes below 0.07 fb are discarded as they would result in less than 10 expected signal events with an integrated luminosity of 139 fb<sup>-1</sup>, not enough to be sensitive to with current electroweak SUSY searches. Models are also discarded for lack of experimental sensitivity if the  $\tilde{\chi}_1^\pm$  is heavier than 1.2 TeV. Next, models are only considered in the following if their lightest supersymmetric particle (LSP) is the  $\tilde{\chi}_1^0$ . Finally, models with long-lived or even stable (on the time scale needed for traversing the ATLAS detector) sparticles<sup>†</sup> are discarded as SUSY searches targeting prompt electroweakino decays (like the 1-lepton search), are not expected to be sensitive to these models.

<sup>†</sup> Not considering the LSP.

Altogether, 90,974 of the pMSSM models sampled satisfy all constraints and are generated at truth-level in a subsequent step.

### 11.4.3 Event generation

Event generation is performed using the software centrally provided by the ATLAS production system. The initial pair of sparticles with two one parton in the Matrix Element (ME) are generated using the MADGRAPH5\_AMC@NLO v2.6.1. [174, 175] generator. Next, PYTHIA8.230 [176] with the A14 tune is used for the hadronisation and Parton Shower (PS), together with the NNPDF 2.3 LO [178] Parton Distribution Function (PDF) set.

### 11.4.4 Truth-level analysis

All models passing event generation are evaluated using the truth-level analysis described in section 11.2.

## 11.5 Results in the pMSSM



## Part IV

# Summary and Outlook







**Part V**

**Appendix**



# Abbreviations

**BSM** beyond the Standard Model. [143](#), [149](#)

**ID** inner detector. [145](#)

**JER** jet energy resolution. [145](#)

**LHC** Large Hadron Collider. [149](#)

**LSP** lightest supersymmetric particle. [150](#)

**MC** Monte Carlo. [144](#), [146–148](#)

**ME** Matrix Element. [151](#)

**MS** muon spectrometer. [145](#)

**NLO** next-to-leading order. [150](#)

**PDF** Parton Distribution Function. [151](#)

**pMSSM** phenomenological Minimal Supersymmetric Standard Model. [143](#), [149–151](#)

**PS** Parton Shower. [151](#)

**SR** signal region. [146](#)

**SUSY** Supersymmetry. [143](#), [144](#), [149](#), [150](#)



# Bibliography

- [1] I. C. Brock and T. Schorner-Sadenius, *Physics at the terascale*. Wiley, Weinheim, 2011. <https://cds.cern.ch/record/1354959>.
- [2] M. E. Peskin and D. V. Schroeder, *An Introduction to quantum field theory*. Addison-Wesley, Reading, USA, 1995. <http://www.slac.stanford.edu/~mpeskin/QFT.html>.
- [3] S. P. Martin, “A Supersymmetry primer,” [arXiv:hep-ph/9709356](https://arxiv.org/abs/hep-ph/9709356) [[hep-ph](#)]. [Adv. Ser. Direct. High Energy Phys.18,1(1998)].
- [4] M. Bustamante, L. Cieri, and J. Ellis, “Beyond the Standard Model for Montaneros,” in *5th CERN - Latin American School of High-Energy Physics*. 11, 2009. [arXiv:0911.4409](https://arxiv.org/abs/0911.4409) [[hep-ph](#)].
- [5] L. Brown, *The Birth of particle physics*. Cambridge University Press, Cambridge Cambridgeshire New York, 1986.
- [6] P. J. Mohr, D. B. Newell, and B. N. Taylor, “CODATA Recommended Values of the Fundamental Physical Constants: 2014,” *Rev. Mod. Phys.* **88** no. 3, (2016) 035009, [arXiv:1507.07956](https://arxiv.org/abs/1507.07956) [[physics.atom-ph](#)].
- [7] P. D. Group, “Review of Particle Physics,” *Progress of Theoretical and Experimental Physics* **2020** no. 8, (08, 2020) , <https://academic.oup.com/ptep/article-pdf/2020/8/083C01/34673722/ptaa104.pdf>. <https://doi.org/10.1093/ptep/ptaa104>. 083C01.
- [8] **Super-Kamiokande** Collaboration, Y. Fukuda *et al.*, “Evidence for oscillation of atmospheric neutrinos,” *Phys. Rev. Lett.* **81** (1998) 1562–1567, [arXiv:hep-ex/9807003](https://arxiv.org/abs/hep-ex/9807003) [[hep-ex](#)].
- [9] Z. Maki, M. Nakagawa, and S. Sakata, “Remarks on the unified model of elementary particles,” *Prog. Theor. Phys.* **28** (1962) 870–880. [,34(1962)].
- [10] N. Cabibbo, “Unitary symmetry and leptonic decays,” *Phys. Rev. Lett.* **10** (Jun, 1963) 531–533. <https://link.aps.org/doi/10.1103/PhysRevLett.10.531>.
- [11] M. Kobayashi and T. Maskawa, “CP Violation in the Renormalizable Theory of Weak Interaction,” *Prog. Theor. Phys.* **49** (1973) 652–657.
- [12] E. Noether and M. A. Tavel, “Invariant variation problems,” [arXiv:physics/0503066](https://arxiv.org/abs/physics/0503066).
- [13] J. C. Ward, “An identity in quantum electrodynamics,” *Phys. Rev.* **78** (Apr, 1950) 182–182. <https://link.aps.org/doi/10.1103/PhysRev.78.182>.

- [14] Y. Takahashi, “On the generalized ward identity,” *Il Nuovo Cimento (1955-1965)* **6** no. 2, (Aug, 1957) 371–375. <https://doi.org/10.1007/BF02832514>.
- [15] G. 'tHooft, “Renormalization of massless yang-mills fields,” *Nuclear Physics B* **33** no. 1, (1971) 173 – 199. <http://www.sciencedirect.com/science/article/pii/0550321371903956>.
- [16] J. Taylor, “Ward identities and charge renormalization of the yang-mills field,” *Nuclear Physics B* **33** no. 2, (1971) 436 – 444. <http://www.sciencedirect.com/science/article/pii/0550321371902975>.
- [17] A. A. Slavnov, “Ward identities in gauge theories,” *Theoretical and Mathematical Physics* **10** no. 2, (Feb, 1972) 99–104. <https://doi.org/10.1007/BF01090719>.
- [18] C. N. Yang and R. L. Mills, “Conservation of isotopic spin and isotopic gauge invariance,” *Phys. Rev.* **96** (Oct, 1954) 191–195. <https://link.aps.org/doi/10.1103/PhysRev.96.191>.
- [19] K. G. Wilson, “Confinement of quarks,” *Phys. Rev. D* **10** (Oct, 1974) 2445–2459. <https://link.aps.org/doi/10.1103/PhysRevD.10.2445>.
- [20] T. DeGrand and C. DeTar, *Lattice Methods for Quantum Chromodynamics*. World Scientific, Singapore, 2006. <https://cds.cern.ch/record/1055545>.
- [21] S. L. Glashow, “Partial-symmetries of weak interactions,” *Nuclear Physics* **22** no. 4, (1961) 579 – 588. <http://www.sciencedirect.com/science/article/pii/0029558261904692>.
- [22] S. Weinberg, “A model of leptons,” *Phys. Rev. Lett.* **19** (Nov, 1967) 1264–1266. <https://link.aps.org/doi/10.1103/PhysRevLett.19.1264>.
- [23] A. Salam and J. C. Ward, “Weak and electromagnetic interactions,” *Il Nuovo Cimento (1955-1965)* **11** no. 4, (Feb, 1959) 568–577. <https://doi.org/10.1007/BF02726525>.
- [24] C. S. Wu, E. Ambler, R. W. Hayward, D. D. Hoppes, and R. P. Hudson, “Experimental test of parity conservation in beta decay,” *Phys. Rev.* **105** (Feb, 1957) 1413–1415. <https://link.aps.org/doi/10.1103/PhysRev.105.1413>.
- [25] M. Gell-Mann, “The interpretation of the new particles as displaced charge multiplets,” *Il Nuovo Cimento (1955-1965)* **4** no. 2, (Apr, 1956) 848–866. <https://doi.org/10.1007/BF02748000>.
- [26] K. Nishijima, “Charge Independence Theory of V Particles\*,” *Progress of Theoretical Physics* **13** no. 3, (03, 1955) 285–304, <https://academic.oup.com/ptp/article-pdf/13/3/285/5425869/13-3-285.pdf>. <https://doi.org/10.1143/PTP.13.285>.
- [27] T. Nakano and K. Nishijima, “Charge Independence for V-particles\*,” *Progress of Theoretical Physics* **10** no. 5, (11, 1953) 581–582, <https://academic.oup.com/ptp/article-pdf/10/5/581/5364926/10-5-581.pdf>. <https://doi.org/10.1143/PTP.10.581>.
- [28] F. Englert and R. Brout, “Broken symmetry and the mass of gauge vector mesons,” *Phys. Rev. Lett.* **13** (Aug, 1964) 321–323. <https://link.aps.org/doi/10.1103/PhysRevLett.13.321>.
- [29] P. W. Higgs, “Broken symmetries and the masses of gauge bosons,” *Phys. Rev. Lett.* **13** (Oct, 1964) 508–509. <https://link.aps.org/doi/10.1103/PhysRevLett.13.508>.

- [30] P. W. Higgs, “Spontaneous symmetry breakdown without massless bosons,” *Phys. Rev.* **145** (May, 1966) 1156–1163. <https://link.aps.org/doi/10.1103/PhysRev.145.1156>.
- [31] Y. Nambu, “Quasiparticles and Gauge Invariance in the Theory of Superconductivity,” *Phys. Rev.* **117** (1960) 648–663. [[132\(1960\)](#)].
- [32] J. Goldstone, “Field Theories with Superconductor Solutions,” *Nuovo Cim.* **19** (1961) 154–164.
- [33] V. Brdar, A. J. Helmboldt, S. Iwamoto, and K. Schmitz, “Type-I Seesaw as the Common Origin of Neutrino Mass, Baryon Asymmetry, and the Electroweak Scale,” *Phys. Rev. D* **100** (2019) 075029, [arXiv:1905.12634 \[hep-ph\]](#).
- [34] G. ’t Hooft and M. Veltman, “Regularization and renormalization of gauge fields,” *Nuclear Physics B* **44** no. 1, (1972) 189 – 213. <http://www.sciencedirect.com/science/article/pii/0550321372902799>.
- [35] F. Zwicky, “Die Rotverschiebung von extragalaktischen Nebeln,” *Helv. Phys. Acta* **6** (1933) 110–127. <https://cds.cern.ch/record/437297>.
- [36] V. C. Rubin and W. K. Ford, Jr., “Rotation of the Andromeda Nebula from a Spectroscopic Survey of Emission Regions,” *Astrophys. J.* **159** (1970) 379–403.
- [37] G. Bertone, D. Hooper, and J. Silk, “Particle dark matter: Evidence, candidates and constraints,” *Phys. Rept.* **405** (2005) 279–390, [arXiv:hep-ph/0404175](#).
- [38] D. Clowe, M. Bradac, A. H. Gonzalez, M. Markevitch, S. W. Randall, C. Jones, and D. Zaritsky, “A direct empirical proof of the existence of dark matter,” *Astrophys. J.* **648** (2006) L109–L113, [arXiv:astro-ph/0608407 \[astro-ph\]](#).
- [39] A. Taylor, S. Dye, T. J. Broadhurst, N. Benitez, and E. van Kampen, “Gravitational lens magnification and the mass of abell 1689,” *Astrophys. J.* **501** (1998) 539, [arXiv:astro-ph/9801158](#).
- [40] C. Bennett *et al.*, “Four year COBE DMR cosmic microwave background observations: Maps and basic results,” *Astrophys. J. Lett.* **464** (1996) L1–L4, [arXiv:astro-ph/9601067](#).
- [41] G. F. Smoot *et al.*, “Structure in the COBE Differential Microwave Radiometer First-Year Maps,” *ApJS* **396** (September, 1992) L1.
- [42] **WMAP** Collaboration, “Nine-year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Final Maps and Results,” *ApJS* **208** no. 2, (October, 2013) 20, [arXiv:1212.5225 \[astro-ph.CO\]](#).
- [43] **WMAP** Collaboration, “Nine-year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological Parameter Results,” *ApJS* **208** no. 2, (October, 2013) 19, [arXiv:1212.5226 \[astro-ph.CO\]](#).
- [44] **Planck** Collaboration, “Planck 2018 results. I. Overview and the cosmological legacy of Planck,” *Astron. Astrophys.* **641** (2020) A1, [arXiv:1807.06205 \[astro-ph.CO\]](#).
- [45] A. Liddle, *An introduction to modern cosmology; 3rd ed.* Wiley, Chichester, Mar, 2015. <https://cds.cern.ch/record/1976476>.
- [46] **Planck** Collaboration, “Planck 2018 results. VI. Cosmological parameters,” *Astron. Astrophys.* **641** (2020) A6, [arXiv:1807.06209 \[astro-ph.CO\]](#).

- [47] H. Georgi and S. L. Glashow, “Unity of all elementary-particle forces,” *Phys. Rev. Lett.* **32** (Feb, 1974) 438–441. <https://link.aps.org/doi/10.1103/PhysRevLett.32.438>.
- [48] I. Aitchison, *Supersymmetry in Particle Physics. An Elementary Introduction*. Cambridge University Press, Cambridge, 2007.
- [49] **Muon g-2** Collaboration, G. Bennett *et al.*, “Final Report of the Muon E821 Anomalous Magnetic Moment Measurement at BNL,” *Phys. Rev. D* **73** (2006) 072003, [arXiv:hep-ex/0602035](https://arxiv.org/abs/hep-ex/0602035).
- [50] H. Baer and X. Tata, *Weak Scale Supersymmetry: From Superfields to Scattering Events*. Cambridge University Press, 2006.
- [51] A. Czarnecki and W. J. Marciano, “The Muon anomalous magnetic moment: A Harbinger for ‘new physics’,” *Phys. Rev. D* **64** (2001) 013014, [arXiv:hep-ph/0102122](https://arxiv.org/abs/hep-ph/0102122).
- [52] J. L. Feng and K. T. Matchev, “Supersymmetry and the anomalous magnetic moment of the muon,” *Phys. Rev. Lett.* **86** (2001) 3480–3483, [arXiv:hep-ph/0102146](https://arxiv.org/abs/hep-ph/0102146).
- [53] S. Coleman and J. Mandula, “All possible symmetries of the  $s$  matrix,” *Phys. Rev.* **159** (Jul, 1967) 1251–1256. <https://link.aps.org/doi/10.1103/PhysRev.159.1251>.
- [54] R. Haag, J. T. Lopuszanski, and M. Sohnius, “All Possible Generators of Supersymmetries of the  $s$  Matrix,” *Nucl. Phys.* **B88** (1975) 257. [,257(1974)].
- [55] J. Wess and B. Zumino, “Supergauge transformations in four dimensions,” *Nuclear Physics B* **70** no. 1, (1974) 39 – 50. <http://www.sciencedirect.com/science/article/pii/0550321374903551>.
- [56] H. Georgi and S. L. Glashow, “Gauge theories without anomalies,” *Phys. Rev. D* **6** (Jul, 1972) 429–431. <https://link.aps.org/doi/10.1103/PhysRevD.6.429>.
- [57] S. Dimopoulos and D. W. Sutter, “The Supersymmetric flavor problem,” *Nucl. Phys. B* **452** (1995) 496–512, [arXiv:hep-ph/9504415](https://arxiv.org/abs/hep-ph/9504415).
- [58] **MEG** Collaboration, T. Mori, “Final Results of the MEG Experiment,” *Nuovo Cim. C* **39** no. 4, (2017) 325, [arXiv:1606.08168](https://arxiv.org/abs/1606.08168) [[hep-ex](#)].
- [59] H. P. Nilles, “Supersymmetry, Supergravity and Particle Physics,” *Phys. Rept.* **110** (1984) 1–162.
- [60] A. Lahanas and D. Nanopoulos, “The road to no-scale supergravity,” *Physics Reports* **145** no. 1, (1987) 1 – 139. <http://www.sciencedirect.com/science/article/pii/0370157387900342>.
- [61] J. L. Feng, A. Rajaraman, and F. Takayama, “Superweakly interacting massive particles,” *Phys. Rev. Lett.* **91** (2003) 011302, [arXiv:hep-ph/0302215](https://arxiv.org/abs/hep-ph/0302215).
- [62] **Super-Kamiokande** Collaboration, K. Abe *et al.*, “Search for proton decay via  $p \rightarrow e^+ \pi^0$  and  $p \rightarrow \mu^+ \pi^0$  in 0.31 megaton-years exposure of the Super-Kamiokande water Cherenkov detector,” *Phys. Rev. D* **95** no. 1, (2017) 012004, [arXiv:1610.03597](https://arxiv.org/abs/1610.03597) [[hep-ex](#)].
- [63] J. R. Ellis, “Beyond the standard model for hill walkers,” in *1998 European School of High-Energy Physics*, pp. 133–196. 8, 1998. [arXiv:hep-ph/9812235](https://arxiv.org/abs/hep-ph/9812235).



- [64] J. R. Ellis, J. Hagelin, D. V. Nanopoulos, K. A. Olive, and M. Srednicki, “Supersymmetric Relics from the Big Bang,” *Nucl. Phys. B* **238** (1984) 453–476.
- [65] D. O. Caldwell, R. M. Eisberg, D. M. Grumm, M. S. Witherell, B. Sadoulet, F. S. Goulding, and A. R. Smith, “Laboratory limits on galactic cold dark matter,” *Phys. Rev. Lett.* **61** (Aug, 1988) 510–513. <https://link.aps.org/doi/10.1103/PhysRevLett.61.510>.
- [66] M. Mori, M. M. Nojiri, K. S. Hirata, K. Kihara, Y. Oyama, A. Suzuki, K. Takahashi, M. Yamada, H. Takei, M. Koga, K. Miyano, H. Miyata, Y. Fukuda, T. Hayakawa, K. Inoue, T. Ishida, T. Kajita, Y. Koshio, M. Nakahata, K. Nakamura, A. Sakai, N. Sato, M. Shiozawa, J. Suzuki, Y. Suzuki, Y. Totsuka, M. Koshihara, K. Nishijima, T. Kajimura, T. Suda, A. T. Suzuki, T. Hara, Y. Nagashima, M. Takita, H. Yokoyama, A. Yoshimoto, K. Kaneyuki, Y. Takeuchi, T. Tanimori, S. Tasaka, and K. Nishikawa, “Search for neutralino dark matter heavier than the  $w$  boson at kamiokande,” *Phys. Rev. D* **48** (Dec, 1993) 5505–5518. <https://link.aps.org/doi/10.1103/PhysRevD.48.5505>.
- [67] **CDMS Collaboration**, D. S. Akerib *et al.*, “Exclusion limits on the WIMP-nucleon cross section from the first run of the Cryogenic Dark Matter Search in the Soudan Underground Laboratory,” *Phys. Rev. D* **72** (2005) 052009, [arXiv:astro-ph/0507190](https://arxiv.org/abs/hep-ph/0507190).
- [68] A. Djouadi, J.-L. Kneur, and G. Moultaka, “SuSpect: A Fortran code for the supersymmetric and Higgs particle spectrum in the MSSM,” *Comput. Phys. Commun.* **176** (2007) 426–455, [arXiv:hep-ph/0211331](https://arxiv.org/abs/hep-ph/0211331).
- [69] C. F. Berger, J. S. Gainer, J. L. Hewett, and T. G. Rizzo, “Supersymmetry without prejudice,” *Journal of High Energy Physics* **2009** no. 02, (Feb, 2009) 023–023. <http://dx.doi.org/10.1088/1126-6708/2009/02/023>.
- [70] J. Alwall, P. Schuster, and N. Toro, “Simplified Models for a First Characterization of New Physics at the LHC,” *Phys. Rev. D* **79** (2009) 075020, [arXiv:0810.3921](https://arxiv.org/abs/0810.3921) [hep-ph].
- [71] **LHC New Physics Working Group Collaboration**, D. Alves, “Simplified Models for LHC New Physics Searches,” *J. Phys. G* **39** (2012) 105005, [arXiv:1105.2838](https://arxiv.org/abs/1105.2838) [hep-ph].
- [72] D. S. Alves, E. Izaguirre, and J. G. Wacker, “Where the Sidewalk Ends: Jets and Missing Energy Search Strategies for the 7 TeV LHC,” *JHEP* **10** (2011) 012, [arXiv:1102.5338](https://arxiv.org/abs/1102.5338) [hep-ph].
- [73] F. Ambrogio, S. Kraml, S. Kulkarni, U. Laa, A. Lessa, and W. Waltenberger, “On the coverage of the pMSSM by simplified model results,” *Eur. Phys. J. C* **78** no. 3, (2018) 215, [arXiv:1707.09036](https://arxiv.org/abs/1707.09036) [hep-ph].
- [74] O. Buchmueller and J. Marrouche, “Universal mass limits on gluino and third-generation squarks in the context of Natural-like SUSY spectra,” *Int. J. Mod. Phys. A* **29** no. 06, (2014) 1450032, [arXiv:1304.2185](https://arxiv.org/abs/1304.2185) [hep-ph].
- [75] **ATLAS Collaboration**, M. Aaboud *et al.*, “Dark matter interpretations of ATLAS searches for the electroweak production of supersymmetric particles in  $\sqrt{s} = 8$  TeV proton-proton collisions,” *JHEP* **09** (2016) 175, [arXiv:1608.00872](https://arxiv.org/abs/1608.00872) [hep-ex].
- [76] **ATLAS Collaboration**, “Summary of the ATLAS experiment’s sensitivity to supersymmetry after LHC Run 1 — interpreted in the phenomenological MSSM,” *JHEP* **10** (2015) 134, [arXiv:1508.06608](https://arxiv.org/abs/1508.06608) [hep-ex].

- [77] **ATLAS** Collaboration, “Mass reach of the atlas searches for supersymmetry.” [https://atlas.web.cern.ch/Atlas/GROUPS/PHYSICS/PUBNOTES/ATL-PHYS-PUB-2020-020/fig\\_23.png](https://atlas.web.cern.ch/Atlas/GROUPS/PHYSICS/PUBNOTES/ATL-PHYS-PUB-2020-020/fig_23.png), 2020.
- [78] **CMS** Collaboration, “Summary plot moriond 2017.” [https://twiki.cern.ch/twiki/pub/CMSPublic/SUSYSummary2017/Moriond2017\\_BarPlot.pdf](https://twiki.cern.ch/twiki/pub/CMSPublic/SUSYSummary2017/Moriond2017_BarPlot.pdf), 2017.
- [79] L. S. W. Group, “Notes lepsusywg/02-04.1 and lepsusywg/01-03.1.” <http://lepsusy.web.cern.ch/lepsusy/>, 2004. Accessed: 2021-02-11.
- [80] **ATLAS** Collaboration, G. Aad *et al.*, “Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC,” *Phys. Lett. B* **716** (2012) 1–29, [arXiv:1207.7214 \[hep-ex\]](#).
- [81] **CMS** Collaboration, S. Chatrchyan *et al.*, “Observation of a New Boson at a Mass of 125 GeV with the CMS Experiment at the LHC,” *Phys. Lett. B* **716** (2012) 30–61, [arXiv:1207.7235 \[hep-ex\]](#).
- [82] CERN, “About cern.” <https://home.cern/about>. Accessed: 2021-01-21.
- [83] CERN, “CERN Annual report 2019,” tech. rep., CERN, Geneva, 2020. <https://cds.cern.ch/record/2723123>.
- [84] O. S. Bruning, P. Collier, P. Lebrun, S. Myers, R. Ostojic, J. Poole, and P. Proudlock, *LHC Design Report*. CERN Yellow Reports: Monographs. CERN, Geneva, 2004. <https://cds.cern.ch/record/782076>.
- [85] M. Blewett and N. Vogt-Nilsen, “Proceedings of the 8th international conference on high-energy accelerators, cern 1971. conference held at geneva, 20–24 september 1971,” tech. rep., 1971, 1971.
- [86] L. R. Evans and P. Bryant, “LHC Machine,” *JINST* **3** (2008) S08001. 164 p. <http://cds.cern.ch/record/1129806>. This report is an abridged version of the LHC Design Report (CERN-2004-003).
- [87] R. Scrivens, M. Kronberger, D. Küchler, J. Lettry, C. Mastrostefano, O. Midttun, M. O’Neil, H. Pereira, and C. Schmitzer, “Overview of the status and developments on primary ion sources at CERN\*,” <https://cds.cern.ch/record/1382102>.
- [88] M. Vretenar, J. Vollaie, R. Scrivens, C. Rossi, F. Roncarolo, S. Ramberger, U. Raich, B. Puccio, D. Nisbet, R. Mompo, S. Mathot, C. Martin, L. A. Lopez-Hernandez, A. Lombardi, J. Lettry, J. B. Lallement, I. Kozsar, J. Hansen, F. Gerigk, A. Funken, J. F. Fuchs, N. Dos Santos, M. Calviani, M. Buzio, O. Brunner, Y. Body, P. Baudrenghien, J. Bauche, and T. Zickler, *Linac4 design report*, vol. 6 of *CERN Yellow Reports: Monographs*. CERN, Geneva, 2020. <https://cds.cern.ch/record/2736208>.
- [89] E. Mobs, “The CERN accelerator complex - 2019. Complexe des accélérateurs du CERN - 2019,” <https://cds.cern.ch/record/2684277>. General Photo.
- [90] **ATLAS** Collaboration, “The ATLAS Experiment at the CERN Large Hadron Collider,” *JINST* **3** (2008) S08003.
- [91] **CMS** Collaboration, S. Chatrchyan *et al.*, “The CMS Experiment at the CERN LHC,” *JINST* **3** (2008) S08004.

- [92] **ALICE** Collaboration, K. Aamodt *et al.*, “The ALICE experiment at the CERN LHC,” *JINST* **3** (2008) S08002.
- [93] **LHCb** Collaboration, J. Alves, A. Augusto *et al.*, “The LHCb Detector at the LHC,” *JINST* **3** (2008) S08005.
- [94] **TOTEM** Collaboration, G. Anelli *et al.*, “The TOTEM experiment at the CERN Large Hadron Collider,” *JINST* **3** (2008) S08007.
- [95] **LHCf** Collaboration, O. Adriani *et al.*, “Technical design report of the LHCf experiment: Measurement of photons and neutral pions in the very forward region of LHC,”.
- [96] **MoEDAL** Collaboration, J. Pinfold *et al.*, “Technical Design Report of the MoEDAL Experiment,”.
- [97] **ATLAS** Collaboration, “ATLAS Public Results - Luminosity Public Results Run 2,”. <https://twiki.cern.ch/twiki/bin/view/AtlasPublic/LuminosityPublicResultsRun2>. Accessed: 2021-01-17.
- [98] **ATLAS** Collaboration, Z. Marshall, “Simulation of Pile-up in the ATLAS Experiment,” *J. Phys. Conf. Ser.* **513** (2014) 022024.
- [99] “First beam in the LHC - accelerating science,”. <https://home.cern/news/news/accelerators/record-luminosity-well-done-lhc>. Accessed: 2021-01-10.
- [100] **ATLAS Collaboration** Collaboration, “Luminosity determination in  $pp$  collisions at  $\sqrt{s} = 13$  TeV using the ATLAS detector at the LHC,” Tech. Rep. ATLAS-CONF-2019-021, CERN, Geneva, Jun, 2019. <https://cds.cern.ch/record/2677054>.
- [101] **ATLAS** Collaboration, M. Aaboud *et al.*, “Luminosity determination in  $pp$  collisions at  $\sqrt{s} = 8$  TeV using the ATLAS detector at the LHC,” *Eur. Phys. J. C* **76** no. 12, (2016) 653, [arXiv:1608.03953](https://arxiv.org/abs/1608.03953) [hep-ex].
- [102] G. Avoni, M. Bruschi, G. Cabras, D. Caforio, N. Dehghanian, A. Floderus, B. Giacobbe, F. Giannuzzi, F. Giorgi, P. Grafström, V. Hedberg, F. L. Manghi, S. Meneghini, J. Pinfold, E. Richards, C. Sbarra, N. S. Cesari, A. Sbrizzi, R. Soluk, G. Uccielli, S. Valentinetti, O. Viazlo, M. Villa, C. Vittori, R. Vuillermet, and A. Zoccoli, “The new LUCID-2 detector for luminosity measurement and monitoring in ATLAS,” *Journal of Instrumentation* **13** no. 07, (Jul, 2018) P07017–P07017. <https://doi.org/10.1088/1748-0221/13/07/p07017>.
- [103] S. van der Meer, “Calibration of the effective beam height in the ISR,” Tech. Rep. CERN-ISR-PO-68-31. ISR-PO-68-31, CERN, Geneva, 1968. <https://cds.cern.ch/record/296752>.
- [104] P. Grafström and W. Kozanecki, “Luminosity determination at proton colliders,” *Progress in Particle and Nuclear Physics* **81** (2015) 97 – 148. <http://www.sciencedirect.com/science/article/pii/S0146641014000878>.
- [105] “New schedule for CERN’s accelerators and experiments,”. <https://home.cern/news/press-release/cern/first-beam-lhc-accelerating-science>. Accessed: 2021-01-10.

- [106] **ATLAS Collaboration**, G. Aad *et al.*, “Luminosity Determination in  $pp$  Collisions at  $\sqrt{s} = 7$  TeV Using the ATLAS Detector at the LHC,” *Eur. Phys. J. C* **71** (2011) 1630, [arXiv:1101.2185 \[hep-ex\]](#).
- [107] **ATLAS Collaboration** Collaboration, G. Aad *et al.*, “Improved luminosity determination in  $pp$  collisions at  $\sqrt{s} = 7$  TeV using the ATLAS detector at the LHC. Improved luminosity determination in  $pp$  collisions at  $\sqrt{s} = 7$  TeV using the ATLAS detector at the LHC,” *Eur. Phys. J. C* **73** no. CERN-PH-EP-2013-026. CERN-PH-EP-2013-026, (Feb, 2013) 2518. 27 p. <https://cds.cern.ch/record/1517411>. Comments: 26 pages plus author list (39 pages total), 17 figures, 9 tables, submitted to EPJC, All figures are available at [CERN-PH-EP-2013-026](#).
- [108] “Record luminosity: well done LHC,” <https://home.cern/news/news/accelerators/new-schedule-cerns-accelerators-and-experiments>. Accessed: 2021-01-10.
- [109] A. G., B. A. I., B. O., F. P., L. M., R. L., and T. L., *High-Luminosity Large Hadron Collider (HL-LHC): Technical Design Report V. 0.1*. CERN Yellow Reports: Monographs. CERN, Geneva, 2017. <https://cds.cern.ch/record/2284929>.
- [110] J. Pequeno, “Computer generated image of the whole ATLAS detector.” Mar, 2008.
- [111] **ATLAS Collaboration**, “ATLAS: Detector and physics performance technical design report. Volume 1,”.
- [112] J. Pequeno, “Computer generated image of the ATLAS inner detector.” Mar, 2008.
- [113] **ATLAS Collaboration** Collaboration, K. Potamianos, “The upgraded Pixel detector and the commissioning of the Inner Detector tracking of the ATLAS experiment for Run-2 at the Large Hadron Collider,” Tech. Rep. ATL-PHYS-PROC-2016-104, CERN, Geneva, Aug, 2016. <https://cds.cern.ch/record/2209070>. 15 pages, EPS-HEP 2015 Proceedings.
- [114] **ATLAS IBL Collaboration**, B. Abbott *et al.*, “Production and Integration of the ATLAS Insertable B-Layer,” *JINST* **13** no. 05, (2018) T05008, [arXiv:1803.00844 \[physics.ins-det\]](#).
- [115] **ATLAS Collaboration**, “ATLAS Insertable B-Layer Technical Design Report,” Tech. Rep. CERN-LHCC-2010-013. ATLAS-TDR-19, Sep, 2010. <http://cds.cern.ch/record/1291633>.
- [116] **ATLAS Collaboration**, G. Aad *et al.*, “ATLAS b-jet identification performance and efficiency measurement with  $t\bar{t}$  events in  $pp$  collisions at  $\sqrt{s} = 13$  TeV,” *Eur. Phys. J. C* **79** no. 11, (2019) 970, [arXiv:1907.05120 \[hep-ex\]](#).
- [117] **ATLAS Collaboration**, “Particle Identification Performance of the ATLAS Transition Radiation Tracker.” ATLAS-CONF-2011-128, 2011. <https://cds.cern.ch/record/1383793>.
- [118] J. Pequeno, “Computer Generated image of the ATLAS calorimeter.” Mar, 2008.
- [119] J. Pequeno, “Computer generated image of the ATLAS Muons subsystem.” Mar, 2008.
- [120] S. Lee, M. Livan, and R. Wigmans, “Dual-Readout Calorimetry,” *Rev. Mod. Phys.* **90** no. 2, (Dec, 2017) 025002. 40 p. <https://cds.cern.ch/record/2637852>. 44 pages, 53 figures, accepted for publication in Review of Modern Physics.

- [121] M. Leite, “Performance of the ATLAS Zero Degree Calorimeter,” Tech. Rep. ATL-FWD-PROC-2013-001, CERN, Geneva, Nov, 2013.  
<https://cds.cern.ch/record/1628749>.
- [122] S. Abdel Khalek *et al.*, “The ALFA Roman Pot Detectors of ATLAS,” *JINST* **11** no. 11, (2016) P11013, [arXiv:1609.00249](https://arxiv.org/abs/1609.00249) [[physics.ins-det](#)].
- [123] U. Amaldi, G. Cocconi, A. Diddens, R. Dobinson, J. Dorenbosch, W. Duinker, D. Gustavson, J. Meyer, K. Potter, A. Wetherell, A. Baroncelli, and C. Bosio, “The real part of the forward proton proton scattering amplitude measured at the cern intersecting storage rings,” *Physics Letters B* **66** no. 4, (1977) 390 – 394.  
<http://www.sciencedirect.com/science/article/pii/0370269377900223>.
- [124] L. Adamczyk, E. Banaś, A. Brandt, M. Bruschi, S. Grinstein, J. Lange, M. Rijssenbeek, P. Sicho, R. Staszewski, T. Sykora, M. Trzebiński, J. Chwastowski, and K. Korcyl, “Technical Design Report for the ATLAS Forward Proton Detector,” Tech. Rep. CERN-LHCC-2015-009. ATLAS-TDR-024, May, 2015.  
<https://cds.cern.ch/record/2017378>.
- [125] **ATLAS Collaboration**, A. R. Martínez, “The Run-2 ATLAS Trigger System,” *J. Phys. Conf. Ser.* **762** no. 1, (2016) 012003.
- [126] **ATLAS Collaboration** Collaboration, *ATLAS level-1 trigger: Technical Design Report*. Technical Design Report ATLAS. CERN, Geneva, 1998.  
<https://cds.cern.ch/record/381429>.
- [127] **ATLAS Collaboration**, G. Aad *et al.*, “Operation of the ATLAS trigger system in Run 2,” *JINST* **15** no. 10, (2020) P10004, [arXiv:2007.12539](https://arxiv.org/abs/2007.12539) [[physics.ins-det](#)].
- [128] **ATLAS Collaboration** Collaboration, P. Jenni, M. Nessi, M. Nordberg, and K. Smith, *ATLAS high-level trigger, data-acquisition and controls: Technical Design Report*. Technical Design Report ATLAS. CERN, Geneva, 2003.  
<https://cds.cern.ch/record/616089>.
- [129] **ATLAS Collaboration**, G. Aad *et al.*, “The ATLAS Simulation Infrastructure,” *Eur. Phys. J. C* **70** (2010) 823–874, [arXiv:1005.4568](https://arxiv.org/abs/1005.4568) [[physics.ins-det](#)].
- [130] T. Gleisberg, S. Hoeche, F. Krauss, M. Schonherr, S. Schumann, F. Siegert, and J. Winter, “Event generation with SHERPA 1.1,” *JHEP* **02** (2009) 007, [arXiv:0811.4622](https://arxiv.org/abs/0811.4622) [[hep-ph](#)].
- [131] A. Buckley *et al.*, “General-purpose event generators for LHC physics,” *Phys. Rept.* **504** (2011) 145–233, [arXiv:1101.2599](https://arxiv.org/abs/1101.2599) [[hep-ph](#)].
- [132] V. N. Gribov and L. N. Lipatov, “Deep inelastic e p scattering in perturbation theory,” *Sov. J. Nucl. Phys.* **15** (1972) 438–450.
- [133] J. Blumlein, T. Doyle, F. Hautmann, M. Klein, and A. Vogt, “Structure functions in deep inelastic scattering at HERA,” in *Workshop on Future Physics at HERA (To be followed by meetings 7-9 Feb and 30-31 May 1996 at DESY)*. 9, 1996. [arXiv:hep-ph/9609425](https://arxiv.org/abs/hep-ph/9609425).
- [134] A. Buckley, J. Ferrando, S. Lloyd, K. Nordström, B. Page, M. Rüfenacht, M. Schönherr, and G. Watt, “LHAPDF6: parton density access in the LHC precision era,” *Eur. Phys. J. C* **75** (2015) 132, [arXiv:1412.7420](https://arxiv.org/abs/1412.7420) [[hep-ph](#)].

- [135] M. Bengtsson and T. Sjostrand, “Coherent Parton Showers Versus Matrix Elements: Implications of PETRA - PEP Data,” *Phys. Lett. B* **185** (1987) 435.
- [136] S. Catani, F. Krauss, R. Kuhn, and B. R. Webber, “QCD matrix elements + parton showers,” *JHEP* **11** (2001) 063, [arXiv:hep-ph/0109231](#).
- [137] L. Lonnblad, “Correcting the color dipole cascade model with fixed order matrix elements,” *JHEP* **05** (2002) 046, [arXiv:hep-ph/0112284](#).
- [138] B. Andersson, G. Gustafson, G. Ingelman, and T. Sjostrand, “Parton Fragmentation and String Dynamics,” *Phys. Rept.* **97** (1983) 31–145.
- [139] B. Andersson, *The Lund Model*. Cambridge Monographs on Particle Physics, Nuclear Physics and Cosmology. Cambridge University Press, 1998.
- [140] D. Amati and G. Veneziano, “Preconfinement as a Property of Perturbative QCD,” *Phys. Lett. B* **83** (1979) 87–92.
- [141] D. Yennie, S. Frautschi, and H. Suura, “The infrared divergence phenomena and high-energy processes,” *Annals of Physics* **13** no. 3, (1961) 379–452. <https://www.sciencedirect.com/science/article/pii/0003491661901518>.
- [142] M. Dobbs and J. B. Hansen, “The HepMC C++ Monte Carlo event record for High Energy Physics,” *Comput. Phys. Commun.* **134** (2001) 41–46.
- [143] **GEANT4** Collaboration, S. Agostinelli *et al.*, “GEANT4: A Simulation toolkit,” *Nucl. Instrum. Meth.* **A506** (2003) 250–303.
- [144] **ATLAS Collaboration** Collaboration, “The new Fast Calorimeter Simulation in ATLAS,” Tech. Rep. ATL-SOFT-PUB-2018-002, CERN, Geneva, Jul, 2018. <https://cds.cern.ch/record/2630434>.
- [145] K. Cranmer, “Practical Statistics for the LHC,” in *2011 European School of High-Energy Physics*, pp. 267–308. 2014. [arXiv:1503.07622 \[physics.data-an\]](#).
- [146] G. Cowan, K. Cranmer, E. Gross, and O. Vitells, “Asymptotic formulae for likelihood-based tests of new physics,” *Eur. Phys. J.* **C71** (2011) 1554, [arXiv:1007.1727 \[physics.data-an\]](#). [Erratum: *Eur. Phys. J.* C73,2501(2013)].
- [147] ATLAS Collaboration, “Reproduction searches for new physics with the ATLAS experiment through publication of full statistical likelihoods.” ATL-PHYS-PUB-2019-029, 2019. <https://cds.cern.ch/record/2684863>.
- [148] **ROOT Collaboration** Collaboration, K. Cranmer, G. Lewis, L. Moneta, A. Shibata, and W. Verkerke, “HistFactory: A tool for creating statistical models for use with RooFit and RooStats,” Tech. Rep. CERN-OPEN-2012-016, New York U., New York, Jan, 2012. <https://cds.cern.ch/record/1456844>.
- [149] W. Verkerke and D. P. Kirkby, “The RooFit toolkit for data modeling,” *eConf* **C0303241** (2003) MOLT007, [arXiv:physics/0306116 \[physics\]](#). [,186(2003)].
- [150] L. Moneta, K. Belasco, K. S. Cranmer, S. Kreiss, A. Lazzaro, D. Piparo, G. Schott, W. Verkerke, and M. Wolf, “The RooStats Project,” *PoS ACAT2010* (2010) 057, [arXiv:1009.1003 \[physics.data-an\]](#).



- [151] F. James and M. Roos, “MINUIT: a system for function minimization and analysis of the parameter errors and corrections,” *Comput. Phys. Commun.* **10** no. CERN-DD-75-20, (Jul, 1975) 343–367. 38 p. <https://cds.cern.ch/record/310399>.
- [152] R. Brun and F. Rademakers, “ROOT: An object oriented data analysis framework,” *Nucl. Instrum. Meth.* **A389** (1997) 81–86.
- [153] I. Antcheva *et al.*, “Root — a c++ framework for petabyte data storage, statistical analysis and visualization,” *Computer Physics Communications* **182** no. 6, (2011) 1384 – 1385. <http://www.sciencedirect.com/science/article/pii/S0010465511000701>.
- [154] M. Baak, G. J. Besjes, D. Côte, A. Koutsman, J. Lorenz, and D. Short, “HistFitter software framework for statistical data analysis,” *Eur. Phys. J.* **C75** (2015) 153, [arXiv:1410.1280](https://arxiv.org/abs/1410.1280) [hep-ex].
- [155] L. Heinrich, M. Feickert, G. Stark, and K. Cranmer, “pyhf: pure-python implementation of histfactory statistical models,” *Journal of Open Source Software* **6** no. 58, (2021) 2823. <https://doi.org/10.21105/joss.02823>.
- [156] L. Heinrich, M. Feickert, and G. Stark, “pyhf: v0.6.0.” <https://github.com/scikit-hep/pyhf>.
- [157] A. Paszke, S. Gross, F. Massa, A. Lerer, J. Bradbury, G. Chanan, T. Killeen, Z. Lin, N. Gimelshein, L. Antiga, A. Desmaison, A. Kopf, E. Yang, Z. DeVito, M. Raison, A. Tejani, S. Chilamkurthy, B. Steiner, L. Fang, J. Bai, and S. Chintala, “Pytorch: An imperative style, high-performance deep learning library,” in *Advances in Neural Information Processing Systems 32*, H. Wallach, H. Larochelle, A. Beygelzimer, F. d'Alché-Buc, E. Fox, and R. Garnett, eds., pp. 8024–8035. Curran Associates, Inc., 2019. <http://papers.neurips.cc/paper/9015-pytorch-an-imperative-style-high-performance-deep-learning-library.pdf>.
- [158] M. Abadi, A. Agarwal, P. Barham, E. Brevdo, Z. Chen, C. Citro, G. S. Corrado, A. Davis, J. Dean, M. Devin, S. Ghemawat, I. Goodfellow, A. Harp, G. Irving, M. Isard, Y. Jia, R. Jozefowicz, L. Kaiser, M. Kudlur, J. Levenberg, D. Mané, R. Monga, S. Moore, D. Murray, C. Olah, M. Schuster, J. Shlens, B. Steiner, I. Sutskever, K. Talwar, P. Tucker, V. Vanhoucke, V. Vasudevan, F. Viégas, O. Vinyals, P. Warden, M. Wattenberg, M. Wicke, Y. Yu, and X. Zheng, “TensorFlow: Large-scale machine learning on heterogeneous systems,” 2015. <https://www.tensorflow.org/>. Software available from tensorflow.org.
- [159] J. Bradbury, R. Frostig, P. Hawkins, M. J. Johnson, C. Leary, D. Maclaurin, and S. Wanderman-Milne, “JAX: composable transformations of Python+NumPy programs,” 2018. <http://github.com/google/jax>.
- [160] A. Wald, “Tests of statistical hypotheses concerning several parameters when the number of observations is large,” *Transactions of the American Mathematical Society* **54** no. 3, (1943) 426–482. <https://doi.org/10.1090/S0002-9947-1943-0012401-3>.
- [161] S. S. Wilks, “The large-sample distribution of the likelihood ratio for testing composite hypotheses,” *Ann. Math. Statist.* **9** no. 1, (03, 1938) 60–62. <https://doi.org/10.1214/aoms/1177732360>.
- [162] G. Cowan, “Statistics for Searches at the LHC,” in *69th Scottish Universities Summer School in Physics: LHC Physics*, pp. 321–355. 7, 2013. [arXiv:1307.2487](https://arxiv.org/abs/1307.2487) [hep-ex].

- [163] A. L. Read, “Presentation of search results: the  $CL_S$  technique,” *J. Phys. G* **28** (2002) 2693.
- [164] R. D. Cousins, J. T. Linnemann, and J. Tucker, “Evaluation of three methods for calculating statistical significance when incorporating a systematic uncertainty into a test of the background-only hypothesis for a Poisson process,” *Nucl. Instrum. Meth. A* **595** no. 2, (2008) 480, [arXiv:physics/0702156](#) [[physics.data-an](#)].
- [165] K. CRANMER, “Statistical challenges for searches for new physics at the lhc,” *Statistical Problems in Particle Physics, Astrophysics and Cosmology* (May, 2006) .  
[http://dx.doi.org/10.1142/9781860948985\\_0026](http://dx.doi.org/10.1142/9781860948985_0026).
- [166] ATLAS Collaboration, “Search for direct pair production of a chargino and a neutralino decaying to the 125 GeV Higgs boson in  $\sqrt{s} = 8$  TeV  $pp$  collisions with the ATLAS detector,” *Eur. Phys. J. C* **75** (2015) 208, [arXiv:1501.07110](#) [[hep-ex](#)].
- [167] ATLAS Collaboration, “Search for chargino and neutralino production in final states with a Higgs boson and missing transverse momentum at  $\sqrt{s} = 13$  TeV with the ATLAS detector,” *Phys. Rev. D* **100** (2019) 012006, [arXiv:1812.09432](#) [[hep-ex](#)].
- [168] CMS Collaboration, “Search for electroweak production of charginos and neutralinos in  $WH$  events in proton–proton collisions at  $\sqrt{s} = 13$  TeV,” *JHEP* **11** (2017) 029, [arXiv:1706.09933](#) [[hep-ex](#)].
- [169] ATLAS Collaboration, “Search for direct production of electroweakinos in final states with one lepton, missing transverse momentum and a Higgs boson decaying into two  $b$ -jets in  $pp$  collisions at  $\sqrt{s} = 13$  TeV with the ATLAS detector,” *Eur. Phys. J. C* **80** (2020) 691, [arXiv:1909.09226](#) [[hep-ex](#)].
- [170] ATLAS Collaboration, “Improvements in  $t\bar{t}$  modelling using NLO+PS Monte Carlo generators for Run 2.” ATL-PHYS-PUB-2018-009, 2018.  
<https://cds.cern.ch/record/2630327>.
- [171] ATLAS Collaboration, “Modelling of the  $t\bar{t}H$  and  $t\bar{t}V(V = W, Z)$  processes for  $\sqrt{s} = 13$  TeV ATLAS analyses.” ATL-PHYS-PUB-2016-005, 2016.  
<https://cds.cern.ch/record/2120826>.
- [172] ATLAS Collaboration, “ATLAS simulation of boson plus jets processes in Run 2.” ATL-PHYS-PUB-2017-006, 2017. <https://cds.cern.ch/record/2261937>.
- [173] ATLAS Collaboration, “Multi-Boson Simulation for 13 TeV ATLAS Analyses.” ATL-PHYS-PUB-2017-005, 2017. <https://cds.cern.ch/record/2261933>.
- [174] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H. S. Shao, T. Stelzer, P. Torrielli, and M. Zaro, “The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations,” *JHEP* **07** (2014) 079, [arXiv:1405.0301](#) [[hep-ph](#)].
- [175] R. Frederix and S. Frixione, “Merging meets matching in MC@NLO,” *JHEP* **12** (2012) 061, [arXiv:1209.6215](#) [[hep-ph](#)].
- [176] T. Sjöstrand, S. Ask, J. R. Christiansen, R. Corke, N. Desai, P. Ilten, S. Mrenna, S. Prestel, C. O. Rasmussen, and P. Z. Skands, “An Introduction to PYTHIA 8.2,” *Comput. Phys. Commun.* **191** (2015) 159–177, [arXiv:1410.3012](#) [[hep-ph](#)].



- [177] L. Lönnblad and S. Prestel, “Matching tree-level matrix elements with interleaved showers,” *JHEP* **03** (2012) 019, [arXiv:1109.4829 \[hep-ph\]](#).
- [178] R. D. Ball *et al.*, “Parton distributions with LHC data,” *Nucl. Phys. B* **867** (2013) 244, [arXiv:1207.1303 \[hep-ph\]](#).
- [179] ATLAS Collaboration, “ATLAS Pythia 8 tunes to 7 TeV data.” ATL-PHYS-PUB-2014-021, 2014. <https://cds.cern.ch/record/1966419>.
- [180] D. J. Lange, “The EvtGen particle decay simulation package,” *Nucl. Instrum. Meth. A* **462** (2001) 152.
- [181] ATLAS Collaboration, “The Pythia 8 A3 tune description of ATLAS minimum bias and inelastic measurements incorporating the Donnachie–Landshoff diffractive model.” ATL-PHYS-PUB-2016-017, 2016. <https://cds.cern.ch/record/2206965>.
- [182] B. Fuks, M. Klasen, D. R. Lamprea, and M. Rothering, “Precision predictions for electroweak superpartner production at hadron colliders with RESUMMINO,” *Eur. Phys. J. C* **73** (2013) 2480, [arXiv:1304.0790 \[hep-ph\]](#).
- [183] J. Fiaschi and M. Klasen, “Neutralino-chargino pair production at NLO+NLL with resummation-improved parton density functions for LHC Run II,” *Phys. Rev. D* **98** no. 5, (2018) 055014, [arXiv:1805.11322 \[hep-ph\]](#).
- [184] B. Fuks, M. Klasen, D. R. Lamprea, and M. Rothering, “Gaugino production in proton-proton collisions at a center-of-mass energy of 8 TeV,” *JHEP* **10** (2012) 081, [arXiv:1207.2159 \[hep-ph\]](#).
- [185] S. Alioli, P. Nason, C. Oleari, and E. Re, “A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX,” *JHEP* **06** (2010) 043, [arXiv:1002.2581 \[hep-ph\]](#).
- [186] S. Frixione, P. Nason, and G. Ridolfi, “A Positive-weight next-to-leading-order Monte Carlo for heavy flavour hadroproduction,” *JHEP* **09** (2007) 126, [arXiv:0707.3088 \[hep-ph\]](#).
- [187] P. Nason, “A New method for combining NLO QCD with shower Monte Carlo algorithms,” *JHEP* **11** (2004) 040, [arXiv:hep-ph/0409146](#).
- [188] E. Bothmann *et al.*, “Event generation with Sherpa 2.2,” *SciPost Phys.* **7** no. 3, (2019) 034, [arXiv:1905.09127 \[hep-ph\]](#).
- [189] S. Höche, F. Krauss, S. Schumann, and F. Siegert, “QCD matrix elements and truncated showers,” *JHEP* **05** (2009) 053, [arXiv:0903.1219 \[hep-ph\]](#).
- [190] S. Höche, F. Krauss, M. Schönherr, and F. Siegert, “QCD matrix elements + parton showers. The NLO case,” *JHEP* **04** (2013) 027, [arXiv:1207.5030 \[hep-ph\]](#).
- [191] NNPDF Collaboration, R. D. Ball *et al.*, “Parton distributions for the LHC run II,” *JHEP* **04** (2015) 040, [arXiv:1410.8849 \[hep-ph\]](#).
- [192] ATLAS Collaboration, “Example ATLAS tunes of PYTHIA8, PYTHIA6 and POWHEG to an observable sensitive to  $Z$  boson transverse momentum.” ATL-PHYS-PUB-2013-017, 2013. <https://cds.cern.ch/record/1629317>.

- [193] M. Czakon and A. Mitov, “Top++: A program for the calculation of the top-pair cross-section at hadron colliders,” *Comput. Phys. Commun.* **185** (2014) 2930, [arXiv:1112.5675 \[hep-ph\]](#).
- [194] M. Cacciari, M. Czakon, M. Mangano, A. Mitov, and P. Nason, “Top-pair production at hadron colliders with next-to-next-to-leading logarithmic soft-gluon resummation,” *Phys. Lett. B* **710** (2012) 612–622, [arXiv:1111.5869 \[hep-ph\]](#).
- [195] P. Kant, O. M. Kind, T. Kintscher, T. Lohse, T. Martini, S. Mölbitz, P. Rieck, and P. Uwer, “HatHor for single top-quark production: Updated predictions and uncertainty estimates for single top-quark production in hadronic collisions,” *Comput. Phys. Commun.* **191** (2015) 74–89, [arXiv:1406.4403 \[hep-ph\]](#).
- [196] N. Kidonakis, “Two-loop soft anomalous dimensions for single top quark associated production with a  $W^-$  or  $H^-$ ,” *Phys. Rev. D* **82** (2010) 054018, [arXiv:1005.4451 \[hep-ph\]](#).
- [197] J. M. Campbell and R. K. Ellis, “ $t\bar{t}W^{+-}$  production and decay at NLO,” *JHEP* **07** (2012) 052, [arXiv:1204.5678 \[hep-ph\]](#).
- [198] A. Lazopoulos, T. McElmurry, K. Melnikov, and F. Petriello, “Next-to-leading order QCD corrections to  $t\bar{t}Z$  production at the LHC,” *Phys. Lett. B* **666** (2008) 62–65, [arXiv:0804.2220 \[hep-ph\]](#).
- [199] R. Gavin, Y. Li, F. Petriello, and S. Quackenbush, “FEWZ 2.0: A code for hadronic  $Z$  production at next-to-next-to-leading order,” [arXiv:1011.3540 \[hep-ph\]](#).
- [200] **LHC Higgs Cross Section Working Group** Collaboration, D. de Florian *et al.*, “Handbook of LHC Higgs Cross Sections: 4. Deciphering the Nature of the Higgs Sector,” [arXiv:1610.07922 \[hep-ph\]](#).
- [201] ATLAS Collaboration, “Performance of the ATLAS track reconstruction algorithms in dense environments in LHC Run 2,” *Eur. Phys. J. C* **77** (2017) 673, [arXiv:1704.07983 \[hep-ex\]](#).
- [202] R. Frühwirth, “Application of Kalman filtering to track and vertex fitting,” *Nucl. Instrum. Methods Phys. Res., A* **262** no. HEPHY-PUB-503, (Jun, 1987) 444. 19 p. <https://cds.cern.ch/record/178627>.
- [203] T. Cornelissen, M. Elsing, I. Gavrilenko, W. Liebig, E. Moyse, and A. Salzburger, “The new ATLAS track reconstruction (NEWT),” *J. Phys.: Conf. Ser.* **119** (2008) 032014. <https://cds.cern.ch/record/1176900>.
- [204] ATLAS Collaboration, “Vertex Reconstruction Performance of the ATLAS Detector at  $\sqrt{s} = 13$  TeV.” ATL-PHYS-PUB-2015-026, 2015. <https://cds.cern.ch/record/2037717>.
- [205] ATLAS Collaboration, “Reconstruction of primary vertices at the ATLAS experiment in Run 1 proton–proton collisions at the LHC,” *Eur. Phys. J. C* **77** (2017) 332, [arXiv:1611.10235 \[hep-ex\]](#).
- [206] ATLAS Collaboration, “Topological cell clustering in the ATLAS calorimeters and its performance in LHC Run 1,” *Eur. Phys. J. C* **77** (2017) 490, [arXiv:1603.02934 \[hep-ex\]](#).

- [207] ATLAS Collaboration, “Electron and photon performance measurements with the ATLAS detector using the 2015–2017 LHC proton–proton collision data,” *JINST* **14** (2019) P12006, [arXiv:1908.00005 \[hep-ex\]](#).
- [208] ATLAS Collaboration, “Measurement of the photon identification efficiencies with the ATLAS detector using LHC Run 2 data collected in 2015 and 2016,” *Eur. Phys. J. C* **79** (2019) 205, [arXiv:1810.05087 \[hep-ex\]](#).
- [209] ATLAS Collaboration, “Electron and photon energy calibration with the ATLAS detector using 2015–2016 LHC proton–proton collision data,” *JINST* **14** (2019) P03017, [arXiv:1812.03848 \[hep-ex\]](#).
- [210] ATLAS Collaboration, “Electron reconstruction and identification in the ATLAS experiment using the 2015 and 2016 LHC proton–proton collision data at  $\sqrt{s} = 13$  TeV,” *Eur. Phys. J. C* **79** (2019) 639, [arXiv:1902.04655 \[hep-ex\]](#).
- [211] ATLAS Collaboration, “Muon reconstruction performance of the ATLAS detector in proton–proton collision data at  $\sqrt{s} = 13$  TeV,” *Eur. Phys. J. C* **76** (2016) 292, [arXiv:1603.05598 \[hep-ex\]](#).
- [212] ATLAS Collaboration, “Muon reconstruction and identification efficiency in ATLAS using the full Run 2  $pp$  collision data set at  $\sqrt{s} = 13$  TeV,” [arXiv:2012.00578 \[hep-ex\]](#).
- [213] M. Cacciari, G. P. Salam, and G. Soyez, “The anti- $k_t$  jet clustering algorithm,” *JHEP* **04** (2008) 063, [arXiv:0802.1189 \[hep-ph\]](#).
- [214] M. Cacciari, G. P. Salam, and G. Soyez, “FastJet user manual,” *Eur. Phys. J. C* **72** (2012) 1896, [arXiv:1111.6097 \[hep-ph\]](#).
- [215] M. Cacciari, “FastJet: A Code for fast  $k_t$  clustering, and more,” in *Deep inelastic scattering. Proceedings, 14th International Workshop, DIS 2006, Tsukuba, Japan, April 20–24, 2006*, pp. 487–490. 2006. [arXiv:hep-ph/0607071 \[hep-ph\]](#). [,125(2006)].
- [216] ATLAS Collaboration, G. Aad *et al.*, “Jet energy scale and resolution measured in proton–proton collisions at  $\sqrt{s} = 13$  TeV with the ATLAS detector,” [arXiv:2007.02645 \[hep-ex\]](#).
- [217] M. Cacciari and G. P. Salam, “Pileup subtraction using jet areas,” *Phys. Lett. B* **659** (2008) 119–126, [arXiv:0707.1378 \[hep-ph\]](#).
- [218] ATLAS Collaboration, “Jet energy measurement with the ATLAS detector in proton–proton collisions at  $\sqrt{s} = 7$  TeV,” *Eur. Phys. J. C* **73** (2013) 2304, [arXiv:1112.6426 \[hep-ex\]](#).
- [219] ATLAS Collaboration, “Determination of jet calibration and energy resolution in proton–proton collisions at  $\sqrt{s} = 8$  TeV using the ATLAS detector,” [arXiv:1910.04482 \[hep-ex\]](#).
- [220] ATLAS Collaboration, “Performance of pile-up mitigation techniques for jets in  $pp$  collisions at  $\sqrt{s} = 8$  TeV using the ATLAS detector,” *Eur. Phys. J. C* **76** (2016) 581, [arXiv:1510.03823 \[hep-ex\]](#).
- [221] ATLAS Collaboration, “Optimisation and performance studies of the ATLAS  $b$ -tagging algorithms for the 2017-18 LHC run.” ATL-PHYS-PUB-2017-013, 2017. <https://cds.cern.ch/record/2273281>.

- [222] ATLAS Collaboration, “ATLAS  $b$ -jet identification performance and efficiency measurement with  $t\bar{t}$  events in  $pp$  collisions at  $\sqrt{s} = 13$  TeV,” *Eur. Phys. J. C* **79** (2019) 970, [arXiv:1907.05120 \[hep-ex\]](#).
- [223] ATLAS Collaboration, “Measurements of  $b$ -jet tagging efficiency with the ATLAS detector using  $t\bar{t}$  events at  $\sqrt{s} = 13$  TeV,” *JHEP* **08** (2018) 089, [arXiv:1805.01845 \[hep-ex\]](#).
- [224] ATLAS Collaboration, “Performance of missing transverse momentum reconstruction with the ATLAS detector using proton–proton collisions at  $\sqrt{s} = 13$  TeV,” *Eur. Phys. J. C* **78** (2018) 903, [arXiv:1802.08168 \[hep-ex\]](#).
- [225] **ATLAS Collaboration** Collaboration, “ $E_{\text{T}}^{\text{miss}}$  performance in the ATLAS detector using 2015–2016 LHC p-p collisions,” Tech. Rep. ATLAS-CONF-2018-023, CERN, Geneva, Jun, 2018. <http://cds.cern.ch/record/2625233>.
- [226] D. Adams *et al.*, “Recommendations of the Physics Objects and Analysis Harmonisation Study Groups 2014,” Tech. Rep. ATL-PHYS-INT-2014-018, CERN, Geneva, Jul, 2014. <https://cds.cern.ch/record/1743654>.
- [227] M. Cacciari, G. P. Salam, and G. Soyez, “The Catchment Area of Jets,” *JHEP* **04** (2008) 005, [arXiv:0802.1188 \[hep-ph\]](#).
- [228] **UA1 Collaboration**, G. Arnison *et al.*, “Experimental Observation of Isolated Large Transverse Energy Electrons with Associated Missing Energy at  $\sqrt{s} = 540$  GeV,” *Phys. Lett. B* **122** (1983) 103–116.
- [229] **Aachen-Annecy-Birmingham-CERN-Helsinki-London(QMC)-Paris(CdF)-Riverside-Rome-Rutherford-Saclay(CEN)-Vienna** Collaboration, G. Arnison *et al.*, “Further evidence for charged intermediate vector bosons at the SPS collider,” *Phys. Lett. B* **129** no. CERN-EP-83-111, (Jun, 1985) 273–282. 17 p. <https://cds.cern.ch/record/163856>.
- [230] D. R. Tovey, “On measuring the masses of pair-produced semi-invisibly decaying particles at hadron colliders,” *JHEP* **04** (2008) 034, [arXiv:0802.2879 \[hep-ph\]](#).
- [231] G. Polesello and D. R. Tovey, “Supersymmetric particle mass measurement with the boost-corrected contranverse mass,” *JHEP* **03** (2010) 030, [arXiv:0910.0174 \[hep-ph\]](#).
- [232] **ATLAS Collaboration**, G. Aad *et al.*, “Performance of the missing transverse momentum triggers for the ATLAS detector during Run-2 data taking,” *JHEP* **08** (2020) 080, [arXiv:2005.09554 \[hep-ex\]](#).
- [233] **ATLAS Collaboration**, G. Aad *et al.*, “Performance of algorithms that reconstruct missing transverse momentum in  $\sqrt{s} = 8$  TeV proton-proton collisions in the ATLAS detector,” *Eur. Phys. J. C* **77** no. 4, (2017) 241, [arXiv:1609.09324 \[hep-ex\]](#).
- [234] ATLAS Collaboration, “ATLAS data quality operations and performance for 2015–2018 data-taking,” *JINST* **15** (2020) P04003, [arXiv:1911.04632 \[physics.ins-det\]](#).
- [235] ATLAS Collaboration, “Selection of jets produced in 13 TeV proton–proton collisions with the ATLAS detector.” ATLAS-CONF-2015-029, 2015. <https://cds.cern.ch/record/2037702>.
- [236] N. Hartmann, “ahoi.” <https://gitlab.com/nikoladze/ahoi>, 2018.

- [237] **ATLAS Collaboration**, “Object-based missing transverse momentum significance in the ATLAS detector,” Tech. Rep. ATLAS-CONF-2018-038, CERN, Geneva, Jul, 2018. <https://cds.cern.ch/record/2630948>.
- [238] A. Roodman, “Blind analysis in particle physics,” *eConf C030908* (2003) TUIT001, [arXiv:physics/0312102](https://arxiv.org/abs/physics/0312102).
- [239] ATLAS Collaboration, “A method for the construction of strongly reduced representations of ATLAS experimental uncertainties and the application thereof to the jet energy scale.” ATL-PHYS-PUB-2015-014, 2015. <https://cds.cern.ch/record/2037436>.
- [240] J. Bellm *et al.*, “Herwig 7.0/Herwig++ 3.0 release note,” *Eur. Phys. J. C* **76** no. 4, (2016) 196, [arXiv:1512.01178](https://arxiv.org/abs/1512.01178) [hep-ph].
- [241] ATLAS Collaboration, “Simulation of top-quark production for the ATLAS experiment at  $\sqrt{s} = 13$  TeV.” ATL-PHYS-PUB-2016-004, 2016. <https://cds.cern.ch/record/2120417>.
- [242] S. Frixione, E. Laenen, P. Motylinski, C. White, and B. R. Webber, “Single-top hadroproduction in association with a  $W$  boson,” *JHEP* **07** (2008) 029, [arXiv:0805.3067](https://arxiv.org/abs/0805.3067) [hep-ph].
- [243] **ATLAS Collaboration** Collaboration, “SUSY July 2020 Summary Plot Update,” Tech. Rep. ATL-PHYS-PUB-2020-020, CERN, Geneva, Jul, 2020. <http://cds.cern.ch/record/2725258>.
- [244] G. Apollinari, I. Béjar Alonso, O. Brüning, M. Lamont, and L. Rossi, *High-Luminosity Large Hadron Collider (HL-LHC): Preliminary Design Report*. CERN Yellow Reports: Monographs. CERN, Geneva, 2015. <https://cds.cern.ch/record/2116337>.
- [245] **LHC Reinterpretation Forum** Collaboration, W. Abdallah *et al.*, “Reinterpretation of LHC Results for New Physics: Status and Recommendations after Run 2,” *SciPost Phys.* **9** no. 2, (2020) 022, [arXiv:2003.07868](https://arxiv.org/abs/2003.07868) [hep-ph].
- [246] ATLAS Collaboration, “RECAST framework reinterpretation of an ATLAS Dark Matter Search constraining a model of a dark Higgs boson decaying to two  $b$ -quarks.” ATL-PHYS-PUB-2019-032, 2019. <https://cds.cern.ch/record/2686290>.
- [247] K. Cranmer and I. Yavin, “RECAST: Extending the Impact of Existing Analyses,” *JHEP* **04** (2011) 038, [arXiv:1010.2506](https://arxiv.org/abs/1010.2506) [hep-ex].
- [248] S. Ovin, X. Roubey, and V. Lemaitre, “DELPHES, a framework for fast simulation of a generic collider experiment,” [arXiv:0903.2225](https://arxiv.org/abs/0903.2225) [hep-ph].
- [249] A. Buckley, J. Butterworth, D. Grellscheid, H. Hoeth, L. Lonnblad, J. Monk, H. Schulz, and F. Siegert, “Rivet user manual,” *Comput. Phys. Commun.* **184** (2013) 2803–2819, [arXiv:1003.0694](https://arxiv.org/abs/1003.0694) [hep-ph].
- [250] A. Buckley, D. Kar, and K. Nordström, “Fast simulation of detector effects in Rivet,” *SciPost Phys.* **8** (2020) 025, [arXiv:1910.01637](https://arxiv.org/abs/1910.01637) [hep-ph].
- [251] D. Dercks, N. Desai, J. S. Kim, K. Rolbiecki, J. Tattersall, and T. Weber, “CheckMATE 2: From the model to the limit,” *Comput. Phys. Commun.* **221** (2017) 383–418, [arXiv:1611.09856](https://arxiv.org/abs/1611.09856) [hep-ph].
- [252] M. Drees, H. Dreiner, D. Schmeier, J. Tattersall, and J. S. Kim, “CheckMATE: Confronting your Favourite New Physics Model with LHC Data,” *Comput. Phys. Commun.* **187** (2015) 227–265, [arXiv:1312.2591](https://arxiv.org/abs/1312.2591) [hep-ph].

- [253] E. Conte, B. Fuks, and G. Serret, “MadAnalysis 5, A User-Friendly Framework for Collider Phenomenology,” *Comput. Phys. Commun.* **184** (2013) 222–256, [arXiv:1206.1599 \[hep-ph\]](#).
- [254] E. Maguire, L. Heinrich, and G. Watt, “HEPData: a repository for high energy physics data,” *J. Phys. Conf. Ser.* **898** no. 10, (2017) 102006, [arXiv:1704.05473 \[hep-ex\]](#).
- [255] **ATLAS** Collaboration, “Simpleanalysis,” <https://gitlab.cern.ch/atlas-sa/simple-analysis>, 2021.
- [256] S. Kraml, S. Kulkarni, U. Laa, A. Lessa, W. Magerl, D. Proschofsky-Spindler, and W. Waltenberger, “SModelS: a tool for interpreting simplified-model results from the LHC and its application to supersymmetry,” *Eur. Phys. J. C* **74** (2014) 2868, [arXiv:1312.4175 \[hep-ph\]](#).
- [257] F. Ambrogio, S. Kraml, S. Kulkarni, U. Laa, A. Lessa, V. Magerl, J. Sonneveld, M. Traub, and W. Waltenberger, “SModelS v1.1 user manual: Improving simplified model constraints with efficiency maps,” *Comput. Phys. Commun.* **227** (2018) 72–98, [arXiv:1701.06586 \[hep-ph\]](#).
- [258] **ATLAS** Collaboration, “Search for direct production of electroweakinos in final states with one lepton, missing transverse momentum and a higgs boson decaying into two  $b$ -jets in  $pp$  collisions at  $\sqrt{s} = 13$  tev with the atlas detector,” 2021. <https://www.hepdata.net/record/ins1755298?version=4>.
- [259] **LHC Reinterpretation Forum** Collaboration, W. Abdallah *et al.*, “Reinterpretation of LHC Results for New Physics: Status and Recommendations after Run 2,” *SciPost Phys.* **9** no. 2, (2020) 022, [arXiv:2003.07868 \[hep-ph\]](#).
- [260] **ATLAS** Collaboration, “1lbb-likelihoods-hepdata.tar.gz,” 2020. <https://www.hepdata.net/record/resource/1408476?view=true>.
- [261] G. Alguero, S. Kraml, and W. Waltenberger, “A SModelS interface for pyhf likelihoods,” [arXiv:2009.01809 \[hep-ph\]](#).
- [262] M. D. Goodsell, “Implementation of the ATLAS-SUSY-2019-08 analysis in the MadAnalysis 5 framework (electroweakinos with a Higgs decay into a  $b\bar{b}$  pair, one lepton and missing transverse energy;  $139\text{ fb}^{-1}$ ),” *Mod. Phys. Lett. A* **36** no. 01, (2021) 2141006.
- [263] J. Y. Araz *et al.*, “Proceedings of the second MadAnalysis 5 workshop on LHC recasting in Korea,” *Mod. Phys. Lett. A* **36** no. 01, (2021) 2102001, [arXiv:2101.02245 \[hep-ph\]](#).
- [264] M. Feickert, L. Heinrich, G. Stark, and B. Galewsky, “Distributed statistical inference with pyhf enabled through funcX,” in *25th International Conference on Computing in High-Energy and Nuclear Physics*. 3, 2021. [arXiv:2103.02182 \[cs.DC\]](#).
- [265] R. Chard, Y. Babuji, Z. Li, T. Skluzacek, A. Woodard, B. Blaiszik, I. Foster, and K. Chard, “funcx: A federated function serving fabric for science,” ACM, Jun, 2020. <http://dx.doi.org/10.1145/3369583.3392683>.
- [266] D. Merkel, “Docker: Lightweight linux containers for consistent development and deployment,” *Linux J.* **2014** no. 239, (Mar., 2014) .
- [267] S. Binet and B. Couturier, “docker & HEP: Containerization of applications for development, distribution and preservation,” *J. Phys.: Conf. Ser.* **664** no. 2, (2015) 022007. 8 p. <https://cds.cern.ch/record/2134524>.



- [268] K. Cranmer and L. Heinrich, “Yadage and Packtivity - analysis preservation using parametrized workflows,” *J. Phys. Conf. Ser.* **898** no. 10, (2017) 102019, [arXiv:1706.01878 \[physics.data-an\]](#).
- [269] ATLAS Collaboration, “Summary of the ATLAS experiment’s sensitivity to supersymmetry after LHC Run 1 — interpreted in the phenomenological MSSM,” *JHEP* **10** (2015) 134, [arXiv:1508.06608 \[hep-ex\]](#).
- [270] ATLAS Collaboration, “Searches for electroweak production of supersymmetric particles with compressed mass spectra in  $\sqrt{s} = 13$  TeV  $pp$  collisions with the ATLAS detector,” *Phys. Rev. D* **101** (2020) 052005, [arXiv:1911.12606 \[hep-ex\]](#).
- [271] ATLAS Collaboration, “Search for electroweak production of charginos and sleptons decaying into final states with two leptons and missing transverse momentum in  $\sqrt{s} = 13$  TeV  $pp$  collisions using the ATLAS detector,” *Eur. Phys. J. C* **80** (2020) 123, [arXiv:1908.08215 \[hep-ex\]](#).
- [272] ATLAS Collaboration, “Search for direct stau production in events with two hadronic  $\tau$ -leptons in  $\sqrt{s} = 13$  TeV  $pp$  collisions with the ATLAS detector,” *Phys. Rev. D* **101** (2020) 032009, [arXiv:1911.06660 \[hep-ex\]](#).
- [273] ATLAS Collaboration, “Search for bottom-squark pair production with the ATLAS detector in final states containing Higgs bosons,  $b$ -jets and missing transverse momentum,” *JHEP* **12** (2019) 060, [arXiv:1908.03122 \[hep-ex\]](#).
- [274] W. Porod, “SPheno, a program for calculating supersymmetric spectra, SUSY particle decays and SUSY particle production at  $e^+e^-$  colliders,” *Comput. Phys. Commun.* **153** (2003) 275–315, [arXiv:hep-ph/0301101](#).
- [275] W. Porod and F. Staub, “SPheno 3.1: Extensions including flavour, CP-phases and models beyond the MSSM,” *Comput. Phys. Commun.* **183** (2012) 2458–2469, [arXiv:1104.1573 \[hep-ph\]](#).
- [276] S. Heinemeyer, W. Hollik, and G. Weiglein, “FeynHiggs: A Program for the calculation of the masses of the neutral CP even Higgs bosons in the MSSM,” *Comput. Phys. Commun.* **124** (2000) 76–89, [arXiv:hep-ph/9812320](#).
- [277] H. Bahl, T. Hahn, S. Heinemeyer, W. Hollik, S. Paßehr, H. Rzehak, and G. Weiglein, “Precision calculations in the MSSM Higgs-boson sector with FeynHiggs 2.14,” *Comput. Phys. Commun.* **249** (2020) 107099, [arXiv:1811.09073 \[hep-ph\]](#).
- [278] T. Hahn, S. Heinemeyer, W. Hollik, H. Rzehak, and G. Weiglein, “High-Precision Predictions for the Light CP -Even Higgs Boson Mass of the Minimal Supersymmetric Standard Model,” *Phys. Rev. Lett.* **112** no. 14, (2014) 141801, [arXiv:1312.4937 \[hep-ph\]](#).
- [279] B. C. Allanach, “SOFTSUSY: a program for calculating supersymmetric spectra,” *Comput. Phys. Commun.* **143** (2002) 305–331, [arXiv:hep-ph/0104145 \[hep-ph\]](#).
- [280] G. Belanger, F. Boudjema, A. Pukhov, and A. Semenov, “MicrOMEGAs 2.0: A Program to calculate the relic density of dark matter in a generic model,” *Comput. Phys. Commun.* **176** (2007) 367–382, [arXiv:hep-ph/0607059](#).
- [281] G. Belanger, F. Boudjema, A. Pukhov, and A. Semenov, “micrOMEGAs: A Tool for dark matter studies,” *Nuovo Cim. C* **033N2** (2010) 111–116, [arXiv:1005.4133 \[hep-ph\]](#).

- [282] P. Athron, M. Bach, H. G. Fargnoli, C. Gnendiger, R. Greifenhagen, J.-h. Park, S. Paßehr, D. Stöckinger, H. Stöckinger-Kim, and A. Voigt, “GM2Calc: Precise MSSM prediction for  $(g - 2)$  of the muon,” *Eur. Phys. J. C* **76** no. 2, (2016) 62, [arXiv:1510.08071 \[hep-ph\]](#).
- [283] F. Mahmoudi, “SuperIso v2.3: A Program for calculating flavor physics observables in Supersymmetry,” *Comput. Phys. Commun.* **180** (2009) 1579–1613, [arXiv:0808.3144 \[hep-ph\]](#).
- [284] W. Beenakker, R. Hopker, and M. Spira, “PROSPINO: A Program for the Production of Supersymmetric Particles in Next-to-leading Order QCD,” Tech. Rep. hep-ph/9611232, Nov, 1996. <https://cds.cern.ch/record/314229>. 12 pages, latex, no figures, Complete postscript file and FORTRAN source codes available from <http://wwwcn.cern.ch/mspira/prospino/>.
- [285] W. Beenakker, M. Klasen, M. Kramer, T. Plehn, M. Spira, and P. M. Zerwas, “The Production of charginos / neutralinos and sleptons at hadron colliders,” *Phys. Rev. Lett.* **83** (1999) 3780–3783, [arXiv:hep-ph/9906298](#). [Erratum: Phys.Rev.Lett. 100, 029901 (2008)].