

1 INTRODUCTION

1.1 Events [2.2]

- (1) The result u_1 of a random trial is called **outcome**.
- (2) The set of all possible outcomes is called **sample space** Ω .
- (3) A collection of outcomes A is called **event**.

Thus an event is a subset of the sample space (or possibly the whole sample space).

The events A_1, \dots, A_n are said to be **mutually exclusive** (or **mutually disjoint**) if all pairs A_i, A_j are exclusive; that is, if it is impossible that two or more of these events take place at the same time.

1.2 Sample spaces [2.2]

If the number of outcomes is finite or denumerably infinite, Ω is said to be a **discrete sample space**. More particularly, if the number is finite, Ω is said to be a **finite sample space**.

If the number of outcomes is neither finite nor denumerably infinite, Ω is said to be a **continuous sample space**.

1.3 Kolmogorov's System of Axioms [2.3]

The following axioms for the probabilities $P(\cdot)$ should be fulfilled:

- A1.** If A is any event, then $0 \leq P(A) \leq 1$.
- A2.** If Ω is the entire sample space, then $P(\Omega) = 1$.
- A3.** ("Addition Formula"). If A and B are mutually exclusive events, then $P(A \cup B) = P(A) + P(B)$.

The sample space Ω and the probabilities $P(\cdot)$ together constitute a **probability space**.

1.4 Results from Kolmogorov's system [2.3]

Complement Theorem

$$P(A^*) = 1 - P(A) \quad (1)$$

$$P(\emptyset) = 0 \quad (2)$$

Addition Theorem

$$P(A \cup B) = P(A) + P(B) + P(A \cap B)$$

$$P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(A \cap C) - P(B \cap C) + P(A \cap B \cap C) \quad (3)$$

Boole's Inequality

$$P(A \cup B) \leq P(A) + P(B)$$

$$P(A \cup B \cup C) \leq P(A) + P(B \cup C) \leq P(A) + P(B) + P(C) \quad (4)$$

1.5 Probabilities in Discrete Sample Spaces [2.4]

$$P(u_i) = 1/m \quad (i = 1, 2, \dots, m) \quad (5)$$

If (5), we have a **uniform probability distribution**.

An event A with g outcomes, has g **favourable cases**. That means there are m **possible cases** in Ω . We then determine $P(A)$ from the addition formula by adding the probabilities m of the g outcomes. Hence,

$$P(A) = g/m \quad (6)$$

1.6 Urn model [2.4]

Let's examine the probability of drawing exactly k white balls from an urn containing a white and b black balls, n balls are drawn.

If one ball at a time is drawn at random without returning it to the urn, the drawings are said to be made **without replacement**. The probability is then:

$$\frac{\binom{a}{k} \binom{b}{n-k}}{\binom{a+b}{n}} \quad (7)$$

If one ball at a time is drawn at random and is returned before the next ball is drawn, the drawings are said to be made **with replacement**. The probability is then:

$$\binom{n}{k} p^k q^{n-k} \quad (8)$$

where


$$p = \frac{a}{a+b}, \quad q = 1 - p = \frac{b}{a+b} \quad (9)$$

1.7 Conditional Probability [2.5]

$$P(B|A) = \frac{P(A \cap B)}{P(A)} \quad (10)$$


$$P(A \cap B) = P(A)P(B|A) = P(B)P(A|B)$$

is called the **conditional probability of B given A**.



From this we can derive the **Total Probability Theorem**. If the events H_1, H_2, \dots, H_n are mutually exclusive, have positive probabilities, and together fill Ω completely, any event A satisfies the formula:

$$P(A) = \sum_{i=1}^n P(H_i)P(A|H_i) \quad (11)$$



And **Bayes' Theorem**:

$$P(H_i|A) = \frac{P(H_i)P(A|H_i)}{\sum_{j=1}^n P(H_j)P(A|H_j)} \quad (12)$$

1.8 Independent Events [2.6]

$$P(A \cap B) = P(A)P(B), \quad (13)$$

If (13), then A and B are said to be **independent events**. This can be extended to three events. Consequence:

If the events A_1, A_2, \dots, A_n are independent and $P(A_i) = p_i$, then the probability that at least one of them occurs is equal to

$$1 - (1 - p_1)(1 - p_2) \cdots (1 - p_n) \quad (14)$$

If the events A_i are independent and each one of them occurs with probability p , then the probability that at least one of them occurs is equal to $1 - (1 - p)^n$.

Two trials as **independent**, if the result of one does not affect, or is not affected by, the result of the other. An important special case is provided by **repeated trials**.

1.9 Theorems in Combinatorics [2.7]

Drawing n elements from N :

	with replacement	without replacement
order	N^n	$N(N - 1) \cdots (N - n + 1)$
no order	$\binom{N + n - 1}{n}$	$\binom{N}{n}$

The **multiplication principle** states: If choice 1 can be performed in a_1 ways and choice 2 in a_2 ways, then there are $a_1 a_2$ ways to perform both choices. In the case of three choices, the number of ways is $a_1 a_2 a_3$, and so on.